

2 **Effect of selected soil amendments on the**  
3 **incidences of *Ralstonia solanacearum* isolates**  
4 **in greenhouse on selected solanaceous crops**

5  
6  
7

8 **ABSTRACT**

**Aims:** The aim of this study was to establish the effect of selected soil amendments on the incidences of *Ralstonia solanacearum* isolates in greenhouse on selected solanaceous crops.

**Study design:** The study was laid out as randomized complete block design (RCBD) in split pot arrangement for two seasons in the greenhouse.

**Place and Duration of Study:** The experiment was carried out in Kenyatta University situated in Kiambu County about 20 km from Nairobi city along Nairobi-Thika road between July, 2017- September, 2017 and between November, 2017- January, 2018.

**Methodology:** The three host crops of interest (potatoes, tomatoes and capsicum) were inoculated with prepared pure bacterial isolates; 18 (2T-Kiambu-Low Land), 71(2A-Nyeri-Low Land), 67 (2A-Nyeri-High Land), 83 (2T-Kirinyaga-Highland) and MX (18/71/67/83). The experiment had a total of 450 pots having a total area of 6.93 M<sup>2</sup>. The treatments were Chalim™, Super-hydro-grow polymer + Metham sodium, Metham sodium, Metham sodium & Orange peel, Super-hydro-grow polymer, Brassica tissues, Chalim™ + Super-hydro-grow polymer, Brassica tissue + Orange peel, Metham sodium + Super-hydro-grow polymer and Control (no amendments).

**Results:** There was significant differences ( $P \leq 0.05$ ) in the bacterial wilt incidences in selected solanaceous crops between control and all the soil amendments used in season 1 and 2. Brassica tissue + Super-hydro-grow polymer was superior in reducing bacterial wilt incidences in selected solanaceous crops in all the *R. solanacearum* isolates from Kenyan highlands and lowlands both in season 1 and 2.

**Conclusion:** Organic and inorganic soil amendments could serve as a viable control of bacterial wilt in solanaceous crops caused by *R. solanacearum* in the greenhouse.

9  
10 **Key:** Bacterial wilt, incidences, *Ralstonia solanacearum*, solanaceous crops

11  
12 **1. INTRODUCTION**

13  
14 The increasing global population coupled with the challenges of environmental degradation and an increasingly variable  
15 climate have created a world-wide need for improved food security [1, 2]. Bacterial wilt disease caused by *Ralstonia*  
16 *solanacearum* is one of the most important constraints in production of vegetables in the tropical and sub-tropical regions  
17 [3]. The pathogen *R. solanacearum* is widespread in tropical, sub-tropical and warm temperate regions and infects more  
18 than 450 plant species in 54 families [4, 5]. *R. solanacearum*, the causal agent of bacterial wilt disease, is considered one  
19 of the most destructive bacterial pathogens due to its lethality, unusually wide host range, persistence and broad  
20 geographical distribution Wei *et al.* [6]. *R. solanacearum* is a diverse species that differs in host range, geographical

21 distribution, pathogenicity and biochemical and physiological properties [7]. This pathogenic species has been divided into  
22 five races based on host range [8] and six biovars based on metabolic profiles, related to the ability to metabolize three  
23 sugar alcohols and three disaccharides. These globally dispersed and heterogeneous strains cause loss of productivity of  
24 many crops, which have major socio-economic impacts [9]

25  
26 Tim *et al.* [10] and Joshi *et al.* [11] reported that bacterial wilt disease is affected by environmental condition like soil  
27 temperature, soil moisture, soil type (which influences soil moisture and microbial populations).The bacterium enters host  
28 plant roots from the soil and colonizes the xylem vessels in the vascular system [12]. Infected plants suffer yellowing,  
29 stunting and wilting, and often die rapidly [13]. Symptoms of *R. solanacearum* on tomato include wilting and necrosis as  
30 well as vascular browning [14]. Typically, stem and tuber cross-sections ooze whitish bacterial exudates [15]. Ramesh *et al.*  
31 [16] further reported that *R. solanacearum* infection could spread through contaminated water and weeds in the  
32 Solanaceae family. *Ralstonia solanacearum* can be disseminated by farm implements, pollinator insects in banana,  
33 irrigation water, infested soil, plant debris, latently infected vegetative propagation materials (such as *Pelargonium*  
34 cuttings, potato tubers and banana corms) and through roots damaged by nematodes [17, 18]. Latent infection is  
35 widespread and has been identified in several asymptomatic host plants including tomato, geranium, squash, and potato  
36 [19, 20]. An incidence of 55 % and 25 % has been recorded on chili and potato crops respectively from the major chili and  
37 potato producing regions of Ethiopia [21]. Assefa *et al.*[22] (2015) reported that, in Ethiopia percentage wilt incidence of  
38 bacterial wilt was as high as 63 % on potato, 55 % on tomato and 100 % on pepper. Singh *et al.* [23] observed in a study  
39 in India that crop losses of up to 90% was reported in the greenhouse compared to losses of 25-60 % reported for open  
40 field tomato

41  
42  
43 Bacterial wilt control in various pathosystems has been possible through use of a combination of diverse methods such  
44 as, host resistance, biofumigation, fertiliser application, soil solarisation, biological control, chemical control, other cultural  
45 practices and integrated  
46 disease management schemes [24,25]. However, traditional control measures are not always effective in the control of  
47 bacterial wilt, such as the application of bactericides, disease-resistant cultivars and crop rotation [26], thus making it  
48 compelling to find a potential soil-borne disease control method to reduce economic loss. Synthetic chemicals have been  
49 used for many years to control agricultural biological agents, however, considerable problems have arisen from the  
50 continued application of these chemicals, including development of resistance by the pathogen, high cost, residual effect  
51 on soil, pollution of the environment and hazard from handling toxic compounds [27]. The use of soil amendments (SAs) is  
52 a widespread means to control soil-borne disease. It has been reported that an SA, composed of urea and calcium oxide  
53 (CaO), is effective to control the bacterial wilt of tomato by affecting the pH and nitrite accumulation in the field [28, 29].  
54 Calcium carbonate (CaCO<sub>3</sub>) could not only serve as a soil amendment to change soil pH but also increase soil Ca<sup>2+</sup>  
55 content. Polymers are widely used for many applications in agriculture: to combat viruses and other crop pathogens, and  
56 functionalized polymers are employed to increase the efficiency of pesticides and herbicides, allowing the application of  
57 lower doses and thus indirectly protecting the environment [30]. Amendments provide energy and nutrients to soil,  
58 drastically changing the environment for the growth and survival of crops and microorganisms [31]. In the current study,  
59 the aim of this study was to establish the effect of selected soil amendments on the incidences of *R. solanacearum*  
60 isolates in greenhouse on selected solanaceous crops. This current study will help to reduce bacteria wilt in very  
61 important selected solanaceous crops by using organic and inorganic amendments.

## 62 63 64 **2. MATERIAL AND METHODS**

### 65 66 **2.1 Study Area**

67  
68 The experiment was carried out in Kenyatta University situated in Kiambu County about 20 km from Nairobi city along  
69 Nairobi-Thika road. The county enjoys a warm climate with temperatures ranging between 12°C and 18.7°C. The rainfall  
70 aggregate for the county is 1000 mm each year. Its geographical coordinates are 1° 10' 0" South, 36° 50' 0" East. Low  
71 fertility soils are mainly found in the middle zone and the eastern part of the county which form part of the semi-arid areas.  
72 The soils in the midland zone are dissected and are easily eroded. The soils are sandy or clay and can support drought  
73 resistant crops such as soya beans and sunflower as well as ranching. The elevation of the main campus is 1720 meters  
74 above sea level (ASL) [32].

### 75 76 **2.2 Experimental Design and Treatments**

77 The experiment was carried out between July, 2017- September, 2017 and between November, 2017- January, 2018 and  
78 was replicated three times for the two seasons. The experiment was laid out in randomized complete block design  
79 (RCBD) in split plot arrangement in the greenhouse. Potatoes, tomatoes and capsicum were planted in pots each with a

radius of 0.07 M (area 0.015 M<sup>2</sup>). The experiment had a total of 450 pots having a total area of 6.93 M<sup>2</sup>. The treatments were Chalim™, Super-hydro-grow polymer and Metham sodium, Metham sodium, Metham sodium +Orange peel, Super-hydro-grow polymer, Control, Brassica tissue, Chalim™ + Super-hydro-grow polymer, Brassica tissue + Orange peel and Metham sodium + Super-hydro-grow polymer. The three host crops of interest (potatoes, tomatoes and capsicum) were inoculated with prepared pure bacterial isolates; 18 (2T-Kiambu-Low Land), 71(2A-Nyeri-Low Land), 67 (2A-Nyeri-High Land), 83 (2T-Kirinyaga-Highland) and MX (18/71/67/83). All agronomic practices including, watering, fertilization, weeds, pests and disease control were well managed.

## 2.3 Preparation of Pot Soil Amendments

Fresh leaves of cabbage plant residues were finely chopped and incorporated into the soil at the rate of 0.01kg per pot of an area of 0.015 M<sup>2</sup>. The inoculated soil was thoroughly mixed with the finely chopped cabbage plant residue, ensuring that all the residues were well incorporated in the soil pot. Metham sodium, a chemical fumigant was applied in pot of an area of 0.015 M<sup>2</sup> at the rate of 3.08 ml per pot. This was the positive control. Chalim™ effect was assessed in the inoculated pots after application at the rate of 0.3 g per pot of an area of 0.015 M<sup>2</sup>. Super-hydro-grow polymer was applied in plot of 0.015 M<sup>2</sup> at the rate of 0.0003 ml per pot using knap-sack sprayer. Combination of Chalim™ + Super-hydro-grow polymer was applied at the rate of 0.3 g per pot and 0.0003 ml per the same pot of an area of 0.015 M<sup>2</sup> respectively. Metham sodium + Super-hydro-grow polymer was applied in a pot of an area of 0.015 M<sup>2</sup> at the rate of 3.08 ml per pot and 0.0003 ml per the same pot respectively.

Metham sodium +Orange peel treatment was applied in a pot of an area of 0.015 M<sup>2</sup> at the rate of 3.08 ml per pot and Orange peel rate of 0.01kg per the same pot respectively. Brassica tissue + Orange peel treatment were applied at a rate 0.01kg per pot of an area of 0.015 M<sup>2</sup> and Orange peel at a rate of 0.01kg per the same pot respectively.

## 2.4 Greenhouse inoculation

The positively identified potato tubers and stem tubers of capsicum and tomato were used to isolate *R. solanacearum*. The five pure bacterial isolates were 18 (2T-Kiambu-Low Land), 71(2A-Nyeri-Low Land), 67 (2A-Nyeri-High Land),83 (2T-Kirinyaga-Highland) and MX (2T-L/2A-L/2A-H/2T-H(18/71/67/83)). The bacteria was purified, multiplied and then harvested to make a composite bacterial inoculum. Pure bacterial was harvested (30 plates per plant sample) into a 10 L of sterile distilled water to make composite bacterial inoculate to be sprayed in 450 pots. The experimental plant pots was performed in glasshouse where the 450 pots were inoculated each with 10 mL of 3.05 x 10<sup>9</sup> cfu/mL of *R. solanacearum* isolates 18, 71, 67, 83 and MX. Metham sodium, a known fumigant was used as a positive control. Randomized complete split plot design was used in the pot layout.

## 2.5 Data Collection and Analysis

Three sample crops (tomato, capsicum and potato) were used. The plants were rated weekly, each Wednesday for bacterial wilt disease incidence from the 18<sup>th</sup> day after planting where wilted plants were uprooted upon total foliage wilt and recorded though only the incidence at 4<sup>th</sup>, 7<sup>th</sup> and 10<sup>th</sup> weeks after planting (WAP) was considered for evaluation. For proper key diagnostic identification of *R. solanacearum* in the field and to distinguish bacterial wilt from vascular wilts caused by fungal pathogens, bacterial wilt symptoms was identified by visual observation of typical bacterial wilt disease symptoms such as wilting, vascular discoloration, bacterial streaming in glass of water and browning of the vascular bundles of the tuber. Milky white strands containing bacteria and extracellular polysaccharide was oozed out from the cut ends of the xylem. The diseased samples were brought to the laboratory and subjected aseptically for detection and confirmation of *Ralstonia solanacearum*. The assessment of the incidences of *R. solanacearum* was estimated using the formulas as adapted from Mwaniki *et al.*, (2016).

$$\text{Disease incidence} = \frac{\text{Total asymptomatic Plants}}{\text{Total Plants per area}} \times 100\%$$

## 2.8 Data analysis

Data that was obtained from soil amendments effect of incidences of *R. solanacearum* on selected solanaceous plants was statistically analyzed by statistical package for social sciences (SPSS) software for Windows, ver. 23 (SPSS, IBM, USA). Chi-square was done to measure the strength of associations between variables. A p-value of <0.05 was considered to be statistically significant.

### 3. RESULTS AND DISCUSSION

#### 3.1 Influence of Organic and Inorganic Soil Amendments on Disease Incidence on Potatoes

The results of incidences of bacterial wilt on potatoes grown under greenhouse for season 1 and 2 are shown by figure 1 and 2 respectively.





Figure 1. Greenhouse Bacterial wilt incidence in potatoes season 1




<sup>1</sup> BT-Brassicae Tissue, BT+OP -Brassica tissue+Orange peel, BT+SHG- Brassicae Tissue+ Super-hydro-grow polymer, CM- Chalim™, CM+OP- Chalim™+ Orange peel, CM+SHG- Chalim™+ Super-hydro-grow polymer,MS- Metham sodium, Ms+OP- Metham sodium+ Orange peel, MS+SHG- Metham sodium+ Super-hydro-grow polymer : 18 (2T-Kiambu-Low Land), 71(2A-Nyeri-Low Land), 67 (2A-Nyeri-High Land),83 (2T-Kirinyaga-Highland) and MX (2T-L/2A-L/2A-H/2T-H(18/71/67/83).

**Figure 2. Greenhouse Bacterial wilt incidence in potatoes season 2**

Significant differences ( $P \leq 0.05$ ) were revealed in the bacterial wilt incidences in potatoes between control and all the soil amendments used in season 1 and 2 in the five *R. solanacearum* isolate from Kenyan highlands and lowlands. These results indicate the suppressive effect of organic and inorganic treatments used in this study. The Brassica tissue + Super-hydro-grow polymer was superior in reducing bacterial wilt incidences in potatoes in all the *R. solanacearum* isolates from Kenyan highlands and lowlands. Brassica species produce glucosinolates which are nematocidal and biocidal. The biocidal action of isothiocyanates produced by Brassica tissue and their potential to manage and suppress phytopathogens has been reported by Brown & Morra [33] and Matthiessen & Kirkegaard [34]. Significant differences ( $P \leq 0.05$ ) were found in the bacterial wilt incidences in potatoes between Brassica tissue alone and Brassica tissue + Orange peel and Brassica tissue + Super-hydro-grow polymer soil amendments used in both season 1 and 2. Brassica tissue + orange peels, Brassica tissue + Super-hydro-grow polymer and Chalim™ + Super-hydro-grow polymer had a synergetic effect in eliminating *R. solanacearum* in potatoes as opposed to Brassica tissue and Chalim™ used individually as seen in Figure 1 and 2. Chalim (CaCO<sub>3</sub>) in soil reduced bacterial wilt incidence, which was accord with the results of Heyman *et al.* [35]. The population of *R. solanacearum* was reduced significantly in the soil with CaCO<sub>3</sub> treatment, which suggested that the effect of CaCO<sub>3</sub> on *R. solanacearum* was mainly related to the role of Ca<sup>2+</sup> in the greenhouse experiment which increased activity of peroxidase (POD) and polyphenoloxidase (PPO) thus reducing incidences of *R. solanacearum*. Isolate 71 was resistant to Brassica tissue + Orange peel soil amendment causing bacterial wilt while the other isolates were susceptible hence no incidence of bacterial wilt in potatoes to the same treatment in season 2. Variations in the incidence of bacterial wilt are attributable to the diversity of *R. solanacearum* strains, variations in soil types in different agro ecological zones. There was no significant differences ( $P \leq 0.05$ ) revealed in the bacterial wilt incidences among the isolate in the greenhouse control experiment in potatoes. There was decline in the incidences of bacterial wilt in the second season with subsequent treatment with the same soil ammendement except for greenhouse control experiement.


**3.2 Influence of Organic and Inorganic Soil Amendments on Disease Incidence on Capsicum**

The results of incidences of bacterial wilt on capsicum grown under gree house for season 1 and 2 are shown by figure 3 and 4 respectively.



<sup>2</sup> BT-Brassicae Tissue, BT+OP -Brassica tissue+Orange peel, BT+SHG- Brassicae Tissue+ Super-hydro-grow polymer, CM- Chalim™, CM+OP- Chalim™+ Orange peel, CM+SHG- Chalim™+ Super-hydro-grow polymer,MS- Metham sodium, Ms+OP- Metham sodium+ Orange peel, MS+SHG- Metham sodium+ Super-hydro-grow polymer: 18 (2T-Kiambu-Low Land), 71(2A-Nyeri-Low Land), 67 (2A-Nyeri-High Land),83 (2T-Kirinyaga-Highland) and MX (2T-L/2A-L/2A-H/2T-H(18/71/67/83).

187  
188 **Figure 3. Greenhouse Bacterial wilt incidence in Capsicum season 1**  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201



202  
203  
204 **Figure 3. Greenhouse Bacterial wilt incidence in Capsicum season 2**  
205

206 There was no significant differences ( $P \leq 0.05$ ) revealed in the bacterial wilt incidences between control and Metham sodium + Super-hydro-grow polymer and Metham sodium + Orange peel in the greenhouse control experiment in  
207 capsicum in both season 1 and 2. The Brassica tissue + Super-hydro-grow polymer was superior in reducing bacterial wilt  
208 incidences in capsicum in all the *R. solanacearum* isolates from Kenyan highlands and lowlands both in season 1 and 2.  
209 Soil amendment in season 2 with Brassica tissue + Orange peel, Brassica tissue + Super-hydro-grow polymer and  
210 Chalim™ + Super-hydro-grow polymer recorded zero incidence of bacterial wilt in capsicum in all the isolates. The use of  
211 Brassica tissue plant residue in the control of bacterial wilt has been conducted and reported to be effective in a study by  
212 [36]. The use of plant tissue in controlling of bacterial wilt has been reported to be eco-friendly relative to the use of  
213 inorganic chemicals as well as being readily available to the resource-poor farmers [37]. Combination of Chalim™ +  
214 Super-hydro-grow polymer significantly ( $P \leq 0.05$ ) reduced bacterial wilt incidences in capsicum as opposed to use of  
215 Chalim™ alone in the two seasons. Metham sodium soil amendment significantly ( $P \leq 0.05$ ) reduced bacterial incidences  
216 as opposed to use a combination of Metham sodium + Super-hydro-grow polymer and Metham sodium + Orange peel in  
217 the two seasons.  
218  
219  
220

### 221 3.3 Influence of Organic and Inorganic Soil Amendments on Disease Incidence on Tomatoes

222 The results of incidences of bacterial wilt on tomatoes grown under greenhouse for season 1 and 2 are shown by figure 5  
223 and 6 respectively.  
224  
225

<sup>3</sup> BT-Brassicae Tissue, BT+OP -Brassica tissue+Orange peel, BT+SHG- Brassicae Tissue+ Super-hydro-grow polymer, CM- Chalim™, CM+OP- Chalim™+ Orange peel, CM+SHG- Chalim™+ Super-hydro-grow polymer, MS- Metham sodium, Ms+OP- Metham sodium+





Figure 5. Greenhouse Bacterial wilt incidence in Tomatoes season 1

4

Orange peel, MS+SHG- Metham sodium+ Super-hydro-grow polymer: 18 (2T-Kiambu-Low Land), 71(2A-Nyeri-Low Land), 67 (2A-Nyeri-High Land),83 (2T-Kirinyaga-Highland) and MX (2T-L/2A-L/2A-H/2T-H(18/71/67/83).

<sup>4</sup> BT-Brassicae Tissue, BT+OP -Brassica tissue+Orange peel, BT+SHG- Brassicae Tissue+ Super-hydro-grow polymer, CM- Chalim™, CM+OP- Chalim™+ Orange peel, CM+SHG- Chalim™+ Super-hydro-grow polymer,MS- Metham sodium, Ms+OP- Metham sodium+ Orange peel, MS+SHG- Metham sodium+ Super-hydro-grow polymer : 18 (2T-Kiambu-Low Land), 71(2A-Nyeri-Low Land), 67 (2A-Nyeri-High Land),83 (2T-Kirinyaga-Highland) and MX (2T-L/2A-L/2A-H/2T-H(18/71/67/83).



**Figure 6. Greenhouse Bacterial wilt incidence in Tomatoes season 2**

Significant differences ( $P \leq 0.05$ ) were revealed in the bacterial wilt incidences in Tomatoes between control and all the soil amendments used in season 1 and 2 except in season 1 where Metham sodium + Orange peel and control did not have significant ( $P \leq 0.05$ ) difference as shown on figure 5 and 6. The Brassica tissue + Super-hydro-grow polymer was superior in reducing bacterial wilt incidences in tomatoes in all the *R. solanacearum* isolates from Kenyan highlands and lowlands both in season 1 and 2. Brassicaceous materials have been reported to have allelopathic effects as well as biofumigation effects to soil biota that includes plant parasitic nematodes [36, 37, 38, 39]. The production of biofumigation products including isothiocyanates (ITCs) that has an active ingredient related to that of Metham sodium and dazomet have been reported to be highly toxic to pests and pathogens. Kim *et al.* [40] reported that, Bacterial wilt of tomatoes caused by *R. solanacearum* is a devastating disease that limits the production of tomato in Korea. Tomato plants are grown worldwide in the field or greenhouse [41]. Isolate 71 was resistant to Brassica tissue + Orange peel soil amendment causing bacterial wilt while the other isolates were susceptible hence no incidence of bacterial wilt in tomatoes to the same treatment in season 2. There was significant difference ( $P \leq 0.05$ ) in bacterial wilt incidence in tomatoes for soil amendment between Brassica tissue soil amendment alone compared to a combination of Brassica tissue + Orange peel and Brassica tissue + Super-hydro-grow polymer in both season 1 and 2. Previous studies have shown that essential oils in citrus, protopine and corydaline alkaloids, lactons, polyacetylene, acyclic sesquiterpenes, hypericin and pseudohypericin compounds are effective toward various bacteria including *R. solanacearum* [42] (Maruti *et al.*, 2011).

In this study, all representative isolates from Kenyan highland and lowland used were pathogenic to tomato seedlings in the greenhouse to varying degrees. Singh *et al.* [23] showed that the micro-climate inside the greenhouse (Temperatures of 28°C-30°C, 80%-90% RH and wet soil) favors rapid pathogen multiplication and disease. Various findings have reported *Ralstonia* strains as differing in their virulence [43, 44]. Morais *et al.* [45] reported that information on the pathogenicity and molecular variability of *Ralstonia* strains will improve our knowledge on the epidemiology and ecology of these pathogens. This is particularly true with respect to latency, survival and aggressiveness of each strain. Bacterial wilt caused by *R. solanacearum* is one of the major diseases of tomato and the disease causes concern for tomato production because it can drastically reduce tomato up to 90% [46, 47].

Chalim soil amendment in tomatoes minimally reduced bacterial wilt between the two seasons. Meanwhile, higher  $\text{Ca}^{2+}$  content in tobacco was associated with less disease, which was agreement with the results of Sugimoto *et al.* [48]; the

<sup>5</sup> BT-Brassicacae Tissue, BT+OP -Brassica tissue+Orange peel, BT+SHG- Brassicae Tissue+ Super-hydro-grow polymer, CM- Chalim<sup>TM</sup>, CM+OP- Chalim<sup>TM</sup>+ Orange peel, CM+SHG- Chalim<sup>TM</sup>+ Super-hydro-grow polymer,MS- Metham sodium,MS+OP- Metham sodium+ Orange peel, MS+SHG- Metham sodium+ Super-hydro-grow polymer: 18 (2T-Kiambu-Low Land), 71(2A-Nyeri-Low Land), 67 (2A-Nyeri-High Land),83 (2T-Kirinyaga-Highland) and MX (2T-L/2A-L/2A-H/2T-H(18/71/67/83).



mechanism may be related to the increased activity of peroxidase (POD) and polyphenoloxidase (PPO) as reported by Jiang *et al.* [49] that the severity of tomato wilt can be reduced by increasing activity of POD and PPO in tomato with the increased calcium concentration in tomato tissues which concurs with our current findings. Mondal *et al.* [50] also found that Disease incidence in tomato crop was higher compared to other solanaceous crops like brinjal, chilli, capsicum and potato.

#### 4. CONCLUSION

In conclusion, our findings showed that organic and inorganic soil amendments could serve as a viable control of bacterial wilt in solanaceous crops caused by *R. solanacearum* in the greenhouse. Brassica tissue + Super-hydro-grow polymer was superior in reducing bacterial wilt incidences in selected solanaceous crops in all the *R. solanacearum* isolates from Kenyan highlands and lowlands both in season 1 and 2. The data presented in this study substantiate the findings that, various *R. solanacearum* isolates from both the Kenyan highland and lowland are causing bacteria wilt disease in various important solanaceous crops grown in the country.

#### COMPETING INTERESTS

The authors have no competing interests to declare

#### REFERENCES

1. Beddington J, Asaduzzaman M, Fernandez A, Clark M, Guillou M, Jahn M, Erda L, Mamo T, Van Bo N, Nobre CA, Scholes R, Sharma R, Wakhungu J. Achieving food security in the face of climate change: Summary for policy makers from the Commission on Sustainable Agriculture and Climate Change. CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS). Copenhagen, Denmark. 2011. Available online at: [www.ccafs.cgiar.org/commission](http://www.ccafs.cgiar.org/commission).
2. Godfray HCJ, Pretty J, Thomas SM, Warham EJ, Beddington JR. Linking policy on climate and food Science. 2011; 331, 1013–1014. (doi:10.1126/science.1202899)
3. Devendra K C, Aundy K, Sajad N. In-vitro Evaluation of Arabidopsis thaliana Ecotypes against *Ralstonia solanacearum* Race4, Int. J. Curr. Microbiol. App. Sci. 2017; 6(5):575-579
4. Guidot A, Coupat B, Fall S, Prior P and Bertolla F. Horizontal gene transfer between *Ralstonia solanacearum* strains detected by comparative genomic hybridization on microarrays. ISME J. 3, 2009; 549–5 doi: 10.1038/ismej.2009.14
5. Prior P, Ailloud F, Dalsing BL, Remenant B, Sanchez B, Allen C. Genomic and proteomic evidence supporting the division of the plant pathogen *Ralstonia solanacearum* into three species. BMC Genom. 2016; 17, 90.
6. Wei Y, Moreno CC, Gongora TJ, Wang K, Sang Y, Duran R, Macho AP. The *Ralstonia solanacearum* csp22 peptide, but not flagellin-derived peptides, is perceived by plants from the Solanaceae family. Journal of Plant Biotechnology. 2018; 1–14.
7. Sikirou R, Zocli B, Paret ML, Deberdt P, Coranson-Beaudu R, Huat J., et al. (2015). First report of bacterial wilt of Gboma (*Solanum macrocarpon*) caused by *Ralstonia solanacearum* in Benin. *Plant Dis.* 11, 1640–1640. doi: 10.1094/PDIS-02-15-0213-PDN
8. Singh D, Sinha S, Yadav DK, Chaudhary G. Detection of *Ralstonia solanacearum* from asymptomatic tomato plants, irrigation water, and soil through non-selective enrichment medium with hrp gene-based bio-PCR. Curr Microbiol. 2014; 69: 127- 134.

- 325 9. Jiang Y, Li B, Liu P, Liao F, Weng Q, Chen Q. First report of bacterial wilt caused by *Ralstonia solanacearum* on fig  
326 trees in China. *For. Pathol.* 2016; 46 256–258. 10.1111/efp.12267  
327
- 328 10. Tim M, Prakash P, Carlos L. North Florida Research And Education Center (Nfrec), Quincy; Prakash Pradhanang  
329 Post-Doctoral associate, Nfrec, Quincy, FL 32351; Carlos A. Lopes, Researcher, Embrapa Hortaliças, Brasília 70359-  
330 970, Brasil. 2008.  
331
- 332 11. Joshi M, Srivastava R, Sharma AK and Prakash A. Screening of Resistant Varieties and Antagonistic *Fusarium*  
333 *oxysporum* for Biocontrol of *Fusarium* Wilt of Chilli. *J Plant Pathol Microb.* 2012; 3:134. doi:10.4172/2157-  
334 7471.1000134  
335
- 336 12. Muthoni J, Hussein S, Melis R. Management of bacterial wilt *Ralstonia solanacearum* Yabuuchi et al., 1995 of  
337 Potatoes: Opportunity for Host Resistance in Kenya *Journal of Agricultural Science*; Vol. 4, No. 9; ISSN 1916-9752 E-  
338 ISSN. 2012; 1916-9760.  
339
- 340 13. Meng F. The virulence factors of the bacterial wilt pathogen *Ralstonia solanacearum*. *J. Plant Pathol. Microbiol.* 2013;  
341 4, 3.  
342
- 343 14. Swanson JK, Yao J, Tans-Kersten J, Allen C, Behavior of *Ralstonia solanacearum* race 3 biovar 2 during latent and  
344 active infection of geranium. *Phytopathology.* 2005; 95, 136– 43.  
345
- 346 15. Genin S, Boucher C. *Ralstonia solanacearum*: secrets of a major pathogen unveiled by analysis of its genome.  
347 *Molecular Plant Pathology.* 2002; 3:111-118.  
348
- 349 16. Ramesh R, Achari GA and Gaitonde S. Genetic diversity of *Ralstonia solanacearum* infecting solanaceous vegetables  
350 from India reveals the existence of unknown or newer sequevars of Phylotype I strains. *Eur. J. Plant Pathol.* 2014;  
351 140: 543-562.  
352
- 353 17. Williamson L, Nakaho K, Hudelson B and Allen C. *Ralstonia solanacearum* race 3, biovar 2 strains isolated from  
354 geranium are pathogenic on potato. *Plant Disease.* 2002; 86:987-991.  
355
- 356 18. Kim SH, Olson TN, *Ralstonia solanacearum* race 3, biovar 2, the causal agent of brown rot of potato, identified in  
357 geraniums in Pennsylvania, Delaware, and Connecticut  
358 (Abstract). *Plant Disease.* 2003; 87, 450  
359
- 360 19. Cruz APZ, Ferreira V, Pianzola MJ, Siri MI, Coll N, Valls M. A novel, sensitive method to evaluate potato germplasm  
361 for bacterial wilt resistance using a luminescent *Ralstonia solanacearum* reporter strain. *Mol. Plant Microbe Interact.*  
362 2014; 27 277–285. 10.1094/MPMI  
363
- 364 20. Vrisman CM, Deblais L, Rajashekara G, and Miller SA. Differential colonization dynamics of cucurbit hosts by *Erwinia*  
365 *tracheiphila*. *Phytopathology.* 2016; 106:684-692. <https://doi.org/10.1094/PHYTO-11-15-0289-R> [Link](#), [ISI](#), [Google](#)  
366 [Scholar](#)  
367
- 368 21. Bekele B, Abate E, Asefa A. Dickinson M, Incidence of potato viruses and bacterial wilt disease in the west Amhara  
369 sub-region of Ethiopia. *Journal of Plant Pathology.* 2011.93(1):149-157.  
370 <http://sipav.org/main/jpp/index.php/jpp/article/view/285/151>  
371
- 372 22. Assefa M, Dawit W, Lencho A and Hunduma T. Assessment of wilt intensity and identification of causal fungal and  
373 bacterial pathogens on hot pepper (*Capsicum annum* L.) in BakoTibbe and Nonno districts of west Shewa zone,  
374 Ethiopia. *International Journal of Phytopathology.* 2015; 4 (1), pp.21-28..  
375
- 376 23. Singh, DR, Kumar K and Birah A. Eco-friendly management modules for bacterial wilt (*Ralstonia solanacearum*)  
377 of tomato for protected cultivation in a tropical island ecosystem. *Biological Agriculture and Horticulture:*  
378 *An International Journal for Sustainable Production Systems.* 2014a.  
379
- 380 24. Getachew A, Chemed F, Seid A, Wydra K. Effects of soil amendment on bacterial wilt caused by *Ralstonia*  
381 *solanacearum* and tomato yields in Ethiopia. *J Plant Prot Res.* 2011; 51(1):72–76  
382

- 383 25. Singh AK, Singh RK, Singh AK, Singh VK, Rawat SS, Mehta KS, Kumar A, Gupta MK, Thakur S. Bio-mulching for  
384 ginger crop management: traditional ecological knowledge led adaptation under rainfed agroecosystems. *Indian J*  
385 *Tradit Know.* 2013; 13(1):111–122.
- 386
- 387 26. Liu Y, Shi J, Feng Y, Yang X, Li X, Shen Q. Tobacco bacterial wilt can be biologically controlled by the application of  
388 antagonistic strains in combination with organic fertilizer. *Biol. Fertil. Soils.* 2013; 49:447–464.
- 389
- 390 27. Tijjani A, Bashir KA, Mohammed I, Muhammad A, Gambo A and Habu M. Biopesticides for pests control: a review.  
391 *Journal of Biopesticides and Agriculture.* 2016; 3 (1): 6 – 13
- 392
- 393 28. Michel VV, & Mew TW. Effect of soil amendment on the survival of *Ralstonia solanacearum* in different soils.  
394 *Phytopathology.* 1998; 88, 300–305.
- 395
- 396 29. Michel VV, Wang JF, Midmore DJ, & Hartman GL. Effects of intercropping and soil amendment with urea and calcium  
397 oxide on the incidence of bacterial wilt of tomato and survival of soil-borne *Pseudomonas solanacearum* in Taiwan.  
398 *Plant Pathology.* 1997; 46, 600–610.
- 399
- 400 30. Manjunatha SB, Biradar DP, Aladakatti YR. Nanotechnology and its applications in agriculture: A review. *Journal of*  
401 *Farm Science.* 2016; 29(1), 1-13.
- 402
- 403 31. Lazarovits G, Tenuta M, Conn KL. Organic amendments as a disease control strategy for soilborne diseases of high-  
404 value agricultural crops. *Australasian Plant Pathology*, 2001; 30(2), 111-117.
- 405
- 406 32. Jaetzold R, Schmidt H, Hornetz B, Shisanya C. Central Kenya. Agroecological Zones and subzones. Ministry of  
407 Agriculture, Farm Management Hand book of Kenya Vol. II. Natural conditions and Farm Management information  
408 2nd Edition Part B, Central Province: 2006; 434-438
- 409
- 410 33. Brown PD and Morra MJ. Control of soil-borne plant pests using glucosinolate containing plants. *Advances in*  
411 *Agronomy.* 1997; 61, pp. 167-231.
- 412
- 413 34. Matthiessen JN and Kirkegaard JA. Biofumigation for managing soil-borne pests - progress, pitfalls and  
414 prospects. In: Zalucki, M., Drew, R., and White, G. eds. Proceedings of the 6th Australasian Applied Entomology  
415 Research Conference. University of Queensland. 1998; 1, pp. 364- 372.
- 416
- 417 35. Heyman F, Lindahl B, Persson L, Wikstrom M and Stenlid J.. Calcium concentrations of soil affect suppressiveness  
418 against *Aphanomyces* root rot of pea. *Soil Biol. Biochem.* 2007; 39: 2222-2229.
- 419
- 420 36. Gruver LS, Weil RR, Zasada IA, Sardanelli S and Momena B. Brassicaceous and rye cover crops altered free-living  
421 soil nematode community composition. *Applied Soil Ecology.* 2010; 45, pp. 1-12.
- 422
- 423 37. Hartman GL, Hong WF and Hayward AC. Potential of biological and chemical control of bacterial wilt. In: Hartman GL  
424 and Hayward AC. eds. Bacterial wilt. Proceedings of ACIAR conference, October 28-31, 1992, Kaohsiung, Taiwan.  
425 ACIAR Proceedings No. 45. Brisbane, Australia, Watson Ferguson and Company. 1993; pp. 322-326.
- 426
- 427 38. Bailey KL and Lazarovits G. Suppressing soil borne diseases with residue management and organic  
428 amendments. *Soil and Tillage.* 2003; 72(2), pp. 169-180.
- 429
- 430 39. Schonfeld J, Gelsomin A, van Overbeek LS, Goris-sen A, Smalla K and van Elsas JD. Effects of compost  
431 addition and stimulated solarisation on the fate of *Ralstonia solanacearum* biovar 2 and indigenous bacteria in soil.  
432 *FMESMicrobiology Ecology.* 2003; 43, pp. 63-74.
- 433
- 434 40. Kim SG, Hur OS, Ro NY, Ko HC, Rhee JH, Sung JS, Ryu KY, Lee SY, Baek HJ. Evaluation of resistance to  
435 *Ralstonia solanacearum* in tomato genetic resources at seedling stage. *Plant Pathology Journal.* 2016; 32(1): 58-64.
- 436
- 437 41. Peralta IE, Spooner DM. History, origin and early cultivation of tomato (Solanaceae). In: Razdan MK, Mattoo AK, eds.  
438 *Genetic improvement of solanaceous crops.* Vol. 2. Tomato. Enfield, NH: Science Publishers. 2007; 1–27.
- 439
- 440 42. Maruti J. Dhanavade CB. Jalkute KD, Sonawane K, Jai SG. Study antimicrobial activity of lemon (Citrus lemon L)  
441 peel extract. *British Journal of Pharmacology and Toxicology.* 2011; 2 (3):119-122

- 442 43. Li Y, Feng J, Liu H, Wang L, Hsiang T, Li X, Huang J. Genetic diversity and pathogenicity of *Ralstonia solanacearum*  
443 causing tobacco bacterial wilt in China. *Plant Dis.* 2016; 100, 1288–1296.  
444
- 445 44. Rodrigues L, Destefano S, da Silva M, Costa G, Maringoni AC. Characterization of *Ralstonia solanacearum* strains  
446 from Brazil using molecular methods and pathogenicity tests. *J. Plant Pathol.* 2012; 94, 505–516.  
447
- 448 45. Morais TP, Lopes CA, Tebaldi ND, Luz JMQ. Occurrence and diversity of *Ralstonia solanacearum* populations in  
449 Brazil. *Biosci. J.* 2015; 31, 1722–1737.  
450
- 451 46. Fujiwara K, Aoyama C, Takano M, Shinohara M. Suppression of *Ralstonia solanacearum* bacterial wilt disease by an  
452 organic hydroponic system. *J. Gen. Plant Pathology.* 2012; 78: 217–220.  
453
- 454 47. Aslam MN, Mukhtar T, Hussain MA, Raheel M. Assessment of resistance to bacterial wilt incited by *Ralstonia*  
455 *solanacearum* in tomato germplasm. *Journal of Plant Disease Protection.* 2017; 124: 585–590.  
456
- 457 48. Sugimoto T, Watanabe K, Yoshida S, Aino M, Furiki M, Shiono M. Field application of calcium to reduce phytophthora  
458 stem rot of soybean, and calcium distribution in plants. *Plant Dis.* 2010; 94, 812–819. doi: 10.1094/PDIS-94-7-0812  
459
- 460 49. Jiang JF, Li JG, Dong YH. Effect of calcium nutrition on resistance of tomato against bacterial wilt induced by  
461 *Ralstonia solanacearum*. *Eur J Plant Pathol.* 2013; 136: 547–555.  
462
- 463 50. Mondal B, Bhattacharya I and Khatua DC. Crop and weed host of *Ralstonia solanacearum* in West Bengal. *Journal of*  
464 *Crop and Weed.* 2011; 7(2): 195-199.  
465  
466  
467  
468  
469