

**Evaluation of the Ornamental Potential of Safflower (*Carthamus tinctorius* L.)**

**ABSTRACT**

**Aims:** to evaluate the ornamental potential of two safflower genotypes (*Carthamus tinctorius* L.): ICA 73, ICA 193, grown under protected environment.

**Place and Duration of Study:** Department of Agronomy of Federal Rural University of Pernambuco, between March and May 2017.

**Methodology:** The methodology addressed evaluated the performance of the two genotypes, through three experiments. The first experiment was carried out in a completely randomized design, being evaluated: plant height; stem diameter; leaf dentin; margin of the spinescent leaves, number of branches, number of heads, margin of the spinescent bracts, flowering, and flower production. The second one was conducted in a randomized complete block design in a factorial scheme, and the following variables were evaluated: plant height; stem diameter; number of branches; number of heads; and flower production. Finally, the third experiment used a completely randomized design in a factorial scheme and evaluated the variables: number of flowers; number of flower buds; and post-harvest durability. The analysis of variance was performed using the F test at 5% of probability and, afterwards, the regression or comparison analysis of averages by the Tukey test at 5% of probability.

**Results:** The ICA 73 genotype showed plants with high flower production and the ICA 193 exhibited plants with weak or moderate margin spinescent leaves and bracts, besides good uniformity of the anthesis of the flowers.

**Conclusion:** Both accessions showed ornamental potential, presenting precocity, beauty and durability of the flowers. The density of a plant was the most favorable for potting and for cutting stem, the cut-off point for semi-open flowers was the best for obtaining quality stems.

*Keywords: Floriculture. Precocity. Flowers. Potted Plant. Cutting Stems.*

**1. INTRODUCTION**

Floriculture is a segment that stands out in the world, with significant growth, constituting an important source of income for several countries and promoting the development of productive poles. In this context, production is mainly intended for export to large consumer centers located in the European Union, the United States of America and more recently Asia and the Middle East [1,2].

In Brazil, in contrast to other developing countries, this sector has grown towards the intern market [3]. With annual growth of around 8% a year, growing exports and significant increase in domestic consumption, the floriculture in the country became one of the most prominent segments in the agribusiness market, moving around R \$ 6.7 billion in 2016 [4].

Most of the market is supplied by plants from few states in the Southeast of the country [5], while other important Brazilian regions, where new floriculture poles emerge, end up with development difficulties, despite their natural aptitude for this sector [6].

29

30 Evidently, the floriculture consists of a dynamic sector with a constant search for new  
31 products that meet the new trends. In this sense, safflower has great potential to offer news  
32 for this market. The safflower (*Carthamus tinctorius*) belongs to the Asteraceae family, it is  
33 an herbaceous annual plant, self-pollinated and capable of developing into various  
34 edaphoclimatic conditions [7,8], tolerating low water availability and high temperatures [9].

35

36 The use of this herbaceous plant covers many possibilities and extends to practically every  
37 part of the plant. Its seeds are the first most valued and exploited part in the market, being  
38 present in food products [10], cosmetics and also in the composition of drugs [11]. Safflower  
39 oil may also be intended for biofuel production [12], and the resulting bagasse in this  
40 process, used in animal feed supplements [13]. The second part with the most prominence  
41 concerns the flowers, which allow the extraction of two dyes from their petals, a soluble and  
42 another hydrophilic [14], and have ornamental potential and can be used fresh or dried for  
43 this purpose [15].

44

45 The variability of the color, size and arrangement of the florets that the safflower possesses,  
46 make the species very attractive to the ornamental market, being able to be used as  
47 ornamental plants, cutting stems and confection of bouquets, with potential for planting in  
48 gardens or pots and trade while fresh or when dry [16]. In Europe, the use of this  
49 herbaceous plant in the flower market is not only common, but also important, with specific  
50 cultivars for this purpose [17].

51

52 Safflower cultivation in Brazil is still very limited and is restricted to basically attending to  
53 some scientific research, moreover usually it covers only the production of oil, thus not  
54 exploiting the potential of ornamental culture. However, considering the importance that the  
55 floriculture is taking in Brazilian agribusiness, the search and insertion of new products to  
56 expand and meet market demand becomes an indisputable necessity and, in this context,  
57 investing in the potential presented by this species concerns a very promising strategy.

58

59 The objective of this work was to evaluate the ornamental potential of two safflower  
60 genotypes cultivated in a protected environment, to plant pot, determining the best density,  
61 and cutting stem, defining the best cutting point.

62

## 63 **2. MATERIAL AND METHODS**

64

65 The safflower accesses (*Carthamus tinctorius*) ICA 73 and 193 used in the study were  
66 imported by the Institute of Agricultural Sciences (ICA) in agreement with the Federal  
67 University of Minas Gerais (UFMG) of germplasm banks of India and Ethiopia, which were  
68 later transferred to the Plant Breeding Program of the Federal University of Pernambuco  
69 (UFRPE) to carry out this work.

70

71 Three experiments were carried out under greenhouse conditions in the Agronomy  
72 Department of the Federal Rural University of Pernambuco - UFRPE, Recife - PE, whose  
73 geographical coordinates are 8°10'52"S latitude, 34°54'47" longitude and 2m altitude.

74

75 Seeding was done manually in pots with 5L capacity, filled with commercial Basaplant™  
76 substrate, the depth of approximately three centimeters [18]. Irrigations were performed  
77 manually and daily.

78

### 79 **2.1 First Experiment (Characterization of Access)**

80

81 The experimental design adopted was completely randomized, with twenty repetitions. The  
82 treatments were composed of the two safflower genotypes. twenty vases were used for each  
83 access (ICA 73 and ICA 193), where each vase represented an experimental unit, totaling  
84 40 parcels.

85  
86 The Emergency Velocity Index (EVI) was calculated according to the formula of [19]:

87  $EVI = \frac{\sum(E_n/Nn)}{n}$ ; Where: n = the number of normal seedlings recorded in the count "n";  
88 Nn = number of days of sowing until the count "n". For this, the number of emerged plants,  
89 with two open cotyledon leaves, was registered until the ninth day after sowing. The  
90 percentage of germination was calculated after stabilization of the emergency, considering  
91 the final number of emerged plants.

92  
93 The evaluations considered all the experimental plots, being carried out sixty days after  
94 sowing (DAS), based on the following characteristics: Plant height (PH (cm)) - performed  
95 with ruler and corresponding to the measurement of the soil to the apex of the plant; Stem  
96 Diameter (SD (cm)) - measured with a digital pachymeter in the base of the stem; Leaf  
97 Dentin (LD) - classified by scale of notes: absent or weak (0); moderate (5); strong (10) [20]  
98 (Figure 1); Margin of the Spinescent Leaves (MSL) - graded by note scale: absent or weak  
99 (0); moderate (5); strong (10) [20] (Figure 2); Number of Branches per plant (NB) - obtained  
100 by counting (Figure 3); Number of Heads (NH) - obtained by counting heads (*capitulum*);  
101 Margin of the Spinescent Bracts (MSB) - graded by grading scale: absent or weak (0);  
102 moderate (5); strong (10) [20] (Figure 4); Flowering (FI) - number of days from sowing to  
103 beginning of flowering; Flower Production (FP) - obtained by counting open inflorescences.

104



105  
106

Figure 1. Leaf Dentin of Safflower: (A) Weak; (B) Moderate e (C) Strong.



107  
108  
109  
110

Figure 2. Margin of the Spinescent Leaves of Safflower: (A) Weak; (B) Moderate; (C) Strong.





111  
112  
113

**Figure 3. Safflower plants: (A) Little branched; (B) Very branched.**



114  
115  
116

**Figure 4. Margin of the Spinescent Leaves of Safflower: (A) Strong; (B) Moderate; (C) Absent.**

## 117 **2.2 Second Experiment (Conduction Pruning)**

118

119 The experimental design was the one of randomized blocks, in a 4 x 2 factorial scheme,  
120 combining four plant densities per vase and two safflower accessions, distributed in 4  
121 blocks. Each block was composed of eight vases, wherein each vase corresponded to an  
122 experimental unit, contemplating all the possibilities of combinations between the treatment  
123 factors, totalizing 32 experimental plots. The densities were evaluated referring to: four;  
124 three; two; and one plant per vase.

125

126 After completing the phase of rosette 30 DAS a pruning was done in all the plants, through a  
127 single bevel cut on the central stem. Subsequently, 30 days after the procedure, the  
128 following characteristics were evaluated: Plant height (PH (cm)); Stem Diameter (SD (cm));  
129 Number of branches per plant (NB); Number of heads (NH); Flower production (FP).

130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151

### 2.3 Third Experiment: (Post-Harvest Characterization)

The experimental design was a completely randomized design, in a 3 x 2 factorial scheme, combining three cutting points of the stems and two safflower accesses, using 4 repetitions. Twelve vases were used for each genotype, wherein each vase received two seeds and corresponded to one experimental unit, the cut-off points of the stems were: closed heads, semi-open heads, and open heads.

Through a bevel cut, removal of the central section of all the plants was carried out, when they were approximately 1 cm in diameter. This procedure, commonly called pinch, is used to stimulate the development of lateral heads.

The cutting of the stems was done according to the respective treatments: Open inflorescences, above 70% of the open florets; semi-open, 30 to 40% of open florets; and closed, 5 to 15% of the open florets (Figure 5). The stems were cut in the basal portion, about 3 cm from the base of the plant, sent to the Laboratory of Gene Expression of UFRPE (LABEG), submitted to evaluations of ornamental interest and later placed in a vessel containing water, leaving about 5 cm from the stem base submerged until the moment of the discard, which was given when the stems presented an unpleasant visual aspect, with flowers, leaves and stem darkened.



152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168

**Figure 5. Cut-off points: (a) Closed head; (b) Semi-open head; (c) Open head.**

The ornamental characteristics evaluated after the cutting of the stems were as follows: Number of Flowers (NF) - obtained by counting open inflorescences; Number of Flower Buds (NB) - obtained by counting inflorescences that did not open; Post-Harvest Durability (PHD) - number of days from stem cutting to disposal.

For the analysis of variance, the effects of the treatments and the averages were considered as fixed and treated according to the statistical model for the specific designs of each experiment.

Using the F test at the 5% probability level, were tested the significance of the mean squares and posteriorly the means were submitted to polynomial regression analysis or comparison of means by the Tukey test using the GENES program [21].

Estimates of variance components and genetic parameters were obtained from the following

expressions:  $\sigma_g^2 = \frac{QMG - QMB}{r}$ ,  $h^2 = \frac{\sigma_g^2}{QMG/r}$ , e  $CV_g = \frac{100\sqrt{\sigma_g^2}}{\mu_n}$ , for the genetic variance among means, heritability coefficient and coefficient of genetic variation, respectively.

169  
170

171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199

### 3. RESULTS AND DISCUSSION

The emergence of seedlings began 4 days after the seeding and continued for two days. The accesses presented 80% and 90% of germination percentage (% G) and 5.47 and 8.22 of Emergency Velocity Index (EVI) for ICA 193 and ICA 73, respectively.

Significant differences were observed between the accesses evaluated at the 5% level by the F test for the analyzed characteristics, except for the diameter of the stem, evidencing the existence of genetic variability (Table 1).

According to [22], the plants are classified as ornamental when they present characteristics that arouse attention and interest, from their aesthetic particularities, referring to the color and shape of leaves and flowers, phenological aspects, among others. Taking these characteristics into consideration, the ICA 73 presented plants with the highest number of branches (9.15), a high number of heads (15.9) and, mainly, high flower production (14.3). On the other hand, it exhibited strong spinescent margin of leaves and bracts (9.25). On the other hand, ICA 193 access presented the most favorable points to the low and moderate spinescent margin of leaves and bracts (2.39 and 3.36, respectively). However, showed low branching (5.25) and consequently, lower number of heads and flower production (7.65 and 6.95, respectively) (Table 2).

According to [23], the leaves and bracts margins of safflower plants are peculiarly spinescent, however the vehemence with which this characteristic is expressed in the plant varies according to the different genotypes, thus allowing the selection and development of varieties that exhibit a weak or moderate character expression, offering attractive materials to the floriculture market.

UNDER PEER REVIEW

200 **Table 1. Summary of variances analysis and genetic parameters estimates for Plant Height (PH), Stem Diameter (SD), Leaf Dentin**  
 201 **(LD), Margin of the Spinescent Leaves (MSL), Number of Branches (NB), Number of Heads (NH), Margin of the Spinescent Bracts**  
 202 **(MSB), Flowering (FI), and Flower Production (FP), Recife, 2017.**

SV	DF	MS								
		PH (cm)	SD (cm)	LD	MSL	NB	NH	MSB	FI	PF
<b>Accesses</b>	1	2822.4	0.006	30.625	122.5	152.1	680.62	90.0	198.02	540.22
<b>Residual</b>	38	6665.2	0.022	5.62	15.20	8.90	36.75	13.35	22.31	28.18
<b>F</b>		16.09**	0.29 <sup>ns</sup>	5.44*	8.06**	17.08**	18.52**	6.74*	8.87**	19.16**
<b>Mean</b>		68.1	0.98	4.12	7.5	7.2	11.77	7.75	56.52	10.62
<b>CV</b>		19.45	15.06	57.50	51.98	41.44	51.48	47.15	8.36	49.97
$\sigma^2_g$		132.35		1.25	5.36	7.16	34.03	3.83	8.78	25.60
$H^2_g$		93.78		81.63	87.60	94.15	94.60	85.16	88.73	94.78
$CV_g$		16.89		27.10	30.88	37.17	48.19	25.26	28.25	47.62
$CV_g/CV_e$		0.87		0.47	0.59	0.90	0.94	0.53	5.24	0.95

203 \* and \*\* significant at the 5% and 1% levels, respectively, of the probability by the F test and "ns" not significant by the F test

204  
 205 **Table 2 - Average of Plant Height (PH), Stem Diameter (SD), Leaf Dentin (LD), Margin of the Spinescent Leaves (MSL), Number of**  
 206 **Branches (NB), Number of Heads (NH), Margin of the Spinescent Bracts (MSB), Flowering (FI), and Flower Production (FP), Recife,**  
 207 **2017.**

Accesses	PH (cm)	SD (cm)	LD	MSL	NB	NH	MSB	FI	FP
<b>ICA 73</b>	59.7b	9.99a	5.0a	9.25a	9.15a	15.9a	9.25a	58.7a	14.3a
<b>ICA 193</b>	76.5a	9.74a	3.25b	2.39b	5.25b	7.65b	3.36b	54.3b	6.95b

208 \*Means followed by the same letter do not differ by Tukey test at 5%.

209 Plants with spines have less acceptance in the market, by virtue of limiting the touch, due to  
210 the possibility of promoting scratches and holes in the skin. In this context, ICA 193 stands  
211 out for having naturally weak or moderate spinescent margins, both in the leaves and in the  
212 bracts, presenting viability for insertion in the market of cut flowers without resistance to  
213 acceptance. In contrast, access ICA 73 needs to be submitted to breeding programs in order  
214 to circumvent this limitation for its use in floriculture.

215  
216 Both accesses presented precocity of flowering, and ICA 73 presented a subtle highlight in  
217 relation to this phenological stage, starting its flowering about 59 DAS, while the ICA 193  
218 access began about 54 DAS. According to [24], this phase starts between 60 and 100 days  
219 after sowing and confirms, therefore, the precocity of the materials evaluated in this work  
220 (Table 2).

221  
222 No early commercial material of *Carthamus tinctorius* intended for the ornamental market is  
223 reported, in addition, it is possible to notice a certain difficulty in the development of cultivars  
224 of this species that present attributes of ornamental interest and initiate this phenological  
225 stage early [15]. Less late varieties, commonly used, show beginning of flowering only 80  
226 days after sowing [25,26]. In this sense, the accesses under study have a scarce and  
227 desired characteristic, offering a further differential to include these in commerce, not only  
228 facilitating acceptance but demonstrating competitiveness with products already available.

229  
230 Regarding plant height, ICA 73 showed lower heights than ICA 193, referring to 59.7 cm and  
231 76.5 cm, respectively (Table 2). Considering also the use for cutting stem, according to [15],  
232 the stem length of products with superior quality must present between 70 cm and 80 cm,  
233 however, there are already commercial safflower varieties specific to the ornamental market  
234 with stems from 60 cm [25]. In this way, the values demonstrated by the accesses of this  
235 research, fit within the allowed for both sides.

236  
237 As to leaf dentin, ICA 73 presented moderate intensity (5.0) and weak to moderate ICA 193  
238 (3.25) (Table 2). In order to make arrangements, the margin of the leaf does not have a fixed  
239 pattern, since even the most unusual can contribute to creative and decorative  
240 combinations, including being something very desired to compose bouquets base [27].

241  
242 During the experiment, the plants were affected by pathogens and pests, since no chemical  
243 control was performed. From the symptoms and a previous microscopic analysis, the  
244 presence of *Cercospora carthami* and aphid (aphis) was observed (Figures 6a and 6b).  
245 However, the inflorescences were not directly affected and the characteristics of interest  
246 could be effectively evaluated. [12] reported that, safflower is the target of many pathogens,  
247 including fungi, bacteria and viruses, but the first group cited is the most prominent.  
248 *Cercospora carthami* is one of the fungi that commonly affect the culture, causing foliar  
249 damage. Among the pests, aphids are said to cause the most recurrent damages, however  
250 they are less worrisome than diseases [28].

251  
252 The summary of the analysis of variance for the second experiment and the estimation of the  
253 main genetic parameters for the agronomic and ornamental characters evaluated in the two  
254 safflower accesses are organized in Table 3. According to the results, it is possible to  
255 observe a significant difference between the accesses at the level of 5% by the F test for the  
256 characteristics analyzed.





257  
258  
259  
260  
261  
262  
263

Figure 6. Safflower plants: (A) Pest attack (aphids); (B) Leaf disease (*Cercospora carthami*).

Tabela 3. Summary of variances analysis and genetic parameters estimates for Plant Height (PH), Stem Diameter (SD), Number of Branches (NB), Number of Heads (NH), and Flower Production (FP), Recife, 2017.


SV	DF	MS				
		PH (cm)	SD (cm)	NB	NH	PF
Blocks	3	36.68	0.03	5.54	7.10	5.78
Accesses	1	570.94*	0.51*	283.52**	264.97**	242.91**
Densities	3	325.76*	0.27*	210.51**	191.36**	188.82**
AccessesxDensities	3	30.41 <sup>ns</sup>	0.04 <sup>ns</sup>	33.31**	27.76**	27.78**
Residual	21	79.10	0.05	2.90	3.58	2.93
Mean		60.30	1.75	10.69	10.43	10.05
CV		14.74	12.61	15.94	18.14	17.04
$\sigma_g^2$		30.74	0.03	17.54	16.34	15.00
H <sup>2</sup>		86.15	90.46	98.97	68.65	98.79
CV <sub>g</sub>		9.19	9.71	39.17	38.73	38.55
CV <sub>g</sub> /CV <sub>e</sub>		0.62	0.77	2.46	2.13	2.26

264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276

\* and \*\* significant at the 5% and 1% levels, respectively, of the probability by the F test and "ns" not significant by the F test.

Figure 7 graphically shows the behavior of the accesses as a function of the different densities of plants per vase, as well as the equations and coefficient of determination (R<sup>2</sup>) that best fit the variables studied, according to the regression analysis. All variables can be explained by the linear equation of the 1<sup>st</sup> degree, with R<sup>2</sup> values higher than 0.80.

Plant height and stem diameter were inversely proportional to plant density per vase, decreasing as the number of plants increased (Figure 7a). The diameter is an important feature because it is related to rigidity and quality of the stem, since low densities can lead to flexibility and breakage [29]. Pruning did not limit the final length of the plants, which reached values characteristic of the species.



277  
 278  
 279  
 280

**Figure 7. Plant Height (PH), Stem Diameter (SD), Number of Branches (NB), Number of Heads (NH), and Flower Production (FP) of ICA 73 and ICA 193 safflower accesses according to four plant densities per pot.**

281 The highest values of height and diameter were reached by ICA 193 access, with values  
 282 ranging from 56.75 to 70.00, and 1.71 to 2.21 (Figure 7a). In the first case, the values  
 283 extrapolate the recommendation of [30] for use in vases, but is suitable for employment in  
 284 other areas of social recreation. In this sense, there are already commercial safflower  
 285 varieties destined for the ornamental market with heights between 60 and 80 cm, such as  
 286 Orange Granade, which is highly prized for beautifying gardens [25]. In contrast, ICA 73  
 287 exhibited the smallest values of height and diameter, ranging from 49.00 to 63.25 and 1.44  
 288 and 1.80 (Figure 2a). Despite the low diameter values, the stems showed to be well lignified  
 289 and no breaks were observed.

290  
 291 The decrease due to the increase in the number of plants per vase for number of branches,  
 292 heads and flower production was also observed (Figure 7b and 7c) and is in agreement with  
 293 the one verified by [31], that in its work with other safflower genotypes evidenced a linear  
 294 reduction proportional to the increase of the density. This result is probably linked to  
 295 competition between plants for nutrients, water and light, limiting their development [29]. For  
 296 use in potted plants or gardens, plants with more branches, provide a aerial part more  
 297 voluminous and visually pleasing. Access ICA 73, presented the highest values for these  
 298 characteristics and the best density for both genotypes, refers to 1 plant per vase.

299  
 300 The high number of branches acts negatively on the uniformity of the opening of the flowers,  
 301 due to the different flowering rates of the heads [29]. For ornamental plants destined to  
 302 gardens, vases or other leisure areas, this particularity becomes attractive, because it makes  
 303 possible that the prestige of the flowers can be realized by a greater period of time, since  
 304 while the first flowers are close to senescence, others will still be at the beginning of the  
 305 anthesis. On the other hand, this factor is not attractive for cutting stems.

306  
 307 In Table 4, the analysis of variance of the third experiment and the estimation of the main  
 308 genetic parameters for characters of ornamental importance evaluated in the two accesses  
 309 of safflower. The results show a significant difference between the accesses at the 5% level  
 310 by the F test for the characteristics analyzed. Table 5 shows the means of the variables that  
 311 were submitted to the Tukey test, depending on the treatment factors: accesses and cut-off  
 312 points.

313  
 314 **Table 4. Summary of variances analysis and genetic parameters estimates for Number**  
 315 **of Flowers (NF), Number of Flowers Buds (NFB), Post-Harvest Durability (PHD),**  
 316 **Recife, 2017.**

SV	DF	MS		
		NF	NFB	PHD
Accesses	1	54.0**	84.37**	88.17**
Cut-off Point	2	26.54**	22.79**	77.17**
AccessesxCut-off	2	13.62**	15.87**	4.67 <sup>ns</sup>
Residual	18	1.17	1.12	1.33
Mean		5.33	3.21	10.58
CV		20.03	33.06	10.91
$\sigma^2_g$		4.40	6.94	7.24
H <sup>2</sup>		97.84	98.67	98.49
CV <sub>g</sub>		39.34	82.10	25.42
CV <sub>g</sub> /CV <sub>e</sub>		1.94	2.48	2.33

317 \* and \*\* significant at the 5% and 1% levels, respectively, of the probability by the F  
 318 test and "ns" not significant by the F test.

319 The maturity of the flowers is a very decisive characteristic on the quality of the product and  
 320 makes it impossible in most cases to perform a mechanized harvest [15], justifying the  
 321 importance of defining the best moment for cutting the stems, as far as the anthesis of  
 322 flowers is concerned. Other important information that should be considered refers to the  
 323 fact that the central flower opens days before the lateral flowers, about one week, leading to  
 324 visual depreciation when the other flowers are opening [32], for this reason the pinch was  
 325 performed, stimulating the anther of the lateral flowers in a more uniform way.  
 326

327 The number of flowers, flower buds and post-harvest durability should be analyzed together  
 328 and as a function of the cut-off point, allowing to establish the most appropriate combination  
 329 for quality of the final product.  
 330

331 The ICA 73 had a higher average number of open flowers (10.25) and a lower number of  
 332 floral buds (1.75) for the cut-off point when flowers were open, however, in this same  
 333 treatment the lower post-harvest durability of the stems was obtained (4.25), making it  
 334 impossible to cut stems of this material at this maturation level, since it does not meet an  
 335 adequate number of days of product life (Table 5). At the point of semi-open flowers, the  
 336 number of flowers (5.75) was reduced by half and the number of flower buds increased  
 337 (6.00), showing a nearly 1: 1 ratio between flowers and buds, indicating little uniformity of  
 338 flower anthesis and opening of a few buds after cutting, but exhibited longer flower durability  
 339 (11.00) (Table 5). Finally, at the point of closed flowers the number of flowers reduced even  
 340 more, evidencing that some of the heads did not even develop buds and the low number of  
 341 buds also confirms this hypothesis, however, the durability of the stems was equivalent to  
 342 the cutting treatment with the semi-open flowers.  
 343

344 **Table 5 - Average of Number of Flowers (NF); Number of Flowers Buds (NFB); Post-**  
 345 **Harvest Durability (PHD), Recife, 2017.**

	Variables					
	NF		NFB		PHD	
	ICA 73	ICA 193	ICA 73	ICA 193	ICA 73	ICA 193
Open Head	10.25Aa	4.25Ba	1.75Ac	1.25Bc	4.25Bb	9.75Ab
Semi-open Head	5.75Ab	4.50Ba	6.00Ab	0.50Bb	11.00Ba	13.50Aa
Closed Head	4.50Ab	2.75Bb	7.50Aa	2.25Ba	10.75Ba	14.25Aa

346 *\*Means followed by the same lower letters in column and capital letters on the lines*  
 347 *do not differ significantly by the Tukey test at 5%.*  
 348

349 ICA 193 presented a lower average number of open flowers when compared to ICA 73 at all  
 350 cut-off points, however, it also exhibited a lower number of flower buds, indicating a greater  
 351 uniformity of flower anthesis. At the cut-off point with open flowers, exhibited the second  
 352 highest value of flowers (4.25), according to lower flower buds values (1.25) and lower post-  
 353 harvest durability of the stems (9.75), however, this useful life is already acceptable to the  
 354 market (Table 5). For the cut-off point with semi-open flowers, it presented the highest  
 355 number of flowers (4.50), although it does not differ statistically from the previous treatment  
 356 for this characteristic, smaller number of flower buds (0.50) and second highest number of  
 357 days of stem durability (13.50). Finally, at the cut-off point with closed flowers, the lowest  
 358 number of flowers (2.75) and the highest number of floral buds (2.25) were observed,  
 359 evidencing the difficulty of developing the buds after cutting the stems, also presented the  
 360 greatest number of days of durability of the stems (14.25), but did not differ statistically from  
 361 the previous treatment for this characteristic.  
 362

363 According to [32], the most favorable cutting stage of the stems corresponds to the period in  
 364 which 20% to 30% of the central florets opened, which is equivalent to the cut-off point

365 denominated in this work as semi-open heads, allowing the others to open in the hands of  
366 the consumer, extending the useful life of the product. In agreement with this affirmation, it is  
367 observed that ICA 193 presented the best combination of factors for this cut-off point,  
368 presenting a higher number of flowers, lower number of buds and greater post-harvest  
369 durability. The same observation can be raised for ICA 73, but with some reservations, such  
370 as the performance of a removal of the secondary and tertiary branches, improving the  
371 aesthetics of the product, since despite an adequate number of flowers and stem durability,  
372 it presented high number of buds, or search for improvements of this characteristic through  
373 an improvement program.

374  
375  
376  
377  
378  
379  
380  
381  
382  
383

The flowers produced by both genotypes showed a yellow color at the beginning of the anthesis, changing to orange shades soon after and presented a very attractive visual aspect, with abundant beauty while fresh and even after a period of drought, offering potential for introduction into the Brazilian flower market, contributing to the supply of news for the sector and the consumer (Figure 8a and 8b). According to [23], the characteristics of greater importance and influence on the ornamental value of safflower are attributed to the color of the flowers, where the oranges and yellows stand out, along with the weak spinescent margin of leaves and bracts.



384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401

**Figure 8. (A) Bouquet of stems after cutting; (B) Change the color of the flower.**

The evaluated characteristics presented high estimates of the genetic parameters of heritability and ratio between the coefficients of genetic and experimental variation, a very favorable point in breeding programs, since it indicates in a general way that these characters can be easily improved through classic methods [33] and provide favorable conditions for realization of selection, allowing to obtain high genetic gain within the first cycles [34].

Considering the differences evidenced between the genotypes, together with the completeness they demonstrate for characteristics of ornamental interest, these genotypes suggest potential for inclusion in an improvement program, in order to obtain a material that groups the positive characteristics presented in both accesses. [35] point out that one of the criteria for success in crossbreeding depends on the divergence between the parents, parallel to the superior performance they present referring to the characteristics of interest of the breeder.



402 **4. CONCLUSION**

403

404 The accesses ICA 73 and ICA 193 have ornamental potential, coupled with the precocity,  
405 beauty and durability of their flowers. For plant vase, the best density for cultivation refers to  
406 one plant per vase, allowing better expression of the plants' ornamental potential. For cutting  
407 stem, the cutting point referring to semi-open heads was the major contributor to obtaining  
408 quality stems.

409

410

411 **REFERENCES**

412

413 1. Romero DHS, Restrepo IME. Local competitive profile as a determining factor for the  
414 development of floriculture in Madrid (Cundinamarca). Research and reflection. 2011; 19 (2):  
415 25-43.

416

417 2. Hortiwise. A Study on the Kenyan-Dutch Horticultural Supply Chain. The Dutch Ministry of  
418 Economic Affairs, Agriculture & Innovation The Netherlands. 2012.

419

420 3. Mitsueda NC, Costa EV; D'oliveira PS. Environmental aspects of agribusiness flowers and  
421 ornamental plants. Magazine in Agribusiness and Environment. 2011; 4 (1): 9-20.

422

423 4. Ibraflor. Brazilian Institute of Floriculture. The flower market in Brazil. 2017. Accessed: 20  
424 Dezember 2018. Available: <http://www.ibraflor.com/site/2017/11/04/florente-de-flores-vera-longuini/>.

425

426 5. Junqueira AH, Peetz MS. The productive sector of flowers and ornamental plants of  
427 Brazil, from 2008 to 2013: updates, balance sheets and perspectives. Brazilian Journal of  
428 Ornamental Horticulture. 2014; 20 (2): 115-120.

429

430 6. Ibraflor. Brazilian Institute of Floriculture. Sector Numbers. 2014. Accessed: 28 November  
431 2017. Available: [http://www.IBRAFLOR.com/ns\\_mer\\_interno.php](http://www.IBRAFLOR.com/ns_mer_interno.php).

432

433 7. Dantas CVS, Silva IB, Pereira GM, Maia JM, Lima JPMS, Macedo CEC. Influence of  
434 sanity and water deficit on seed germination of *Carthamus tinctorius* L. Brazilian Journal of  
435 Seeds. 2011; 33 (3): 574-582.

436

437 8. Santos RF, Silva MA. *Carthamus tinctorius* L.: An alternative crop for Brazil. Acta Iguazu.  
438 2015; 4 (1): 26-35.

439

440 9. Zareie S, Mohammadi-Nejad G, Sardouie-Nasab S. Screening of Iranian safflower  
441 genotypes under water deficit and normal conditions using tolerance indices. Australian  
442 Journal of Crop Science. 2013;7(7):1032-1037.

443

444 10. Landau S, Friedman S, Brenner S, Bruckental I, Weinberg ZG, Ashbell G, Hen Y, Dvash  
445 L, Leshem Y. The value of safflower (*Carthamus tinctorius* L.) hay and silage grow under  
446 Mediterranean conditions as forage for dairy cattle. Livestock Production Science.  
447 2004;88:263-271.

448

449 11. Asgarpanah J, Kazemivash N. Review: Phytochemistry, Pharmacology and Medicinal  
450 Properties of *Carthamus tinctorius* L. Chin J Integr Med. 2013;19(2):153-159.

451

452 12. Mündel, H. H.; Blackshaw, R. E.; Byers, J. R.; Huang, H. C.; Johnson, D. L.; Keon, R.;  
453 Kubik, J.; Mckenzie, R.; Otto, B.; Roth, B; Stanford, K. Safflower production on the Canadian

- 454 prairies: revisited in 2004. Agriculture and Agri-Food Canada. Lethbridge Research Centre:  
455 Lethbridge; 2004.  
456
- 457 13. Pintão AM, Silva IF. The Truth About Saffron. In: Workshop medicinal and  
458 phytotherapeutic plants in the tropics. Lisbon. Summaries ... Lisbon: IICT / CCCM, 2008.  
459 Electronic Version.  
460
- 461 14. Jadhav BA, Joshi AA. Extraction and Quantitative Estimation of Bio Active Component  
462 (Yellow and Red Carthamin) from Dried Safflower Petals. Indian Journal of Science and  
463 Technology. 2015;8(16)1-5.  
464
- 465 15. Uher, J. Safflower in European floriculture: a review. In: Proceedings of the 7th  
466 International Safflower Conference, Wagga Wagga, New South Wales, Australia. 2008.  
467
- 468 16. Bradley VL, Guenther RL, Johnson RC, Hannan R M. Evaluation of safflower  
469 germplasm for ornamental use. In: Perspectives on new crops and new uses (J. Janick, ed.).  
470 ASHS, Press: Alexandria. 1999. 433-435.  
471
- 472 17. Oliveira GG. Trichoderma spp. plant growth and biocontrol of Sclerotinia sclerotiorum  
473 and pathogens in safflower seeds (*Carthamus tinctorius*). 2007. 79 p. Dissertation (Master in  
474 Plant Production), Federal University of Santa Maria, Santa Maria, 2007.  
475
- 476 18. Zoz T. Correlation and analysis of yield traits in grains and their components and plant  
477 characteristics in safflower (*Carthamus tinctorius* L.) and castor bean (*Ricinus communis* L.).  
478 2012. 54 p. Dissertation (Master's degree in Agronomy) Faculty of Agronomic Sciences-  
479 UNESP; 2012.
- 480
- 481
- 482 19. Maguire JD. Speed of germination aid in selection and evaluation for seedling  
483 emergence and vigor. *Crop Science*. 1962;2(2):176-77.  
484
- 485 20. Stumpf ERT, Heiden G, Barbieri RL, Fischer SZ, Neitzke RS, Zanchet B, Grolli PR.  
486 Method for evaluating the ornamental potential of native and unconventional flowers and cut  
487 foliage. *Brazilian Journal of Ornamental Horticulture*. 2007; 13: 143-148.  
488
- 489 21. Cruz CD. GENES - versão Windows. Viçosa- MG: Editora UFV; 2006.  
490
- 491 22. Gonçalves MF, Melo AGC. Floristic analysis of ornamental plants implanted in Garça  
492 Forest / SP. *Electronic Journal of Forest Engineering*. 2013; 21 (1): 12-24.
- 493
- 494 23. Pahlavani MH, Mirlohi AF, Saeidi G. Inheritance of flower colour and spininess in  
495 safflower. *Journal of Heredity*. 2004;95:265-267.  
496
- 497 24. Emongor V. Safflower (*Carthamus Tinctorius* L,) the underutilized and neglected crop: A  
498 review. *Asian Journal of Plant Science*. 2010;9(6):299-306.  
499
- 500 25. Sakata. Sakata's reliable seeds. Flower seed. Sakata seed corporation, Yokohama,  
501 Japan. 2010. 87 p.  
502
- 503 26. Geneses Seeds Ltd. Catalog Flowers. Accessed: 20 November 2017. Available:  
504 [http://www.genesisseeds.com/PDFs/GENESIS\\_CATALOG\\_FLOWERS.pdf](http://www.genesisseeds.com/PDFs/GENESIS_CATALOG_FLOWERS.pdf).

- 505  
506 27. Morais ÉB, Castro ACR, Silva TF, Soares NS, Silva JP. Evaluation of potential use of  
507 native Anthurium foliage. Ornamental Horticulture. 2017;23:07-14.  
508  
509 28. Coronado LM. El cultivo Del cártamo (*Carthamus tinctorius* L.) em México. Instituto  
510 nacional de investigacion esforestales, agrícolas y pecuárias. México, 2010.  
511  
512 29. Bellé RA, Rocha EK, Backes FAAL, Neuhaus M, Schwab NT. Safflower grown in  
513 different sowing dates and plant densities. Ciência Rural. 2012;42(12).  
514  
515 30. Motos J, Oliveira MJG. Produção de crisântemos em vaso. Holambra, Flortec. 1998. 34  
516 p.  
517  
518 31. Sampaio MC. Cultivo de cártamo (*Carthamus tinctorius* L.) sob variação de adubações,  
519 densidades e épocas de plantio. 2016. 63p. Dissertação (Mestrado em Engenharia de  
520 Energia na Agricultura), Universidade Federal do Oeste do Paraná, Cascavel, 2016.  
521  
522 32. Wien HC. Cut Flower Cultural Practice Studies and Variety Trials 2012. Dept. of Hort.,  
523 Cornell Univ. Ithaca, 2012.  
524  
525 33. Fox GP, Bowman J, Kelly A, Inkerman A, Poulsen D, Henry R. Assessing for genetic and  
526 environmental effects on ruminant feed quality in barley (*Hordeum vulgare*). Euphytica.  
527 2008;163:249-257.  
528  
529 34. Vencovsky R, Barriga P. Genética Biométrica no Fitomelhoramento. Ribeirão Preto:  
530 SBG; 1992.  
531  
532 35. Cruz CD, Carvalho SP, Vencovsky R. Estudos sobre divergência genética II. Eficiência  
533 da predição do comportamento de híbridos com base na divergência de progenitores.  
534 Revista Ceres.1994;41(234):183-190.