

2

3 **Nitrogen fertilization and inoculation of seeds**

4 **with *Rhizobium tropici* on the agronomic**

5 **performance of common beans**

6

7

8

10

11 **ABSTRACT**

12

Aims: This study aimed to evaluate the influence of nitrogen fertilization and inoculation of seeds with *Rhizobium tropici* on the agronomic performance of common bean (*Phaseolus vulgaris* L.).

Study design: The experimental design consisted of a randomized block in a 4x2 factorial layout, with 4 replicates, the first factor referring to inoculant doses (0, 50, 100 and 200 mL for each 25 kg of seed), while the second refers to nitrogen fertilization (0 and 40 kg ha of N).

Place and Duration of Study: The study was conducted to field in a no-till system area, in the southwestern region of Paraná, Brazil. The soil is classified as a Purple Latosol, with a clayey texture.

Methodology: The adopted spacing was 0.45 m between rows, and the seeding density used was 12 seeds per furrow meter. The inoculants NITRO 1000 *Rhizobium tropici* SEMIA 4077 and SEMIA 4088 were applied, varied according to the treatments. The cultivar used was IPR – Tangará.

Results: As for grain yield, it was observed that nitrogen fertilization did not contribute to the increase in grain yield, and higher productivity was obtained in the absence of N. This was probably due to the high content of organic matter present in the soil.

Conclusion: Nitrogen fertilization at sowing and seed inoculation with *Rhizobium tropici* did not influence the plant population and the 1000-grain mass. In the absence of nitrogen fertilization at sowing, pods with longer length and higher grain yield were obtained. Inoculation of the seeds with *Rhizobium tropici* exerts a positive influence on plant height, number of nodes of the main stem, number of pods per plant, and number of beans per pod, the dose recommended by the manufacturer (100 mL) being efficient, with the possibility of applying a dose of 50 mL, to satisfactory results.

13

14 *Keywords: Phaseolus vulgaris L., Biological Nitrogen Fixation, Inoculation, IPR-Tangará*

15

16

17

18 **1. INTRODUCTION**

19

20 The common bean (*Phaseolus vulgaris* L.) is grown in more than 100 countries, although

21 over 60% of world production is takes place in only five countries, Brazil being the largest

22 producer and consumer in the world. The area cultivated with beans in Brazil as of the

23 2017/18 harvest was estimated at 3.2 million hectares, 0.5% higher than in the 2016/17

24 harvest. The mean productivity is projected at 1,043 kg ha⁻¹, 2.4% less than in the last
25 harvest. Considering the sown area and expected productivity, bean production in the
26 2017/18 harvest is expected to reach 3.35 million metric tons, 1.9% less than in the previous
27 harvest [1].

28

29 A number of factors interfere in the yield of common bean, including nitrogen fertilization
30 management. Inadequate nitrogen fertilization is a factor that often determines the failure of
31 common bean crops. While a few producers apply excessive doses of N, others apply
32 insufficient amounts of this element, limiting crop yield even if other factors of production are
33 optimized [2]. Furthermore, it is necessary to correctly manage the N, as it is a nutrient that
34 can easily be lost through the leaching, volatilization or denitrification processes [3].

35

36 N is the nutrient required in greatest quantity by beans [4] and although there is a
37 recommendation for the use of nitrogen fertilizer for common bean crops, research results
38 suggest that it is possible for this crop to benefit from biological nitrogen fixation (BNF), it
39 may contribute to increased crop yield and reduce or replace the use of nitrogen fertilizers
40 [5], thereby decreasing production costs and the environmental impact of the use of
41 chemical fertilizers. It should be noted that the high dependence on fertilizers by crops, for
42 example, causes an increase in the energy costs for conversion of atmospheric nitrogen, in
43 addition to the extraction process of the other elements, such as phosphorus and potassium
44 [6]. Moreover, excess nutrients applied in conventional agriculture may cause environmental
45 problems. Environmental costs of all nitrogen losses in Europe have been estimated at €70-
46 320 billion per year, which outweighs the direct economic benefits of nitrogen in agriculture
47 [7].

48

49 The practice of seed inoculation with rhizobia represents a low-cost alternative to increase
50 the yield of common bean, in addition to avoiding the contamination of water resources by
51 nitrogen fertilizer and reducing greenhouse gas emissions. It is proven to be a cheap
52 technology, simple to use, and with good economic return. Being a legume, the common
53 bean presents conditions to benefit from symbiotic association with *Rhizobium*, which
54 contributes specifically to N savings. Nevertheless, according to Rabelo et al. [8], there is still
55 a need for improvements in order to develop an inoculation that fully replaces nitrogen
56 fertilization in common beans, as is the case with soybeans. FBN contributes positively to
57 the development of this culture, but not to a sufficient degree to dispense N applications.

58

59 This study aimed to evaluate the influence of nitrogen fertilization and inoculation of seeds
60 with *Rhizobium tropici* on the agronomic performance of common beans.

61

62 2. MATERIAL AND METHODS

63

64 The experiment was conducted in an area under a no-till system, located on the São Pedro
65 da Bandeira line, in the municipality of Dois Vizinhos, Paraná, Brazil. The municipality is
66 located in the Southwest region of the State, between the geographic coordinates 25°44' S
67 and 53°03' W, with an altitude of 509 m.

68

69 The climate of the region according to the Köppen classification subtropical hot humid (Cfa),
70 with a mean annual rainfall of 2,044 mm, well distributed throughout the year. The mean
71 daytime temperature is 19.6°C and the mean nighttime temperature is 15.2°C [9]. The mean
72 rainfall and temperature in the months of the experiment are shown in Table 1.

73

74

75

76 **Table 1. Rainfall (mm), maximum temperature (MAX), minimum temperature (MIN), and**
 77 **mean (MDA) in °C, in the months of execution of the experiment.**

Month	Precipitation (mm)	Temperature		
		MAX	MIN	MDA
November	112	31.4	21.4	26.4
December	266	32.03	24.09	28
January	200	32.96	22.23	27.5

78 Data obtained at the property where the experiment was conducted.

79

80 The soil is classified as a Purple Latosol, with a clayey texture [10]. The area is
 81 characterized by presenting a slightly uneven topography, with a level curve and no erosion.
 82 Prior to the sowing of beans, the area had been sown with an oat crop. Prior to the bean
 83 sowing, a soil sample was collected at a 0-20 cm depth. The results obtained were: pH
 84 CaCl_2 6,7, P 40.05 mg dm^{-3} , organic matter 38.28 g dm^{-3} , Ca, Mg, K, Al and H+Al, respectively,
 85 8.18, 3.33, 0.78, 0.0, 2.37 cmolc dm^{-3} and V 83.83%.

86

87 A Semeato Personale seed drill was used for sowing the beans. The adopted spacing was
 88 0.45 m between rows, and the seeding density used was 12 seeds per furrow meter. The
 89 seeds were treated with thiamethoxan 350 g L^{-1} at the dose 0.7 L of the commercial product
 90 for 100 kg of seed. Following that, the inoculants NITRO 1000 *Rhizobium tropici* SEMIA
 91 4077 and SEMIA 4088 were applied. The inoculant dose, as well as the nitrogen fertilization,
 92 varied according to the treatments. For phosphate and potassium fertilization, 40 kg ha^{-1} was
 93 used. The fertilizer was applied to the side and below the seed.

94

95 The cultivar used was IPR – Tangará. This cultivar belongs to the Carioca group and
 96 presents an indeterminate growth habit, erect plants with long vines, a mean cycle of 87
 97 days from emergence to harvest, and mean productive potential of 3,326 kg ha^{-1} . It has
 98 resistance to common mosaic, *Curtobacterium* wilt, *Fusarium* wilt and rust, mild resistance to
 99 powdery mildew and angular leaf spot, and susceptibility to anthracnose and common
 100 bacterial blight. It presents intermediate tolerance to high temperatures and droughts
 101 occurring during the reproductive phase. The seeds have a light beige coat with light brown
 102 stripes.

103

104 The experimental design consisted of a randomized block, with a 4x2 factorial layout, with
 105 four replications. The first factor refers to four levels of seed treatment with different doses of
 106 inoculant (0 mL, 50 mL, 100 mL, and 200 mL for every 25 kg of seed). The recommended
 107 dose according to the manufacturer is 100 mL for every 25 kg of seed. The second factor refers to
 108 two levels of nitrogen fertilization (0 and 40 kg ha^{-1} of N at sowing). The treatments were as
 109 follows: $T_1 = 0 \text{ kg ha}^{-1} \text{ N}$ and 0 mL inoculant; $T_2 = 0 \text{ kg ha}^{-1} \text{ N}$ and 50 mL inoculant; $T_3 = 0 \text{ kg}$
 110 $\text{ha}^{-1} \text{ N}$ and 100 mL inoculant; $T_4 = 0 \text{ kg ha}^{-1} \text{ N}$ and 200 mL inoculant; $T_5 = 40 \text{ kg ha}^{-1} \text{ N}$ and 0
 111 mL inoculant; $T_6 = 40 \text{ kg ha}^{-1} \text{ N}$ and 50 mL inoculant; $T_7 = 40 \text{ kg ha}^{-1} \text{ N}$ and 100 mL
 112 inoculant; and $T_8 = 40 \text{ kg ha}^{-1} \text{ N}$ and 200 mL inoculant.

113

114 Each experimental plot had 5 m of length composed of 5 lines and spacing of 0.45 m,
 115 totaling an area of 11.25 m^2 . The three central lines were considered as usable area of the
 116 plot, excluding 0.5 m from each end of the line, totaling 4.5 m^2 .

117

118 During the development of the crop, two manual weeding treatments were performed, at 10
 119 and 25 days after sowing. Phytotoxic treatments (125 mL ha^{-1}) were also prepared aiming at
 120 the control of pests such as whitefly, white mite, leafhoppers, Cucurbit beetle and thrips with
 121 thiamethoxam (125 mL ha^{-1}) and diseases such as angular leaf spot, anthracnose and rust
 122 with fentin hydroxide (700 mL ha^{-1}), these comprising pests and diseases of major

123 importance, which may cause greater damage to the common bean in the southwestern
124 region of Paraná.

125

126 The evaluations were carried out at the end of the crop cycle. The following variables were
127 evaluated: plant population, plant height, number of main stem nodes, number of pods per
128 plant, number of grains per pod, mean pod length, 1000-grain mass (13% moisture), and
129 productivity.

130

131 To determine the plant population, the number of plants was counted in 2 m of the usable
132 plot. Plant height was determined by averaging 20 plants harvested from each usable plot,
133 measured from the plant neck to the end of the main stem. The number of nodes of the main
134 stem, number of pods per plant, number of beans per pod and the mean pod length were
135 determined by averaging 20 plants randomly collected from each usable plot. The 1000-
136 grain mass was determined by averaging 2 samples of each usable plot, each sample
137 having 250 grains, then weighed in a semi-analytical balance and then calculated the
138 moisture content corrected to 13%, estimating the 1000-grain mass. Grain productivity was
139 determined throughout the plot area, correcting for moisture content of 13% and,
140 subsequently, estimated productivity in kg ha⁻¹.

141

142 3. RESULTS AND DISCUSSION

143

144 There was no significant effect of the interaction between the inoculant doses and nitrogen
145 fertilization for the variables of plant population, mean pod length, one 1000-grain mass, and
146 productivity. Thus, the results are presented independently. For these variables, there was
147 also no significant effect of inoculant doses. A significant difference was observed for
148 nitrogen fertilization only for the variables of mean pod length and grain yield (Table 2).

149

150 Nitrogen application at sowing had no effect on the plant population (Table 2). This can be
151 explained by the fact that the seed contains the nutrients necessary for the establishment of
152 the seedling, such as in the Fabaceae (beans, soybean), in which the endosperm is partially
153 or fully absorbed during the development of the seed in favor of the cotyledons, which
154 assumes the function of tissue [11]. Additionally, the temperature, which had an average
155 value of 26.4 °C (Table 1) may have contributed to the homogeneity of the booth, it being
156 noted that the optimal temperature for bean germination is around 28 °C, and rainfall
157 occurred shortly after sowing (Table 1) [12].

158

159 **Table 2. Population of plants per hectare, mean pod length (cm), 1000-grain mass (g)**
160 **and yield (kg ha⁻¹) of common beans as a function of nitrogen fertilization.**

161

	Plant population (ha)	Mean pod length (cm)	1000-grain mass (g)	Yield (kg ha ⁻¹)
Without N	239.777 a	9.12 a	284.31 a	1,716.6 a
With N	237.333 a	8.88 b	287.37 a	1,686.6 b
CV	3.7	2.31	2.33	1.05

162

163

164

Mean values followed by the same lowercase letter in the columns do not differ significantly from each other by the F test (P<0.05).

165

166

167

168

169

170

For mean pod length, it was observed that there were no significant differences in the different doses of the inoculant (Table 2). This can be explained by the fact that this component presents high genetic heritability, being intrinsically linked to the characteristic of the cultivar [13]. Notwithstanding, a significant difference was observed for nitrogen fertilization, in which the best result was observed in the absence of N, with 9.12 cm (Table 2). This was probably due to the high content of organic matter present in the soil.

171
172 The 1000-grain mass and grain yield were also not influenced by nitrogen fertilization at
173 sowing (Table 2), as well as by seed inoculation. As for grain yield, it was observed that
174 nitrogen fertilization did not contribute to the increase in grain yield, and higher productivity
175 ($1,716.6 \text{ kg ha}^{-1}$) was obtained in the absence of N (Table 2). This was probably due to the
176 high content of organic matter present in the soil.
177

178 It was also possible to observe that the different inoculant doses did not present significant
179 differences. The high production levels obtained, even in the absence of inoculation,
180 demonstrate the great capacity of competition and effectiveness in symbiotic N fixation by
181 native strains. The yields obtained in the absence of inoculation were similar to high doses in
182 some treatments. The non-response to seed inoculation in relation to productivity can be
183 explained by the presence of native *Rhizobium* strain in the soil, which is usually more
184 aggressive than the introduced strain. The presence of native strains in the soil makes it
185 difficult to perform the introduced strain, as they compete for the sites of nodular infection,
186 their presence in the nodulation in plants that did not receive the inoculant being evident,
187 even in a smaller number of nodules, which may be equivalent to other treatments thanks to
188 their greater efficiency [14].
189


190 Ferreira et al. [14] studying *Rhizobium* strains, concluded that the inoculation of efficient
191 strains in common bean-nodulating cultivar, or their cultivation in soils with an efficient native
192 population, may allow the non-use of nitrogen in the bean crop without affecting productivity.
193 Rabelo et al. [8] mention that FBN contributes positively to the development of this crop, but
194 not enough to dispense N applications.
195

196 Peres et al. [15], in three years of cultivation and using several cultivars, verified that the
197 inoculation with rhizobia contributed to the increase in grain yield. These data show that the
198 biological nitrogen fixation may supplement the nitrogen fertilization in the bean crop,
199 allowing a reduction in nitrogen fertilizer rates, without losses in grain yield. Nevertheless,
200 Matoso and Kusdra [16] did not find positive results regarding the inoculation of common
201 bean with *R. tropici*.
202

203 There was a significant effect of the interaction between inoculant doses and nitrogen
204 fertilization for the variables of plant height, number of main stem nodes, number of pods per
205 plant, and number of grains per pod.
206

207 Following the interaction between nitrogen fertilization and seed inoculation, a similar
208 behavior can be observed for plant height, number of nodes of the main stem per plant,
209 number of pods per plant, and number of grains per pod. Nitrogen fertilization obtained the
210 lowest mean values in all studied variables, even in inoculated treatments. In the absence of
211 nitrogen fertilization, inoculation provided a considerable increase in some agronomic
212 variables of importance, such as the number of pods per plant and number of grains per
213 pod.
214

215 The height of the plants showed a significant interaction ($P < 0.05$). It was noted that, in the
216 absence of N, the values were higher, and that these values decreased when increasing the
217 inoculant dose (Figure 1), thereby reinforcing the idea of action of the native strains in the
218 area. The adaptation of rhizobia to the soil depends on the biotic and abiotic conditions of
219 the environment and the wild or cultivated legume species, both in size and variability. As
220 the cultivated area has a high content of organic matter and has been worked under a no-till
221 system for a long time, the biotic, physical and chemical conditions of the soil are in
222 favorable conditions for the development of the native strains.
223




224
225
226
227

Fig 1. Plant height of cv. IPR Tangará bean as a function of inoculant doses and nitrogen fertilization.

228 When the same inoculant doses associated with nitrogen fertilization (40 kg ha^{-1} of N) were
229 studied, the best inoculation response was obtained, and the best result was observed when
230 applied at half the recommended inoculant dose (50 mL). Although the mean values were
231 lower than in treatments without N use (Figure 1). N fertilizer may cause a reduction in
232 symbiotic efficiency but, when applied in small amounts in the bean crop, allows an increase
233 in nodule growth and higher BNF, and very low levels of nitrate in the soil may be limiting to
234 symbiotic activity. According to Novo et al. [17], root nodulation supplies the needs of the
235 plants, avoiding the nitrogen fertilization, as it inhibits the formation of the nodules, affecting
236 BNF the biomass production.

237
238 As for the number of nodes of the main stem, results similar to those already studied were
239 observed, the best results being observed in the absence of nitrogen fertilization. In the
240 latter, the lower dose of inoculant stands out over the others (Figure 2), followed by half of
241 the recommended dose (50 mL), as previously discussed on the aggressiveness of the
242 native strains in the area, with the best performance of these strains in the absence of
243 nitrogen fertilization at sowing.

244




245
246
247
248

Fig 2. Number of stems per plant of the cv. IPR Tangará as a function of inoculant doses and nitrogen fertilization.

249 By adding 40 kg ha^{-1} of nitrogen at sowing, a 15.8% reduction was obtained in the number of
250 nodes, achieving the best results associated with the recommended inoculant dose (100 mL)
251 (Figure 2).

252
253
254
255
256
257
258
259
260
261

Regarding the number of pods per plant, significant differences were observed between treatments, with better results obtained in the absence of nitrogen fertilization (Figure 3). In this sense, Souza et al. [18] verified that seed inoculation only increased the number of pods per plant in the absence of N application. Nevertheless, in this variable, it is possible to observe that nitrogen fertilization with the aid of the recommended inoculant dose (100 mL) was able to reach values similar to the recommended inoculant dose without N in fertilization (Figure 3). Nevertheless, the cheapest form of fertilization, without N, with the influence of the inoculation, comprising half of the dose (50 mL), exceeded all other treatments.



262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290


Fig 3. Number of pods per cv. IPR Tangará as a function of inoculant doses and nitrogen fertilization.

Soratto et al. [19] and Alvarez et al. [20] observed positive results for the number of pods per plant when the N doses were increased. Araújo et al. [21] observed no effect of inoculation on the number of pods per plant. The number of pods per plant depends on the cultivar being used.

The number of grains per pod, despite being a characteristic of high genetic heritability [13], and thus related to the cultivar used, showed an interaction between the treatments with the two forms of fertilization and the different doses of the inoculant. In the treatment without nitrogen fertilization, the results were higher (Figure 4), in accordance with those found by Soratto et al. [22] which also did not have significant effects on the number of grains per pod with the use of different levels of N.

With the interaction between inoculant doses, it can be observed that half of the recommended dose (50 mL) was superior compared to the others, reaching 5.7 grains per pod (Figure 4). Similar results were found by Araújo et al. [21] and Romanini Júnior et al. [23], who noted that the inoculation with rhizobia contributed to the increase in productivity. Binotti [24] did not find significant differences for the number of grains per pod, using *R. tropici* in the inoculation of the bean seeds.

In the treatment that received nitrogen fertilization, the results were lower, but half of the inoculant dose again presented superior results compared to others in this treatment, reaching 5.1 grains per pod. A different result was found in a study carried out by Farinelli et al. [25], in the state of São Paulo, Brazil, evaluating the application of N doses in the bean crop, in a no-till system and in conventional planting, finding maximum yield with the application of 185 kg ha⁻¹ of N.



292 **Fig 4. Number of grains per pod of cv. IPR Tangará bean as a function of inoculant**
 293 **doses and nitrogen fertilization.**

295 4. CONCLUSION

296 Nitrogen fertilization at sowing and seed inoculation with *Rhizobium tropici* did not influence
 297 the plant population and the 1000-grain mass.

298 In the absence of nitrogen fertilization at sowing, pods with longer length and higher grain
 299 yield were obtained

300 Inoculation of the seeds with *Rhizobium tropici* exerts a positive influence on plant height,
 301 number of nodes of the main stem, number of pods per plant, and number of beans per pod,
 302 the dose recommended by the manufacturer (100 mL) being efficient, with the possibility of
 303 applying a dose of 50 mL, to satisfactory results.

308 CONSENT (WHERE EVER APPLICABLE)

309 Not applicable.

312 ETHICAL APPROVAL (WHERE EVER APPLICABLE)

313 Not applicable.

316 REFERENCES

- 317 1. National Supply Company. Follow-up of the Brazilian Grain Harvest. Safra 2017/18. 9^o
 318 Survey. Brasília: Conab. 2018; 1-178. Accessed 01 July 2018. Available:
 319 <https://www.conab.gov.br/>
- 320 2. War AF, Silva DB, Rodrigues GC. Management of irrigation and nitrogen fertilization for
 321 common bean in the Cerrado region. Pesquisa Agropecuária Brasileira. 2000; 35 (6): 1229-
 322 36. English. DOI: <http://dx.doi.org/10.1590/S0100-204X200000600020>.
- 323 3. Salgado FHM, Silva J, Oliveira TC, Barros HB, NG Steps, Fidelis RR. Efficiency of
 324 common bean genotypes in response to nitrogen fertilization. Tropical Agriculture Research.
 325 2012; 42 (4): 368-74. English. DOI: <http://dx.doi.org/10.1590/S1983-40632012000400007>.

- 332
333 4. Vieira RF, Lima MS, Neves JCL, Andrade MJB. Fertilizing. In: Carneiro JE, Paula Júnior
334 TJ, Borém A. editors. Beans from planting to harvest. Viçosa: Editora UFV; 2015.
335
- 336 5. Pelegrin R, Merchant FM, Otsubo IMN, Otsubo AA. Bean culture response to nitrogen
337 fertilization and rhizobium inoculation. Brazilian Journal of Soil Science. 2009; 33 (1): 2219-
338 26. English. DOI: <http://dx.doi.org/10.1590/S0100-06832009000100023>
339
- 340 6. Macdonald GK, Bennett EM, Potter PA, Rramankutty N. Agronomic phosphorus
341 imbalances across the world's croplands. Proceedings of the National Academy of Sciences
342 of the United States of America. 2011; 108 (7): 3086-91. DOI: 10.1073 / pnas.1010808108
343
- 344 7. Foley JA, Ramankutty N, Brauman KA, Cassidy ES, Gerber JS, Johnston M, Mueller ND,
345 O'connell C, Ray DK, West PC, Balzer C, Bennett EM, Carpenter SR, Hill J, Monfreda C,
346 Polasky S, Rockstrom J, Sheehan J, Siebert S, Tilman D, Zaks DPM. Solutions for a
347 cultivated planet. Nature. 2011; 478 (7369): 337-42. DOI: 10.1038 / nature10452
348
- 349 8. Rabelo ACR, Ribeiro DF, Rezende RM, Alcantara E, Freitas AS. Nitrogen fertilization on
350 bean crop. Journal of the Vale do Rio Verde University. 2017; 15 (1): 825-41. English.
351
- 352 9. Caviglione JH, Kilhl LRB, Caramori PH, Oliveira D. Climatic charts of Paraná. Londrina:
353 Iapar; 2000.
354
- 355 10. Santos HG, Almeida JA, Oliviera JB, Lumberras JF, Ljos Anjos, Coelho MR, Jacomine
356 PKT, Cunha TJF, Oliveira VA. Brazilian system of soil classification. 3rd ed. Brasília:
357 Embrapa; 2013.
358
- 359 11. Federal University of Santa Maria. The seed and its germination. Accessed 18 August
360 2018. Available: <http://coral.ufsm.br/sementes>.
361
- 362 12. Andrade MJB, Oliveira DP, Figueiredo MA, Martins FAD. Edafoclimatic requirements. In:
363 In: Carneiro JE, Paula Júnior TJ, Borém A. editors. Beans from planting to harvest. Viçosa:
364 Editora UFV; 2015.
365
- 366 13. Araújo JLS, Stradiotto R, Franco AA. Selection of common bean cultivars (*Phaseolus*
367 *vulgaris* L.) for biological fixation of nitrogen under high temperature conditions. In: 4th
368 National Bean Research Meeting. 1993. Proceedings. Londrina: Iapar; 136.
369
- 370 14. Ferreira AN, Arf O, Carvalho MAC, Araújo RS, Sá MS, Buzetti S. Rhizobium tropici
371 strains on bean inoculation. Scientia Agricola. 2000; 57 (3): 507-12. English. DOI:
372 <http://dx.doi.org/10.1590/S0103-90162000000300021>.
373
- 374 15. Peres JRR, Suhet AR, Mendes IC, Vargas MAT. Effect of inoculation with rhizobium and
375 nitrogen fertilization on seven bean cultivars on cerrado soil. Brazilian Journal of Soil
376 Science. 1994; 18 (3): 415-20. English.
377
- 378 16. Matoso SCG, Kusdra JF. Nodulation and bean growth in response to the application of
379 molybdenum and rhizobial inoculant. Brazilian Journal of Agricultural and Environmental
380 Engineering. 2014; 18 (6): 567-73. English. DOI: <http://dx.doi.org/10.1590/S1415-43662014000600001>.
381
382

- 383 17. New MCSS, Tanaka RT, Mascarenhas HAA. Nitrogen and potassium in symbiotic N
384 fixation by soybean cultivated in winter. *Scientia Agricola*. 1999; 56 (1): 143-56. English.
385 DOI: <http://dx.doi.org/10.1590/S0103-90161999000100021>.
386
- 387 18. Souza EFC, Soratto RP, Pagani FA. Nitrogen application and inoculation with rhizobia in
388 common bean cultivated after maize intercropped with brachiaria. *Pesquisa Agropecuária*
389 *Brasileira*. 2011; 11 (4): 370-77. English. DOI: [http://dx.doi.org/10.1590/S0100-](http://dx.doi.org/10.1590/S0100-204X2011000400005)
390 [204X2011000400005](http://dx.doi.org/10.1590/S0100-204X2011000400005).
391
- 392 19. Soratto RP, Silva TRB, Arf O, Carvalho MAC. Levels and times of application of nitrogen
393 in cover in common bean irrigated in no-tillage. *Agronomic Culture*. 2001; 10 (1): 89-99.
394 English.
395
- 396 20. Alvarez ACC, Arf O, Alvarez RCF, Pereira JCR. Response of the bean to the application
397 of doses and sources of nitrogen under cover in the no-tillage system. *Acta Scientiarum*
398 *Agronomy*. 2005; 27 (1): 69-75. English. DOI:
399 <http://dx.doi.org/10.4025/actasciagron.v27i1.1927>.
400
- 401 21. Araujo FF, Carmona FG, Tiritan CS, Creste JE. Biological fixation of N in common bean
402 submitted to inoculant dosages and chemical treatment in the seed compared to nitrogen
403 fertilization. *Acta Scientiarum Agronomy*. 2007; 29 (4): 535-40. English. DOI:
404 <http://dx.doi.org/10.4025/actasciagron.v29i4.416>
405
- 406 22. Soratto RP, Carvalho MAC, Arf O. Chlorophyll content and yield of common bean due to
407 nitrogen fertilization. *Pesquisa Agropecuária Brasileira*. 2004; 39 (9): 895-901. English. DOI:
408 <http://dx.doi.org/10.1590/S0100-204X2004000900009>.
409
- 410 23. Romanini Junior A, Arf O, Binotti FFS, Sá ME, Buzetti S, Fernandes FA. Evaluation of
411 rhizobium inoculation and nitrogen fertilization in common bean development under no -
412 tillage system. *Bioscience Journal*. 2007; 23 (4): 74-82. English.
413
- 414 24. Binotti FFS. Nitrogen management in winter bean in succession to corn and Brachiaria
415 under no - tillage system. 2009; 178. Thesis (Doctorate in Agronomy / Production Systems) -
416 Faculty of Engineering. State University of São Paulo, Ilha Solteira; 2009.
417
- 418 25. Farinelli R, Lemos LB, Penariol FG, Egéa MM, Gasparoto MG. Coverage nitrogen
419 fertilization in common bean and no - tillage. *Pesquisa Agropecuária Brasileira*. 2006; 41 (2):
420 307-12. English. DOI: <http://dx.doi.org/10.1590/S0100-204X2006000200016>.
421