

***Henosepilachna vigintioctopunctata* (Coleoptera: Coccinellidae: Epilachninae) attacking cultivated and wild crops in Brazil**

ABSTRACT

Henosepilachna vigintioctopunctata was recorded on *Solanum melongena* L. for the first time in cultivated crops areas in Espírito Santo State, southeastern Brazil, in the first semester of 2017. In an one-year research, *H. vigintioctopunctata* was recorded in eight plants, host of the families *Amaranthaceae* and *Solanaceae*. These new records represent a high potential establishment of this pest in Brazil.

Key words: 28-spot lady bird, *Solanum melongena*, Phytophagy, Plant Protection.

1. INTRODUCTION

Henosepilachna vigintioctopunctata (Fabricius, 1775) (Coleoptera: Coccinellidae: Epilachini), the 28-spot lady bird or hadda beetle has been reported as one of the most important pest of cultivated and wild Solanaceae and Cucurbitaceae plants in Asia, with records as pest in China [1], India [2-4], Indonesia [5,6], Japan [7,8] and Pakistan [9] and in Oceania, with records in Australia [10]. Cultivated plants in these regions include eggplants, potatoes, tomatoes, tobacco and cucumbers.

In Western Hemisphere, the first record of *H. vigintioctopunctata* was made in 1990 in Curitiba, Paraná State, Brazil, in an unidentified wild cucurbit. In 1991, in Paranaguá, on *Piper nigrum* L. (Piperaceae) and in 1992, in Itajaí, Santa Catarina State, on *Solanum americanum* Mill. (Solanaceae) [11]. On the three mentioned records only adults were reported. *Henosepilachna vigintioctopunctata* was probably introduced from Port of Paranaguá, Paraná State (L.M. Almeida, personal communication).

In 2002, adults and larvae were over again collected in *S. americanum* in Itajaí. In 2010, all development stages of *H. vigintioctopunctata* were collected on *Brugmansis suaveoleus* (Humb. and Bonpl. ex Willd.) (Solanaceae) in Campinas and São Paulo, São Paulo State [12].

During field researches at Instituto Federal do Espírito Santo, Campus Itapina, in Colatina, Espírito Santo State, in the first semester of 2017, adults and larvae were observed attacking eggplants *Solanum melongena* L. (Solanaceae). New records were made through one year in different host plants. Adults and larvae collected in July, 2018, in Colatina, in eggplants were sent to Dra. Lucia Massutti Almeida, Department of Zoology, Federal University of Paraná (UFPR), Curitiba, Brazil, to species identification. Vouchers specimens were deposited in Entomology Collection Prof. Dr. Pe. Jesus Santiago Moure, UFPR.

In a one-year survey, *H. vigintioctopunctata* was recorded in eight plants, host of the families *Amaranthaceae* and *Solanaceae*, five of them with all development stages of the pest (Table 1).

Table 1. Host plant and development stages of *Henosepilachna vigintioctopunctata* found in Espirito Santo, Brazil, July 2017-Agost 2018.

Botanical family	Host plant	Development stage found			
		Egg	Larva	Pupa	Adult
Amaranthaceae	<i>Amaranthus viridis</i> L.	A	A	A	P
Solanaceae	<i>Brugmansia suaveolens</i> (Willd.)	P	P	P	P
	<i>Physalis angulata</i> L.	P	P	P	P
	<i>Solanum aethiopicum</i> L.	P	P	P	P
	<i>Solanum lycopersicum</i> L.	A	A	A	P
	<i>Solanum melongena</i> L.	P	P	P	P
	<i>Solanum nigrum</i> L.	P	P	P	P
	<i>Solanum paniculatum</i> L.	A	A	A	P

P – present, A – absent.

Attacked leaves showed typical symptoms mentioned to *H. vigintioctopunctata*, characterized by a network aspect due to scraped occasioned by larval and adults (Fig. 1.). According to [13], in *S. melongena* leaves, larvae feed on phloem, epidermal and parenchymal tissues, while adults scraped upper and lower sides of the leaves. Fruit reduction up to 60% in eggplants field has been reported in India [14].




Fig. 1. Attacked plants for *Henosepilachna vigintioctopunctata*: *Solanum melongena* L. (A), *Solanum aethiopicum* L. (B), *Physalis angulata* L. (C) and *Solanum nigrum* L. (D) in Espirito Santo, Brazil, July 2017-Agost 2018.

2. Conclusion

Since its first record, *H. vigintioctopunctata* has not been mentioned in the list of quarantine pests of the Ministry of Agriculture. The new records of this present study, up to 1000 km from the first occurrence in Brazil, in agriculture areas and in eight hosts, represents a high potential of establishment of this pest in Brazil. The Brazilian Association of Seed and Seedling Trade (ABCSEM) estimated 820 thousand hectares of vegetable crop in 2016. Eggplant production has increased in Brazil in last year mainly due to its medicinal importance. Thus, efforts to avoid the spread of *H. vigintioctopunctata* should be made.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of

the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

References

1. Zhou L, Xie BG, Wang XP Population dynamic of *Henosepilachna vigintioctopunctata* in different host plants in Jiangnan plain. *China J. Northern Hort.* 2015;11:103–105 (in Chinese).
2. Rajagopal D, Trivedi TP . Status bioecology and management of *Epilachna beetle*, *Epilachna vigintioctopunctata* (Fabricius) (Coleoptera: Coccinellidae) on potato in India: A review. *Trop. Pest Management.* 1989;35(4):410-413.
3. Bhagat KC, Munshi SK. Host preference of spotted leaf eating beetle, *Henosepilachna vigintioctopunctata* (Fabricius) on brinjal varieties. *Pest Management and Economic Zoology.* 2004;12:77-81.
4. Jamwal VVS, Ahmad H, Sharma D. Host biology interactions of *Epilachna vigintioctopunctata* (Fabricius). *The Bioscan.* 2013;8(2):513-517.
5. Nakamura K, Abbas I, Hasyim A. Population dynamics of the phytophagous lady beetle *Epilachna vigintioctopunctata* in an eggplant field in Sumatra. *Researches on Population Ecology.* 1988; 30:25-41.
6. Katakura H, Nakano S, Kahono S, Aabbas I, Nakamura N. Epilachnine Ladybird Beetles (Coleoptera: Coccinellidae) of Sumatra and Java, Tropics. 2001;10(3): 325-352.
7. Hirano K. Study on the movements of the 28-spotted lady-beetle *Henosepilachna vigintioctopunctata* by the mark-recapture method. *Applied Entomology and Zoology.* 1985;29:7–13 (in Japanese).
8. Hirano K. Population Dynamics of a phytophagous Lady-beetle, *Epilachna vigintioctopunctata* (Fabricius) (Coleoptera: Coccinellidae), Living in Spatio-temporally Heterogeneous Habitats: II. Seasonal Changes in Adult Movement and Spatial Distribution of the Population. *Applied Entomology and Zoology.* 1993;28(2):131.
9. Naz F, Inayatullah M, Rafi MA, Ashfaq M, Ali A. *Henosepilachna vigintioctopunctata* (Fabricius) (Epilachninae: Coccinellidae) its taxonomy, distribution and host plants in Pakistan. *Sarhad Journal of Agriculture.* 2012; 28(3):421-427.
10. Richards AM. The *Epilachna Vigintioctopunctata* Complex (Coleoptera: Coccinellidae). *International Journal of Entomology.* 1983;25:11-41.
11. Schroder RFW, Athanas MM, Pavan C. *Epilachna vigintioctopunctata* (Coleoptera: Coccinellidae), new record for Western Hemisphere, with a review of host plants. *Entomological News.* 1993;104(2):111-112.
12. Casari SA, Teixeira EP. Immatures of *Epilachna chevrolat* (Coleoptera: Coccinellidae, Epilachninae). *Revista Brasileira de Entomologia.* 2015; 59:113-120.
13. Mohanchander, Ponnuvel et al. Effect of Hadda beetle, *Epilachna vigintioctopunctata* Fab. (Coleoptera: Coccinellidae) infestation on eggplant leaf (*Solanum melongena* L.) and bio-control potential of essential oil formulations. *Pakistan Journal of Biological Sciences.* 2013;16:991-997.
14. Mall NP, Pandey RS, Singh SV, Singh SK. Seasonal incidence of insect pests and estimation of the losses caused by shoot-and-fruit borer on brinjal. *Indian Journal of Entomology.* 1992; 54(3):241-247.