

1 **Obstructive Sialadenitis of Submandibular Gland Due to a Nail-Like Fish**
2 **Bone Foreign Body: A Rare Case Report**

3 **Running title:** Obstructive sialadenitis of submandibular gland
4

5 **Abstract**

6 **Introduction:**

7 Foreign body-associated sialadenitis of submandibular gland is not often and scarce within the
8 literatures. In this study, a report of piece of Nail-like fish bone foreign body entering the
9 Wharton’s duct causing an acute sialadenitis is presented.

10 Foreign bodies must be explored and all suspected areas must be examined carefully for
11 avoiding secondary problems and surgeries in the future.

12 Foreign bodies in the oral and maxillofacial region are often experienced after trauma and
13 dental treatment.

14 **Case report:**

15 We describe a case of obstructive sialadenitis in the submandibular gland caused by penetration
16 of a fish bone in a 68-year-old man. Hhe had swelling and spontaneous pain in the left
17 submandibular region. Radiographic examination didn’t show foreign body in the
18 submandibular gland. Initially we diagnosed obstructive sialadenitis in the left submandibular
19 gland and we guess something like salivary stone may be the cause of this swelling so
20 compressing and milking of Wharton duct was done and suddenly a tip of foreign body was
21 appeared. The foreign body measured 1.3cm *3 mm*2mm and was a nail-like object. On
22 pathological examination, the foreign body was found to be a fish bone (cartilage-like organic
23 material).

24 **Conclusion**

25 This case demonstrated that precise and proper inspection and examination, milking and then
26 pay attention to secretion of salivary gland lead to proper diagnosis and after that suitable
27 treatment, so this could reduce costly assessment and treatment, also lessen bewilderment of the
28 patient.

29 **Keywords:** Obstructive sialadenitis, foreign body, foreign body-associated sialadenitis,
30 submandibular gland, nail-like fish bone

31
32 **1. Introduction**

33 Obstructive sialadenitis of the submandibular gland is usually due to sialoliths, but foreign body-
34 associated sialadenitis is not often and rare in the literatures [1–14]. The mechanism of foreign-

35 body entry is generally traumatic [4-6, 15], reports of a foreign body entering the salivary gland
36 intraorally through Wharton's or Stensen's duct are not common and usual. [4, 5, 7, 16]. Foreign
37 body-induced sialoliths are even more rare [6, 8, 9, 17, 18]. As we know Sialoendoscopy ,is one
38 of minimally invasive procedure, that has recently been applied for direct diagnosis of pathologic
39 features in the ductal system and removal of sialoliths and foreign bodies in the duct of the
40 salivary gland but it use for distal of the salivary's duct [4-6,12,19].

41 Here we report a patient has cured with milking and compressing Wharton's duct and suddenly
42 the foreign body like fish bone-induced sialoliths came out of the duct and after that prescribing
43 antibiotic.

44

45 **2. Case Report**

46 A 68-year-old male patient was referred to Department of Oral and maxillofacial Medicine and
47 surgery with a chief complaint of a swelling in left side of neck since 14 days ago and he was
48 suffering pain since 2 weeks ago, by the way Pain was increased in intensity while swallowing
49 and eating meals. (Figure 1A, 1B)Patient gave history of fever and malaise, difficulty in eating
50 and also speaking. He expressed that the swelling was small in size and immediately increase to
51 present size of 5-6 cm.

52 The patient also suffered from Diabetes ,Hypertension and he had history of Cardiac
53 Arrhythmia. He consulted with his dentist and get antibiotic (Cap Amoxicillin 500 mg) every 8
54 hour and also panoramic view radiography was taken but the dentist couldn't find the cause
55 (figure 2A). For a short duration partial recovery was done but after that the swelling was
56 recurred.

57 Clinical examination of intraoral revealed that ovoid shape swelling in floor of the mouth and it
58 measured 5-6 cm in diameter. Extra oral findings reveal that enlargement of lateral neck
59 extended from lower border of mandible (5 cm anterior to angle of mandible) to lateral upper
60 border of thyroid cartilage. (Figure 1A) The border of enlargement was well-defined and regular
61 border, surface was smooth and skin over the swelling was intact like adjacent tissues. It was
62 tender on palpation but temperature was not raised. Consistency of swelling was soft and rubbery
63 and fluctuation was present but it was not fixed to overlying skin (Figure 1a, 1b).

64 Intra oral examination showed swelling of Wharton's duct in left floor of the mouth (Figure 2b).
65 it was tender and painful on palpation, and consistency of swelling was soft and fluctuant.

66 It is important to formulate the differential diagnosis when swelling and mass is seen at the side
67 of neck since this would help further evaluation of the condition and management of the patient.
68 After considering all clinical findings following entities were considered in differential
69 diagnosis—acute submandibular sialadenitis and benign swelling of neck.

70




a

b

71 **Figure 1:** Swelling of submandibular region that extended to lateral neck, and the patient had
72 tenderness on palpation (a). Asymmetry and swelling of left submandibular region, frontal view
73 of patient (b).

74 Compressing and Milking of Wharton duct was performed and suddenly the nail-like fish bone
75 foreign body came out of the duct and after that the pus were pushed out and we waited for about
76 15 minutes to get out of whole pus then we rinsed the orifice and duct (Figure 3a,3b, Figure 4a)
77 High dose of oral antibiotic; cap Amoxicillin (2 g) every 6 hours was prescribed for one day
78 and then it was tapered until one week after culture test .We emphasize use antibiotics 2 hours
79 before eating meals for better Gastrointestinal absorption. Also use of adequate hydration and
80 sialagogues, pure honey as a mouth rinse for 3-4 time a day, was advised to him.

81 The recalled sessions for follow-up was 3 days and 1 week later. Significant improvement was
82 achieved. Foreign body sent for histopathological examination. The report of biopsy was
83 interpreted as a fishbone (Figure 4b, Figure 5a, 5b).Final diagnosis of obstructive submandibular
84 sialadenitis was given. There is no residual or recurrent swelling apparent in the area of
85 intervention after a follow-up period of 6 months.



a



b

86 **Figure 2:** (close view) Panoramic view of patient Without illustrating opacity of foreign body in left
87 submandibular duct (a). Photography of swelling of orifice of Wharton's duct in left floor of the mouth
88 (compare with right floor of the mouth) (b).



a




b

89 **Figure 3:** photography (a) illustrating tip of the foreign body in orifice of Wharton's duct that appeared
90 after milking of this orifice .Close view of Nail-like foreign body (bone fish) that different derbies
91 Surrounded it around (b)

92



93 **Figure 4)** Photography from floor of the mouth illustrating pushing out of pus from orifice of Wharton's
 94 duct that appeared after removing the foreign body (a). Improvement of swelling of the floor of the mouth
 95 after removal of foreign body and treatment with antibiotic for a period of 2 weeks (b).



96 **Figure 5:** The report of biopsy was interpreted as foreign body (fish bone) (a). Microscopic view of
 97 foreign body (fish bone) (b)

98

99 **3. Discussion**

100 A search of Medline using the key words foreign body, fish bone and submandibular gland
 101 revealed first case published on 1990[10].

102 Many different kind of things as foreign bodies have been found and reported in salivary gland
 103 such as paper clips, feathers ,toothbrush bristles, spikes of wheat, blades of grass, hairs ,pencil
 104 lead, plastic pen tops, plant material, splinters of wood, pieces of metal, slivers of fingernail, and
 105 fish bones [4,6,7,10,11,13,16].

106 2 reasons and hypotheses considered for entering foreign bodies into the salivary gland: one of
107 them is penetrating trauma [4, 6, and 20] and the other reason is retrograde migration [7, 13, and
108 21]

109 Perhaps In our case because of partial edentulous and inability to complete mastication and also
110 lessening of salivary gland's secretion due to presumption of drugs, contingency of entering
111 foreign body and retrograde migration will be increased.

112 When traumatic injury is a cause and history of obstructive sialadenitis so diagnosis is relatively
113 easy [6]. However, in cases without traumatic injury, it is obscure whether the cause is retrograde
114 migration for these 4 reasons: (1) there is almost steady salivary flow; (2) the orifice of duct is
115 mobile and can twist in all directions; (3) the diameter of the duct at the orifice is miniature; and
116 (4) in most submandibular glands, there is a sphincter-like system in the first 3 cm of Wharton's
117 duct that prevents the retrograde migration of substances [17,21,22,23]. Findings support the
118 possibility that some sialoliths might result from retrograde migration of a fish bone through the
119 orifice of Wharton's duct. Fish bones are one of the most common foreign bodies found in the
120 pharynx and esophagus, but they are very rarely found in the salivary gland or the duct [4-10, 13,
121 14, 24, and 25]. In previous studies, the incidence of a fish bone encompassed by a sialolith
122 ranged from 2.8% (12/423) to 4.4% (5/114) of patients with sialoliths of the submandibular
123 gland [6, 8, and 26].

124 There are interesting statistics about fish bone as a foreign body, for example, in one of review of
125 English-language literature, migration of a fish bone into the salivary gland was more common
126 in men than in women [4-10, 13, 26]. Fish bones were more often present in the submandibular
127 gland than the parotid gland [4-10,13,14,26] and were more often located in the left side of the
128 submandibular gland than in the right side [4-10,13,14,26]. Stone formation induced by a fish
129 bone tended to be obviously related to occupation (fisherman), dietary habit (seafood), and
130 history of injury (recollection of a fish-bone injury and subsequent symptoms) [6]. Fish bone-
131 induced sialoliths were previously reported to be 3–18 mm in size [5-8, 10, 26].

132 Our case was matched to this literature, in our male patient, the nail-like fish bone foreign body
133 was a total of 1.2 cm in length and 0.3 cm in diameter in the left submandibular duct.

134 In evaluating the patient with sialadenitis, these steps should be taken in the following order:

135 1. History, 2. physical examination, 3. culture, 4. Laboratory investigation, 5. Radiography, and if
136 indicated, 6. fine-needle aspiration biopsy.

137 There are wide range of approaches for management and treatment of sialadenitis, these include
138 conservative medical management to more invasive surgical intervention.

139 One management scheme is as follows:

140 • Acute sialadenitis –

141 Medical management (hydration, antibiotics [oral versus parenteral], warm compresses and
142 massage, sialagogues);

143 surgical management (consideration of incision and drainage versus excision of the gland in
144 cases refractory to antibiotics, incision and drainage with abscess formation, gland excision in
145 cases of recurrent acute sialadenitis)

146 Conservative therapies for acute management of obstructive sialadenitis, include: hydration,
147 analgesia (NSAIDs), sialagogues to stimulate salivary secretion, and regular, gentle gland
148 massage.

149 As we know the first- line therapy for stones in distal ducts of salivary glands is interventional
150 sialadenoscopy, also we can use this approach for removal of foreign bodies.

151 By the way, if infection is present, empiric antibiotic therapy should be given after proper
152 cultures have been obtained.

153 In our case, because of the improvement of sign and symptoms, we continued the same antibiotic
154 that we prescribed before for our patient .By the way with removal of fish bone foreign body,
155 most of the pus was pushed out. As we know the foreign body was the main cause.

156

157 **4. Conclusion**

158 This case demonstrated that precise and proper inspection and examination lead to proper diagnosis and
159 after that suitable treatment, so this could reduce costly and expensive assessment and treatment, also
160 lessen bewilderment of the patient.

161 Another important matter, Patients with any form of sialadenitis should be educated as to the worthiness
162 of hydration and excellent oral hygiene.

163 At the end, Milking and pay attention to transparency (glassiness) and canescent of secretion of salivary
164 gland are helpful for achievement of proper diagnosis.

165 **References**

166 1. Taneja M, Taneja MK. Foreign body Wharton's duct. Indian J Otolaryngol Head Neck Surg 2011;
167 63:300–301.

168 2. Ozturk, Kayhan MD; Erdur, Omer MD; Aksoy, Ceren MD. Foreign Body of Submandibular Gland.
169 Journal of Craniofacial Surgery: October 2016 - Volume 27 - Issue 7 - p e600–e601.

170 3. Shameeka Thopte, Shams Ul Nisa, Abhijeet Jadhav, Rohan Chaudhari. Sialolithiasis Of
171 Submandibular Gland With Acute Suppurative Sialadenitis: A Case Report. World Journal Of Pharmacy
172 And Pharmaceutical Sciences 5(4) · March 2016

173 4. Gill AS, Kieliszak C, Joshi AS. Sialendoscopy as a management tool in patients with foreign body
174 impaction of the salivary gland. Am J Otolaryngol 2016; 37:369–71.

175 5. Yamano Y, Uzawa K, Ito H, Tanzawa H. Endoscopically assisted removal of a fish bone penetrating
176 the parotid duct: an unusual case. J Oral Maxillofac Surg 2014; 72:1343–9.

177 6. Xie L, Zheng L, Yu C, Yang C, Chen Z, Yun B, et al. Foreign body induced sialolithiasis treated by
178 sialoendoscopic intervention. J Craniofac Surg 2014; 25:1372–5.

- 179 7. Su YX, Lao XM, Zheng GS, Liang LZ, Huang XH, Liao GQ. Sialoendoscopic management of
180 submandibular gland obstruction caused by intraglandular foreign body. *Oral Surg Oral Med Oral Pathol*
181 *Oral Radiol* 2012; 114:e17–21.
- 182 8. Yu C, Yang C, Zheng L. Sialendoscopic findings in patients with obstructive sialadenitis: long-term
183 experience. *Br J Oral Maxillofac Surg* 2013; 51:337–41.
- 184 9. Sato K, Umeno H. Clinical photographs. Fish bone-induced sialolith. *Otolaryngol Head Neck Surg*
185 2009; 141:539–40.
- 186 10. Abe K, Higuchi T, Kubo S, Oka M. Submandibular sialoadenitis due to a foreign body. *Br J Oral*
187 *Maxillofac Surg* 1990; 28:50–2.
- 188 11. Ozturk K, Erdur O, Aksoy C. Foreign body of submandibular gland. *J Craniofac Surg* 2016; 27:e600–
189 601.
- 190 12. Ardekian L, Klain H, Peled M. Obstructive sialadenitis of submandib-ular gland due to foreign body
191 successfully treated by sialoendoscopic intervention. *J Oral Maxillofac Surg* 2009; 67:1337–9.
- 192 13. Derin S, Sahan M, Kule M, Koseoglu S, Celik OI. Fish bone induced sialolith in Warthon duct. *J*
193 *Craniofac Surg* 2015; 26:e663–664.
- 194 14. Matsuo T. Acute suppurative parotitis caused by a fish bone: a case report. *Int J Oral Maxillofac Surg*
195 1997; 26:54.
- 196 15. Amarbir S.GillBS^aChristopher R.KieliszakDO^bArjun S. Sialendoscopy as a management tool in
197 patients with foreign body impaction of the salivary gland. *American Journal of Otolaryngology*. Volume
198 37, Issue 4, July–August 2016, Pages 369-371.
- 199 16. Sivapatha Sundaram Sreetharan , Rajan Philip.Unusual Foreign Body of Parotid Gland Presenting as
200 Sialolithiasis: Case Report and Literature Review. *Case Reports in Otolaryngology*. Volume 2012 (2012),
201 Article ID 367349, 3 pages.
- 202 17. Marchal F, Kurt AM, Dulguerov P, Lehmann W. Retrograde theory in sialolithiasis formation. *Arch*
203 *Otolaryngol Head Neck Surg* 2001; 127:66–8.
- 204 18.Yasufumi KOSUGI,Toshinori Iwai, Shinsuke OHTA,Iwai TOHNAI.A case of an endoscopically
205 removed parotid duct sialolith. *Nippon Koku Geka Gakkai zasshi*· March 2017; 63(3):153-157.
- 206 19.Maria E.PapadakiDMD, MD,Joseph P.McCainDMD,KingKimDMD,Ronald L.KatzDMD&Leonard
207 B.KabanDMD, MD||Maria J.TroulisDMD, MSc .Interventional Sialoendoscopy: Early Clinical
208 Results.*Journal of Oral and Maxillofacial Surgery*. Volume 66, Issue 5, May 2008, Pages 954-962.
- 209 20. P Capaccio, Torretta S, Ottaviani F,et al.Modern management of obstructive salivary diseases. *Acta*
210 *Otorhinolaryngol Ital*. 2007 Aug; 27(4): 161–172.
- 211 21. Yu-xiongSuMD, DDS^aXiao-meiLaoBDS, DDS^bGuang-senZheng Li-zhongLiangDDS^bXing-
212 huaHuangBDS^bGui-QingLiaoMD, DDS^cSialoendoscopic management of submandibular gland
213 obstruction caused by intraglandular foreign body. *Oral Surgery, Oral Medicine, Oral Pathology and Oral*
214 *Radiology* Volume 114, Issue 5, November 2012, Pages e17-e21.
- 215 22. F. Marchal, A.M. Kurt, P. Dulguerov, W. Lehmann. Retrograde theory in sialolithiasis formation
216 .*Arch Otolaryngol Head Neck Surg*, 127 (2001), pp. 66-68.

- 217 23. Mahabaleshwara C.H, Jayadeep Nidyalmale, Abhishek P.T, Ashoka G. 'FISH BONE': THE
218 REASON BEHIND SUBMANDIBULAR SIALADENTITS - A UNIQUE CASE REPORT. International
219 Journal of Clinical And Diagnostic Research ISSN 2395-3403 Volume 5, Issue 3, May-June 2017. *Intl. J.*
220 *Clin. Diag. Res.* 2017; 5(3): 1.
- 221 24. Ikenberry SO, Jue TL, Anderson MA, et al; Management of ingested foreign bodies and food
222 impactions. *Gastrointest Endosc.* 2011 Jun73 (6):1085-91. doi: 10.1016/j.gie.2010.11.010.
- 223 25. Marc H Hohman, Wayne J Harsha, K Linnea Peterson. Migration of Ingested Foreign Bodies into the
224 Thyroid Gland: Literature Review and Case Report. *The Annals of otology, rhinology, and laryngology*
225 February 2010.119(2):93-8 .
- 226 26. Yu C, Yang C, Zheng L, Wu D. Endoscopic observation and strategic management of obstructive
227 submandibular sialadenitis. *J Oral Max-illofac Surg* 2010; 68:1770–5.
- 228