

1 **Effect of some Botanical Extracts on the management of *Meloidogyne incognita* and on**
2 **Growth of Tomato**

3 **ABSTRACT**

4 Tomato (*Solanum lycopersicum* L.) is an important and widely grown vegetable crop all over the
5 world. Although tomato is nutritionally and economically important, its production is
6 constrained by biotic and abiotic constraints leading to poor marketable quantity and quality
7 worldwide. Nematodes have one of the major pests affecting tomato production worldwide,
8 especially, in the tropical and sub-tropical regions. Screen house experiments were laid out in
9 Complete Block Design (CBD) with a 3x7 factorial arrangement replicated three times carried
10 out at the Department of Crop and Environmental Protection, University of Agriculture, Makurdi
11 screen house. The soil was sterilized before the experiment. Fresh leaves and seeds of Moringa,
12 Castor bean and Jatropha were washed with tap water, 15 g from each of leaves and seeds of the
13 different botanicals was macerated separately in an electric blender at high speed for 4 minutes
14 in 100ml distilled water. The mixtures were passed through a Whatman filter paper; the filtrates
15 of the leaves/seeds were then collected. Three tomato varieties viz: Roma Vf, Rio Grande and
16 UC82B were inoculated with approximately 5,000 freshly hatched *Meloidogyne incognita*
17 juveniles, two weeks after transplanting. Thirty percent 30%v/v aqueous extract of each of
18 Castor, Moringa and Jatropha leaves and seeds was used, while distilled water (0%) served as the
19 control. Thirty ml 30ml of various each leaf and seed aqueous extract was applied, 48 hours
20 after inoculation as soil drench. Application was done at 1weeks intervals thereafter for a period
21 of 16weeks. Data collected include number of fruits per plant, root gall index, nematode
22 reproductive factor, and final nematode population. The results showed that various leaves and
23 seed extracts significantly (P<0.05) reduced root gall index, final population of *M. incognita* in
24 the soil and nematode reproductive factor than the control. Application of the various treatments
25 led to significant increase in mean number of fruits and mean fruit weight yield of all the three
26 tomato varieties. Therefore, the application of leaf and seed aqueous extracts of Moringa,
27 Jatropha and Castor will serve as good alternative for the management of root knot nematode
28 population.

29 Keywords: Tomato, Botanical extracts, *M. incognita*

30
31
32 **INTRODUCTION**

33 **Tomato fruit is an important source of minerals, essential amino acids, sugars, dietary fibers,**
34 **and vitamins as it contains vitamin A as carotene, vitamin B1 (thiamin), B2(riboflavin), niacin**
35 **and vitamin C (1). Tomato seed contains 24 percent edible oil, used for manufacturing of**
36 **salad creams, margarine and soap, the residual pressed cake is used for fertilizer and**
37 **livestock feeding (2). It is also found to have medicinal value as it is important in weight**
38 **management and reduction of cancer risk (3).**

39 Although tomato is nutritionally and economically important, its production is constrained by
40 biotic and abiotic factors leading to poor marketable quantity and quality worldwide (27). (4)
41 Identified nematodes as one of the major pests affecting tomato production worldwide,
42 especially, in the tropical and sub-tropical regions. (5) stated that plant parasitic nematodes
43 attack seedling roots after emergence **there** by affecting crop productivity. (6) reported that a
44 lot of plant species such as tomato, amaranth, soy bean, yam tubers, cassava, maize, rice e.t.c
45 are prone to attack by the root-knot nematode.

46 The need for farmers to adopt strategies that do not pollute the environment has increased
47 urgency in the search for alternative sustainable methods to manage nematodes (7; 8 and 9).
48 Alternatively, research has focused on antagonistic plants (10, 26). These compounds can be
49 developed for use as nematicides or they can serve as model compounds for the
50 development of derivatives with enhanced activity or environmental friendliness (11). The
51 objective of the study was to determine the effect of botanical extracts from leaves and seeds
52 of Moringa, Jatropha and castor on root-knot nematode **infecting** three Tomato varieties
53
54

55 MATERIALS AND METHODS

57 Experimental Layout and Location

58 The experiment was carried out at the Department of Crop and Environmental Protection Laboratory
59 University of Agriculture, Makurdi, laid out in 3x7 factorial arrangement laid out in Completely
60 Randomized Design (CRD) replicated three times

61 Soil Sterilization

62 The Soil for the pot experiment was sterilized using the steam sterilization method, **using by** a metal
63 barrel steam sterilizer. The steam sterilizer has two chambers, the lower chamber contained water
64 and the upper part **contained** the soil. The soil was covered with wet jute sacks to conserve steam in
65 the chamber. **Fired** wood was used as the source of heat. The soil was sterilized for 30 minutes at 100°
66 C; it was then left for 48hrs before **using**.

67 Nursery Preparations and Agronomic Practice

68 Tomato seedlings were raised in three different buckets containing sterilized soils for the
69 three different varieties, the seeds of the various varieties were spread and soil **was** lightly
70 poured on **it the seeds** about two inches was used to cover the seeds. Mulch was provided to
71 protect the seeds from excessive sunlight and to serve as moisture conserver. Watering of the
72 young seedlings was done daily until when the seedlings were ready for transplanting.

73 Transplanting and Inoculation of Seedlings

74 Tomato seedlings were transplanted into 72 plastic buckets. Seven kg of sterilized soil was
75 put into each plastic buckets. The seedlings were transplanted, four weeks after germination.
76 Each of the tomato stands contained in the pots was inoculated with approximately 5,000
77 root-knot nematode (*M. incognita*) juveniles. Inoculation took place, one week after
78 transplanting with the use of a syringe and by pulling away the soil around the roots 2 cm
79 deep and 3 cm from the root. The juveniles were inoculated into the hole and the soil was
80 gently covered. Each bucket contained one seedling.

81 Source and Preparation of Plant Aqueous Extracts

82 Seeds/leaves of Moringa, Castor and Jatropha were obtained from the University Research
83 Farm.

84 Fresh leaves and seeds of Moringa, Castor bean and Jatropha were washed with tap water.
85 Fifteen grams from each of leaves and seeds of the different plant (Castor, Jatropha and
86 Moringa) was macerated separately in an electric blender at high speed for 4 minutes in
87 100ml distilled water. The mixture was left for 12 hours (overnight). Each mixture was passed
88 through a Whatman filter paper. Filtrates of the leaves/seed collected served as standard
89 solution 'S' for the experiment

90

91 Application of Treatments

92 The stands were treated with 30ml of each of various leaves and seeds aqueous extracts, 48
93 hours after inoculation. The untreated seedlings/stand served as the controls. Application of
94 the botanical extracts was done at weekly intervals thereafter until harvest. What is number
95 of replicates for each treatment?

96

97 Harvesting of tomato plants

98 The tested plants were harvested when fully matured (16 weeks). To ensure easy removal of
99 the plants from the soil, the sides of the plastic pots was pressed in order to loosen the soil.
100 The soil was then removed from the roots by gently shaking the plants.

101

102

103 Data Analysis

104 Data collected were analyzed, using the Genstat statistical package (Discovery edition
105 7). Least significant difference (LSD) at 5% was used for comparing mean differences.

106

107

108

109

110

111

RESULTS

112

113

114

115

116

117

118

119

120

121

122

123

124

Table 1 shows no significant ($P \leq 0.05$) differences among the varieties on root gall index. However, there are differences ($P < 0.05$) between among the varieties in nematode final populations and reproductive factor, Roma VF recorded the highest final population and reproductive factor of 2228 $J_2/200g$ of soil and 0.45, while Rio Grande recorded the least population and reproductive factor of 1461 $J_2/200g$ and 0.29, respectively.

On the mean number of fruits, there were no significant differences ($P > 0.05$). However, differences ($P < 0.05$) were observed on the mean fruit weight (kg/ha) with Rio Grande yielding, higher fruit weight of 39.63 closely followed by UC28B with 36.87, while the least was observed from Roma VF with 34.95.

There were significant differences ($P < 0.05$) between among the botanical extracts on root-knot nematode parameters and yield parameters (Table 2). Jatropha seed recorded the lowest root gall index of 1.00, while the control had a root gall score of 5.0 (>75% of the root

125 system galled). The untreated control plant recorded the highest nematode reproductive
 126 factor (2.4) and final population of 11978.00 J₂/200g of soil. Moringa leaf recorded the least
 127 final root-knot nematode population of 200 J₂/200g of soil. Similarly, Moringa leaf recorded
 128 0.04 reproductive factors.

129 There **were** significant differences (P<0.05) on the mean number of fruit and fruit weight
 130 (kg/ha). Moringa leaf had the highest number of fruits/ plant and weight of fruit, 8.56 and
 131 47.22, respectively. The control recorded the least number of fruits/plant (1.89) and fruit
 132 weight (17.51) (Table 2).

133
 134 Table 3 shows that there **were** no significant differences (P<0.05) **between among the** three
 135 varieties treated with different botanical extracts (leaf and seeds) on the root gall index,
 136 mean number of fruits and fruits weight. However, there was significant difference between
 137 the treated and the untreated pots. With the control having **the** highest root gall index and
 138 lower mean number of fruits and fruit weight for all varieties. On nematode final population
 139 and reproductive factor however, significant differences were observed **between among** the
 140 varieties treated with different botanical extracts. Plants treated with Jatropha leaves and
 141 **seeds** recorded the final population of 0.00 for both the three varieties as compared to their
 142 respective untreated control (10233, 15533.00 and 10167.00 J₂/200 g of soil).

143
 144
 145
 146
 147
 148
 149
 150 **Table 1: Effect of botanical extracts applied on Tomato Varieties on Root-Knot Nematode**
 151 **on Final Population, Reproductive Factor, Root Gall Index and Yield as Affected by Three**
 152 **Tomato Varieties in Pot Experiment**

Varieties	Root-Knot Nematode Parameters				Yield parameters	
	Root Gall Index (RGI)	Initial Population (Pi)	Final Population (Pf)/200g of soil	Reproductive Factor (RF)	Mean Number of Fruits	Mean Fruit Weight/pot(g)
UC28B	1.67	5000.00	1474.00	0.29	6.90	36.63
ROMA VF	1.71	5000.00	2228.00	0.45	6.95	34.87
RIOGRANDE	1.67	5000.00	1461.00	0.29	6.29	39.95
FLSD _{0.05}	ns	Ns	491.90	0.098	Ns	2.984

154 Each value is average of ? replicates. NS= Not Significant, FLSD= Fishers Least
 155 Significant Difference

Table 2: Effect of Botanical Extracts on Root-Knot Nematode Final Population, Reproductive Factor, Root Gall Index and Yield of Infected Tomato (**what is variety) in Pot Experiment**

Botanical Extracts	Root- Knot Nematode Parameters				Yield Parameters	
	Root Gall Index (RGI)	Initial Population (Pi)	Final Population (PF)/200g of soil	Reproductive Factor (RF)	Mean Number of Fruits	Mean Fruit Weight/pot(g)
Castor Leaf	1.11	5000	230.00	0.05	7.33	44.41
Castor Seed	1.33	5000	260.00	0.05	7.56	37.32
Jatropha Leaf	1.11	5000	350.00	0.07	7.33	38.90
Jatropha Seed	1.00	5000	500.00	0.10	7.78	36.33
Moringa Leaf	1.11	5000	200.00	0.04	8.56	47.22
Moringa Seed	1.11	5000	450.00	0.09	6.56	38.36
Control	5.00	5000	11978.00	2.40	1.89	17.51
FLSD _{0.05}	0.28		751.40	0.15	1.53	4.56

Each value is average of ? replicates. FLSD= Fishers Least Significant Difference.

Table 3: Effect of Botanical Extracts on Root-Knot Nematode Final Population, Reproductive Factor, Root Gall Index and Yield of Tomato Varieties in Pot Experiment

	Botanical Extracts	Root-knot Nematode parameters			Yield Parameters	
		Root Gall Index (RGI)	Final Population (Pf)/200g of soil	Reproductive factor (RF)	Mean Number of Fruits	Mean Fruit Weight/pot(g)
UC28B	Castor Leaf	1.33	22.00	0.00	8.67	48.60
	Castor Seed	1.00	28.00	0.01	6.67	38.77
	Jatropha Leaf	1.00	34.00	0.00	8.33	40.40
	Jatropha Seed	1.00	54.00	0.00	8.00	38.03
	Moringa Leaf	1.00	20.00	0.00	8.67	49.73
	Moringa Seed	1.33	12.00	0.00	6.00	40.33
	Control	5.00	10233	2.05	2.00	21.53
ROMA VF	Castor Leaf	1.00	17.00	0.00	6.00	48.66
	Castor Seed	1.67	8.00	0.67	9.67	39.33
	Jatropha Leaf	1.00	7.00	0.00	7.33	33.29
	Jatropha Seed	1.00	5.00	0.00	6.33	35.54
	Moringa Leaf	1.00	27.00	0.01	9.00	50.20
	Moringa Seed	1.00	5.00	0.00	8.00	35.26
	Control	5.00	15533.00	3.11	2.33	15.80
RIO GRANDE	Castor Leaf	1.00	5.00	0.00	7.33	35.97
	Castor Seed	1.33	7.00	0.00	6.33	33.85
	Jatropha Leaf	1.00	65.00	0.00	6.00	35.85
	Jatropha Seed	1.00	55.00	0.00	9.00	43.00
	Moringa Leaf	1.33	38.00	0.01	8.33	43.40
	Moringa Seed	1.00	8.00	0.00	5.67	39.50
	Control	5.00	10167.00	2.03	1.33	15.20
	FLSD _{0.05}	Ns	1301.40	Ns	Ns	Ns

Each value is average of ? replicates. NS= Not significant

DISCUSSION

Scientists are resorting to use botanicals for the control of pest since synthetic pesticides are expensive and hazardous. The use of botanicals as control measures against parasitic nematodes is now the focus of researchers because they are eco-friendly, easy degradable, cost effective and also available. (27)

From this research, leaf and seed extracts from castor, Moringa and Jatropha were able to have effect on final population of root-knot nematodes, root gall index and nematode reproductive factor as well as improved the growth and yield of root-knot nematode infected tomato in the screen house. This is in line with different promising results carried out by different researchers on the management of root knot nematode using different botanicals extracts. On a studies carried out by (12) indicated that some pant parts caused reduction in gall formation by the root knot nematode. (13) also, found that application of sesame seed extract reduced the incidence of root knot nematodes and the severity of galling on okra roots. Previous studies indicated that some pant parts caused reduction in gall formation by the root knot nematode (12).

Report by (14) pointed out that Moringa leaf powder was not phytotoxic to sweet pepper plants but led to increase (?) and even suppress nematode population. The soil populations of root-knot nematode juveniles were found to be significantly reduced within six months under the host plants mostly damaged by *Meloidogyne* spp. (15).

(16) Also stated that plant extracts of basil, marigold, pyrethrum, neem and china berry are were effective in the reduction of nematode population in soil. (17) reported a reduction in number of egg masses, number of females and final larval population of the soil is as a strong indication of the ability of neem leaf extract to control root-knot nematode in tomato.

(18) Found a negative correlation between number of Jatropha species plants and nematode final population as well as root gall index caused by the root-knot nematode. So, from his study, application of 4 plants caused the lowest number of galls (0.8 and 1.3 for *J. curcas* and *J. gossypifolia*, respectively). According to (19), sesame seed extracts have a systemic nematocidal activity against nematodes which may have accounted for the lower number of galls and mean population in treated plants. The study agrees with that of (20) who stated that the application of neem led to decrease in nematode population, reduced galling index leading to increase plant growth of the plots as compared to that of the untreated plants. (21) reported the extracts of neem seeds, leaves and barks had the ability to significantly suppress root galling index and reduce population of *M. javanica* juveniles on sweet gourd.

The reduction in population of root-knot nematodes, nematode reproductive factor and number of root galls formed could be due to the ability of active ingredients present in the botanical extracts (Castor, Jatropha and Moringa) to get in contact with juveniles. This agrees with the findings of (13) that reported reduction on nematode final population and root galling on the root of sweet paper and attributed it to the direct contact of the extracts with the eggs and juveniles of the root-knot nematodes which ensured that the active ingredients

in the Moringa leaf extracts were effectively delivered to the nematode. The reduction in the nematode final population may be due to nematotoxic substances found in the different botanical extracts used. All extracts were effective in increasing yield of the three varieties of tomato as compared to the untreated pots.

The differences in the effectiveness of the different tested plant extracts could be due to the differences in the chemical compositions and concentrations of toxic components contained in the plant material which led to achieved higher yield from the treated plants as compared to the untreated plants. This agrees with the findings of (22) who stated that application of coffee (*Cassia occidentalis*) and lemon grass (*Cymbopogon citrates*) led to provision of the necessary nutrients required by the plant for optimum yield.

Root-knot infestation led to caused stunted growth of all untreated plants and reduced vigor as well as tomato yield, decrease in nematode final population accompanied by increase in yield of tomato plants might be due to the nematicidal potential of the tested plant extract. The effects of the tested plant extracts on yield of tomato plant were different, in some cases.

Galling and reproductive responses are important indicators of host plant reaction than just root-knot galling index. From this study, nematode multiplication rate was high in the untreated plots as compared to those treated by the different botanical extracts. This is in line with the findings of (15) who also reported a lower reproductive factor (Pf) in tomato when treated with *Crotalaria*, *C. juncea* and African marigold (*T. erecta*) plants. The finding of this study agrees with (23) who pointed out that some natural plants can control root-knot nematode in the laboratory or when incorporated into the soil under field condition.

The results of this study is also in line with that of (24) who evaluated some botanicals extract on nematode affecting cowpea. They pointed out that active ingredients contained in the plant materials suppressed the nematode populations in the field. (25) Also reported similar findings of the efficacy of neem compost on nematodes in spinach. Several other plants and organic materials have also been reported to contain different metabolites necessary for plant growth, better yield and at same time toxic to pathogenic microorganisms in the soil (22).

CONCLUSION

The botanical extracts (Castor, Jatropha and Moringa) leaves and seeds used were able to decreased nematode population, reduced root galling as well as reproductive factor and increased yield of the tomato varieties. The botanical extracts can therefore serve as alternative to synthetic nematicides.

REFERENCES

1. Sesso H.D, Lius, Gaziamo J.M. and Buring J.E. (2003) Dietary lycopene tomato- based food products and cardiovascular disease in women. **Journal of Nutrition 133(7):2336-2341.**
2. Tindall H.D. (2005) Commercial vegetable growing. Oxford University press.
3. Mumtaz K.J. (2003). The facts of Tomatoes **your INDIAN COOK, Bawarchi** health and nutrition. [www:bawarchi.com/health/tomato/htmlgk](http://www.bawarchi.com/health/tomato/htmlgk), 2003).
4. Berlinger, M. J. (1986). Pest in the tomato crop: A scientific basis for improvement. Atherton, J.G and Rudich, J. (eds). Chapman and Hall, London. pp 391-432.
5. Ploeg, A.T. (2002). Effect of selected marigold varieties on root-knot nematodes and tomato and melon yields. *Plant Diseases* 86: 505-508.
6. Sasser, J. N. and Carter, C.C. (1985). An **Advanced** Treatise on *Meloidogyne* Vol 1, Biology and control. North Carolina State Uni. Raleigh N. Carolina. Pp. 178-187.
7. Onalo, J.J.I. (1999). Non-chemical control of yam nematode (*Scutellonema brady*). Ph.D Thesis, University of Ibadan, 176pp.
8. Mashela, P.W., Shimelis, H.A. and Mudau, F.N. (2008). Comparison of the efficacy of ground wild cucumber fruits, aldicarb and fenamiphos on suppression of *Meloidogyne incognita* in tomato. *Phytopathology* 156:264-267.
9. Pakeerathan, K., Mikunthan, G. and Tharshani, N. (2009). Eco-Friendly Management of Root-knot Nematode *Meloidogyne incognita* (Kofid and White) Chit- wood Using Different Green Leaf Manures on on Tomato under Field Conditions. *American -Eurasian Journal of Agriculture & Environmental Science* 6 (5): 494-497.
10. Ismail, A.E., (2013). Feasibility of growing *moringa oleifera* as a mix-crop along with tomato for control of *Meloidogyne incognita* and *Rotylenchulus reniformis* in Egypt. *Archives of Phytopathology and Plant Protection* 46(12): 1403-1407.
11. Chitwood, D.J. (2002). Phytochemical based strategies for nematode control. *Annals review of Phytopathology*, 40: 221–249.
12. Zawam, H.S., M.M.A. Youssef and M.H. El-Hamawi, (2003). Effect of lantana (*Lantana camara*) and castor (*Ricinus communis*) as green manure plants on *Meloidogyne javanica* infecting sunflower (*Helianthus annus*) plant. In the Tenth Congress of Phytopathology. *Egyptian Phytopathological Society* of December 2003, Giza. (Egypt), pp: 97-104.
13. Sikora, R. A and E. Fernandez. (2005). Nematode parasites of vegetables. Pp 319-392. In Luc, M., R. A. Sikora and J. Bridge (eds). *Plant-Parasitic Nematodes in Subtropical and Tropical Agriculture*. 2nd Edition. CAB International.
14. Sowleya, E.N.K., Kankama, F. and Adomakob J. (2013) Management of root-knot nematode (*Meloidogyne* spp.) on sweet pepper (*Capsicum annum* L.) with moringa (*Moringa oleifera* Lam.) leaf powder. *Archives of Phytopathology and Plant Protection* **47(13):1531-1538.**
15. Bello, T.T. Fawole B., and Claudius-Cole, A. (2014). Management of Root-Knot Nematodes (*Meloidogyne* Spp.) on Tomato using Antagonistic Plants. *Journal of Biology, Agriculture and Healthcare* 4 (24):**97-100.**
16. Azhar, R.M. and Seddiq, M. (2007). Nematicidal Effects of Some Botanicals against Root-Knot Nematode, (*Meloidogyne javanica*) on Tomato. *International Journal of plant Science* 2: 49-52

17. Fassuliotis, G. (1985). The role of the nematologist in the development of resistant cultivars. Pp. 233-240 in Sasser, J. N. and Carter, C. C. (eds.) **An Advanced Treatise on Meloidogyne**, Vol. I Biology and Control, Raleigh: North Carolina State University Graphics.
18. Ismail, A. and S.A. Hassabo, (2014). Eco-friendly management of reniform nematode, *Rotylenchulus reniformis* using as an intercropping with sunflower in Egypt. *Acta Advances in Agricultural Sciences: In Press*, 2(6). Under Pressing.
18. Ismail, A.E. (2014). Growing *Jatropha curcas* and *Jatropha gossypifolia* as interculture with sunflower for control *M.javanica* in Egypt. *International Journal of sustainable Agricultural Research* 1(2): 39-44.
19. Akhtar, M. and Mahmood, I. (1993). Control of plant parasitic nematodes with “Nimin” and some plant oils by bare- root dip treatment. *Nematologia Mediterranea* 21: 89-92
20. Abid, M: M, A. Zaki and M.A M.A Magbool (1995): Effect of bare root dip treatment in oil-cake and neem leaf extract on root development and growth of tomato and eggplant. *Nematology* 9 (1) 13-16.
21. Yasmin, L., Rashid, M.H., Nazim Uddinm, M., Hossain, M.S. Hossain, M.E. and Ahmed, M.U. (2003). Use of Neem Extract in Controlling Root-knot Nematode (*Meloidogyne javanica*) of Sweet - gourd. *Pakistan Journal of Plant Pathology* 2 (3): 161-168.
22. Izuogu N.B., Yakubu L.B., Abolusoro S.A. and Nwabia I. W. (2015) Efficacy of aqueous leaf extracts of Negro coffee (*Cassia occidentalis*) and lemon grass (*Cymbopogon citratus*) in the Management of Nematode Pests of Okra (*Abelmoschus esculentus* L. Moench). *Science, Technology and Arts Research Journal* 4(3):67-70.
23. Olabiyi, T.I. and E.E.A. Oyedunmade (2005). Effects of carbofuran and aqueous extracts of Basil (*Ocimum gratissimum* L.) in the control of a root- knot nematode, *Meloidogyne incognita* on *Celosia argentea* (L.). *Journal of Agricultural Research and Development* 4(2): 228-235.
- 24 Yakubu, L.B. and Izuogu, N.B. (2013). Comparative Evaluation of some botanicals and carbofuran in the control of root-knot nematode, *Meloidogyne* spp. on cowpea (*Vigna unguiculater*). *Journal of Agriculture and Veterinary Sciences* 5(2): 205-115
25. Olabiyi, T.I., Atungwu, J.J., Izuogu, N.B., Akintola, J. and Abolusoro, S.A (2013). Efficacy of neem compost on root knot nematode pest of Lagos spinach, *Celosia argentea*. *Archives of Phytopathology and Plant Protection* 46(18): 2253-2258.
26. Oluwatayo, J. I., Asiedu, R. and Adesiyan S. O (2011) Allelopathic potential of plant extracts against *Scutellonema bradys*. *African Journal of Root and Tuber Crops* 9(1) 30-37. International society for root and tuber crops – Africa Branch (ISTRC-AB). IITA, LW. Lambourn and co., Dingwall Road, Croydon CR9 3EE, England UK (28-35)
27. Jidere,C.I. and Oluwatayo J.I. (2018) The effectiveness of botanical extracts on root-knot nematode (*Meloidogyne incognita*) egg-Hatch and juvenile mortality in Vitro. *Asian Journal of Agriculture and Horticultural research* 2(3):1-13.