

Stomach content, length-weight relationship  
and **CONDITION FACTORS OF *TILAPIA***  
**SPECIES FOUND IN THE SOMBREIRO RIVER**  
**ACROSS THREE COMMUNITIES IN RIVERS**  
**STATE, NIGERIA.**

**Abstract**

**Aim:** To determine the condition factor incomplete of *Tilapia guineensis* and *Sarotherodon melanotheron* in Sombreiro river across Buguma, Abonnema and Degema communities in Rivers State.

**Study design:** This study employs field work, laboratory experimental design, statistical analysis and interpretation of data.

**Place and duration of study:** live fish samples were caught by fishermen in Buguma, Abonnema and Degema communities in Kalabari kingdom of Rivers State, and were conveyed in a rectangular plastic aquarium containing ice blocks and oxygen bags to the department of Applied and Environmental Biology, Rivers State University. The duration of study lasted for twelve weeks (65days). clearly define the period

**Methodology:** One hundred and eighty fish samples give their average weights and lengths were identified. The fish samples were weighed using an electronic weighing balance, the total length and length of intestine were measured with a meter rule (cm), color of fish, spines and rays of fish were also observed. Their stomach contents were analyzed viewing under a microscope. The frequency of occurrence method and the numerical method were used for analyzing the food items.

**Results :** The stomach content analysis indicated that the major food were phytoplankton. The length and weight relationship in the three sampled stations for *Tilapia guineensis* and *Sarotherodon melanotheron* showed a negative allometric growth, ( $<3$ ). The condition factor for all the fish samples were greater than one. The physico-chemical parameters showed that there were significant differences in the various physico-chemical parameters across the study stations except for turbidity ( $P=0.744$ ) and salinity ( $P=0.922$ ), that showed no significant difference across the study stations.

Comment [AP1]: Give the results in order

**Conclusions:** The length-weight relationship in fishes can be affected by a number of factors including season, habitat, gonad maturity, sex, diet, stomach fullness, health and differences in length ranges of the specimen caught. The exact relationship between length and weight differs among species of fish according to their inherited body shape, and within a species according to the condition (robustness) of individual fish.

Comment [AP2]: Also in order

The stomach content analysis of the sampled fishes provided a baseline study of food and feeding habits of *Tilapia* species in the sampled stations.

**Keywords:** [Tilapia stomach content](#), Condition factor [K](#), [Sombreiro river](#), [monthly communities](#), [Tilapia guineensis](#), [Sarotherodon melanotheron](#), [Sombreiro river](#)

## 1. INTRODUCTION

Tilapia is a common name derived from the Latin word Thiape, Tswana word for fish [reference?](#). This species of fish are mostly dominant in Middle East and Africa among other regions [reference?](#). Tilapia is a fish from the family Cichlidae, one of the famous and common fish found in the marine and fresh water [reference?](#). Tilapia is a major source of protein for man. Therefore, it is necessary that absolute attention is given to the feeding habit of Tilapia. In Nigeria Tilapia is used to compliment the protein source in normal diet (20). According to (22), some Tilapia species feed on wide range of food organism in their aquatic habitat. (7) reported that the principal food items of Tilapia are diatoms, algal filaments and unidentified organic matter(debris).

Stomach content studies can provide useful information to determine the level of Tilapia in the food web and help [us to formulate strategies to manage multispecies in fish farming system](#).

(18) reported that some Tilapia are mostly found in the benthopelagic, potamodromous , fresh water and brackish environment with a depth of five meters and they feed on mostly benthic-algae and phytoplankton. Also, Tilapia is omnivorous with a tendency to be herbivorous.

However, the feeding rate and the body weight, decreases as the fish size increase [reference?](#). Small Tilapia *rendali* consumed significantly more diatom than larger individual (5). (24) reported that the amount of phytoplankton in the Nile Tilapia stomach content is inversely proportional to the body size. However, the stomach content analysis is a search for variation in food consumed by Tilapia. But food and feeding habit of Tilapia are altered by seasonal change as well as the body size development which reflects on its condition factor [reference?](#).

[You have only described the context of the study. What problem do you want to deal with? Where are the objectives of this work? Incomplete introduction in general](#)

Comment [AP3]: use the impersonal form to write

## 2. MATERIALS AND METHODS

The materials used in this study were rectangular plastic aquarium, dissecting sets, dissecting board, microscope (model), meter rule and a weighing balance ( Denver Instrument Germany) TP-512A.

Comment [AP4]: this way is outdated for years. we do not just quote the material

### 2.1. Study area

Describe the zone of the study, by giving the study period, geographical coordinates, climat (quote and describe the duration of seasons), temperature, rainfall, humidity, diagram of the study area...

### 2.21 SAMPLE COLLECTIONS

Fish samples of *Tilapia guineensis* and *Sarotherodon melanotheron* were obtained from fishermen across the three sample stations namely Buguma, Abonema and Degema respectively. The fishermen employed cast and gill nets methods and the live fishes were immediately transferred into a 30 litre rectangular plastic aquarium containing brackish water from the same river, the aquariums were fortified with ice blocks to maintain water temperature and oxygen bags for better aeration and were conveyed to the laboratory of Applied and Environmental Biology of the Rivers State University for analysis. All fishes arrived without mortality to the departmental laboratory where sorting of fish species as well as measurements immediately commenced. The fishing gears used includes cast nets and gill nets. The fish samples were collected in a rectangular plastic aquarium and were conveyed to the laboratory of Applied and Environmental Biology of the Rivers State University for analysis.

### 2.32 Essay conduct METHODOLOGY

One hundred and eighty (180) *Tilapia* fishes weighing between the range of 21.98g-289.45g with total length ranging between 8cm-28cm for *Tilapia guineensis* and 20.54-250.75g with total length of 9cm-27cm of *Sarotherodon melanotheron* were analyzed. **Live fish samples were caught by fishermen across the three sampled stations, namely, Buguma, Abonnema and Degema communities.** All fishes arrived without mortality to the departmental laboratory where sorting of fish species as well as measurements immediately commenced. ~~The fishermen employed cast net method and the live fishes were immediately transferred into a 30 litre rectangular plastic aquarium containing brackish water from the same river, the aquariums were fortified with ice blocks to maintain water temperature and oxygen bags for better aeration. All fishes arrived without mortality to the departmental laboratory where sorting of fish species as well as measurements immediately commenced.~~

The fish samples were identified with the aid of keys according to (11) and (19). Morphological characteristics such as the total length (TL) of the fish ( from the tip of the snout to the tip of the tail fin) was measured with a meter rule. Number of lateral lines, number of rays and spines, mouth part, color of fish and the bands on the fishes were taken. Denver Instrument Germany TP-512A was used to weigh the fishes. Each fish sample was placed in a clear dissecting board and dissected using a dissecting kit. During the dissection; a mild ventral cut was made on each fish from the anal aperture to the base of the operculum, the body wall was cut open at both sides enabling a thorough exposure of alimentary canals, each stomach was slit open. The content was poured into a Petri dish and observation of the food items was carried out with the aid of a dissecting microscope. The

Comment [AP5]: your results show the opposite, we think that you have mixed the contents and observe at the same time. It is therefore necessary to present separately the results of the stomach contents of the 2 tilapias

frequency of occurrence method and the numerical methods were used in analyzing the food items.

In the frequency of occurrence method, the stomach contents were examined and the individual food items were sorted and identified. Thereafter, the number of stomachs containing one or more of each food item were recorded and expressed as a percentage of all stomachs containing the food items.

% occurrence of a food item = Total number of stomach with a particular food /Total number of stomach with food x100

In the numerical method, the number of individual food item in the stomach was sorted and counted. A total of all the food items were recorded and expressed as percentage number of individual food items in the stomach.

% number of a food item = Total number of a particular food item /Total number of all food items x100

The physico-chemical parameters were measured both insitu and exsitu. The insitu measurements were done with instruments such as HANNA instrument model HI 8424, and Extech model DO 700. BOD and COD were done exsitu. Water samples were taken with a clean plastic rubber and were immediately transported to the laboratory for analysis.

The condition factor (K) of the fish was calculated according to (9).

$$K = W \times 100 / L^3$$

Describe: How do you get the length/weight relationships in your results?

you must use the allometric coefficient b of the weight / length relationship obtained in each station to calculate the condition factor K in each station, to be sure of the reliability of the results

## 2.43 STATISTICAL ANALYSIS


Data generated were subjected to one way analysis of variance (ANOVA) with Duncan's Multiple Range Descriptive Test. The results were further analysed using statistic package for social science (SPSS) VERSION 23. Differences among mean were separated with Turkey HSD at P=0.05.

## 3. RESULTS AND DISCUSSION

### 3.1 Stomach content analysis

Results on the stomach content (Figure 1) shows that in the month of August, station 1, *Tilapia guineensis* and *Sarotherodon melanotheron* fed more on phytoplankton, then others and then plant materials. In station 2, *Tilapia guineensis* and *Sarotherodon melanotheron* fed more on phytoplankton, others and then plant materials. In station 3, *Tilapia guineensis* and *Sarotherodon melanotheron* fed more on plant materials, phytoplankton, protozoa and others.

**Comment [AP6]:** Announce the figures or tables in the text before presenting them




**Fig 1: Stomach Content Analysis for August 2018.**

[Present results by species to facilitate understanding. You can present the 2 curves side by side to form a single figure; for each month](#)

In the month of September, station 1, *Tilapia guineensis* and *Sarotherodon melanotheron* fed more on others, phytoplankton and plant materials. In station 2, *Tilapia guineensis* and *Sarotherodon melanotheron* fed more on plant materials, phytoplankton, others and insect parts. In station 3, *Tilapia guineensis* and *Sarotherodon melanotheron* fed more on phytoplankton, others, protozoa and plant materials.

[you do not have a good way to present the results. We can say that you count results](#)



**Fig 2: Stomach Content Analysis for September 2018.**

In the month of October, station 1, *Tilapia guineensis* and *Sarotherodon melanotheron* fed more on phytoplankton, others and plant materials. In station 2, *Tilapia guineensis* and *Sarotherodon melanotheron* fed more on plant materials, phytoplankton and others. In station 3, *Tilapia guineensis* and *Sarotherodon melanotheron* fed more on phytoplankton, others, protozoa and plant materials.




Fig 3: Stomach Content Analysis for October 2018.

### 3.1 PHYSICO-CHEMICAL PARAMETERS

Announce your table before describe the results

e.g. Physico-chemical parameters are presented in table1. It appears that  $p^H$  levels showed a narrow range of variation between different stations, with the highest ( $6.63 \pm 0.2$ ) mean  $p^H$  and the lowest ( $5.6 \pm 0.2$ ). There were however significant differences in  $p^H$  between the stations ( $P=0.00146$ ). Temperature levels differed significantly across the stations ( $P<0.0001$ ) which range from  $25.33- 26.7^\circ\text{C}$ . Conductivity in the three stations vary significantly ( $P=<0.0001$ ) and ranged between  $140 \mu\text{S} / \text{cm} -416.67 \mu\text{S} / \text{cm}$ .

**Table 1: Results of Physiochemical Parameters of the three Sampled Stations.**

Parameters	Temperature (°c)	pH	Conductivity (µS/cm3)	Turbidity (NTU)	Salinity	Chemical Oxygen Demand (mg/L)	Biological Oxygen Demand (mg/L)
Station 1	26.7±0.31 <sup>b</sup>	5.6±0.2 <sup>c</sup>	416.67±10 <sup>a</sup>	7.4±2.13 <sup>a</sup>	8.7±5.12 <sup>a</sup>	16.53±0.1 <sup>a</sup>	3.95±0.1 <sup>a</sup>
Station 2	25.33±0.1 <sup>c</sup>	6.2±0.15 <sup>b</sup>	140±7.64 <sup>c</sup>	8.8±1.82 <sup>a</sup>	8.93±3.46 <sup>a</sup>	42.6±0.01 <sup>c</sup>	1.3±0.01 <sup>b</sup>
Station 3	27.3±0.2 <sup>a</sup>	6.63±0.2 <sup>a</sup>	208.33±15.28 <sup>b</sup>	9.5±5.03 <sup>a</sup>	10.07±4.5 <sup>a</sup>	20.57±0.06 <sup>b</sup>	3.94±0.01 <sup>a</sup>
P	< 0.0001	0.00146	< 0.0001	0.744	0.922	< 0.0001	< 0.0001
Significant Difference	Yes	Yes	Yes	No	No	Yes	Yes

**Key: (abcd) Superscript of different alphabets along columns shows Mean comparisons that are significantly different.**

### 3.2 CONDITION FACTOR **present only your results here. The discussion is not need in this paragraph**

The condition factor (K) for *Tilapia guineensis* found in Abonnema had a mean and standard deviation of  $2.17 \pm 0.71$ , while *Sarotherodon melanotheron* had a mean and standard deviation of  $2.37 \pm 0.36$ . The results showed that the fishes in Abonnema were in a good condition.

The condition factor (K) for *Tilapia guineensis* found in Buguma had a mean and standard deviation of  $2.39 \pm 0.4$ . while *Sarotherodon melanotheron* had a mean and standard deviation of  $2.25 \pm 0.52$ . The results showed that the fishes in Buguma were in a good condition.

The condition factor (K) for *Tilapia guineensis* found in Degema had a mean and standard deviation of  $2.35 \pm 0.39$ , while *Sarotherodon melanotheron* had a mean and standard deviation of  $2.27 \pm 0.33$ . The results showed that the fishes in Degema were in a good condition.

*Tilapia guineensis* sampled in all three stations had a mean and standard deviation of  $2.29 \pm 0.54$  implying that the fishes sampled during the study were in good condition. While *Sarotherodon melanotheron* sampled in all the three stations had a mean and standard deviation of  $2.29 \pm 0.42$  implying that the fishes sampled during this study were in a good condition.

Comment [AP7]: Send to the discussion

Comment [AP8]: This is the discussion, s

There is no significant difference in the condition factor of the fishes across months and stations as shown in the table below.

UNDER PEER REVIEW

**Table 2: Anova (Tukey) Comparisons across Months and Stations**

	Weight	Length	Condition factor K
August	127.49±37.08 <sup>a</sup>	17.11±1.7 <sup>a</sup>	2.58±0.98 <sup>a</sup>
September	80.066±30.34 <sup>b</sup>	14.72±2.37 <sup>b</sup>	2.48±0.73 <sup>a</sup>
October	63.27±27.74 <sup>c</sup>	13.18±2.12 <sup>c</sup>	2.65±0.55 <sup>a</sup>
Pr> F(Model)	< 0.0001	< 0.0001	0.48
Significant Difference	Yes	Yes	No

	Weight	Length	Condition factor K
Station 1	92.61±44.41 <sup>a</sup>	15.33±2.72 <sup>a</sup>	2.41±0.49 <sup>a</sup>
Station 2	90.81±41.89 <sup>a</sup>	14.86±2.59 <sup>a</sup>	2.73±1.01 <sup>a</sup>
Station 3	87.41±39.75 <sup>a</sup>	14.82±2.57 <sup>a</sup>	2.59±0.71 <sup>a</sup>
Pr> F(Model)	0.79	0.492	0.072
Significant Difference	No	No	No

### 3.3 Graphical Expression of Weight- Length Relationship

Figure 4-11 is a graphical expression of length –weight relationship. This also showed the allometric growth of all different species of sampled fishes across the sampled stations. The empirical values were plotted against their respective weight on an arithmetic scale using the Natural logarithm of the length and weight of the fishes.

For the Abonnema as shown in Fig 4 & 5, *Tilapia guineensis* had R<sup>2</sup> Value of 0.912 which shows a strong relationship between the weight and length of the fishes. It also has a b=2.26 (b<3), which shows a negative allometry indicating that the *Tilapia guineensis* in Abonnema grows faster in length than in weight. *Sarotherodon melanotheron* had R<sup>2</sup> Value of 0.952 which shows a strong relationship between the weight and length of the fishes. It also has a b=2.91 (b<3), which shows a negative allometry indicating that the *Sarotherodon melanotheron* in Abonnema grows faster in length than in weight.





Fig 4: Log Length and Log Weight relationship (Bivariate Fit) of *Sarotherodon melanotheron* in Abonema.



**Fig 5: Log Length and Log Weight relationship (Bivariate Fit) of *Tilapia guineensis* in Abonema.**

Formatted: Indent: Left: 0", Hanging: 0.39"

For the Buguma as shown in Fig 6 & 7, *Tilapia guineensis* had  $R^2$  Value of 0.943 which shows a strong relationship between the weight and length of the fishes. It also has a  $b=2.82$  ( $b < 3$ ), which shows a negative allometry indicating that the *Tilapia guineensis* in Buguma grows faster in length than in weight. *Sarotherodon melanotheron* had  $R^2$  Value of 0.774 which shows a strong relationship between the weight and length of the fishes. It also has a  $b=2.58$  ( $b < 3$ ), which shows a negative allometry indicating that the *Sarotherodon melanotheron* in Buguma grows faster in length than in weight





Fig 6: Log Length and Log Weight relationship (Bivariate Fit) of *Tilapia guineensis* in Buguma.

Formatted: Indent: Left: 0", Hanging: 0.49"

UNDER REVIEW



**Fig 7: Log Length and Log Weight relationship (Bivariate Fit) of *Sarotherodon melanotheron* in Buguma.**

Formatted: Indent: Left: 0", Hanging: 0.39"

For the Degema as shown in Fig 8 & 9, *Tilapia guineensis* had  $R^2$  Value of 0.952 which shows a strong relationship between the weight and length of the fishes. It also has a  $b=2.62$  ( $b < 3$ ), which shows a negative allometry indicating that the *Tilapia guineensis* in Degema grows faster in length than in weight. *Sarotherodon melanotheron* had  $R^2$  Value of 0.948 which shows a strong relationship between the weight and length of the fishes. It also has a  $b=2.88$  ( $b < 3$ ), which shows a negative allometry indicating that the *Sarotherodon melanotheron* in Degema grows faster in length than in weight.




Fig 8: Log Length and Log Weight relationship (Bivariate Fit) of *Tilapia guineensis* in Degema




Fig 9: Log Length and Log Weight relationship (Bivariate Fit) of *Sarotherodon melanotheron* in Degema.

Formatted: Indent: Left: 0", Hanging: 0.39"

UNDER

The total *Tilapia guineensis* found in all 3 stations as shown in Fig 10 had  $R^2$  Value of 0.927 which shows a strong relationship between the weight and length of the fishes. It also has a  $b=2.48$  ( $b<3$ ), which shows a negative allometry indicating that the *Tilapia guineensis* in all sampled stations grows faster in length than in weight.




Fig 10: Log Length and Log Weight relationship (Bivariate Fit) of *Tilapia guineensis* sampled in all stations.

Formatted: Indent: Left: 0", Hanging: 0.49"

The total *Sarotherodon melanotheron* found in all 3 stations as shown in Fig 11 had  $R^2$  Value of 0.891 which shows a strong relationship between the weight and length of the fishes. It also has a  $b=2.75$  ( $b<3$ ), which shows a negative allometry indicating that the *Sarotherodon melanotheron* in all sampled stations grows faster in length than in weight.




Fig 11: Log Length and Log Weight relationship (Bivariate Fit) of *Sarotherodon melanotheron* sampled in all stations.

Formatted: Indent: Left: 0", Hanging: 0.49"

#### 4. DISCUSSION [the discussion also follows the chronological order of the specific objectives](#)

Two Species of Tilapia were identified in this study. They Include *Tilapia guineensis* and *Sarotherodon melanotheron*.

##### 4.1 Stomach Content Analysis

The stomach contents of Tilapia species found in this study across stations were mainly plants compositions ranging from phytoplankton, plants parts, etc. indicate that they were herbivorous. They feed mainly on plant food substances such as phytoplankton, plant parts, leaf parts and some percentages of animal food include insect pupae, insect larva and protozoa. This agrees with (10), (17) and (7) who reported that *Oreochromis niloticus* are obligate herbivores that feed on algal filaments, diatoms and unidentified organic matter. Similarly, (2) reported that the main stomach content of *Synodontis niloticus* is phytoplankton.

(12) and (16) reported that insects and crustaceans comprise a large portion of the diet of *Oreochromis niloticus* and also reported that *O. niloticus* have the ability of feeding on either small or bulky particles and can efficiently filter, and utilize a broad range of particle size (13). (21) reported that Blackchin Tilapia feeds primarily on phytoplankton and filamentous algae. The stomach contents of these fish also include granules of mud and sand implying that they suction feed on the bottom of their aquatic habitat.

## 4.2 Physico-chemical Parameters.

pH levels showed a narrow range of variation between different stations, with the highest ( $6.63 \pm 0.2$ ) mean  $p^H$  and the lowest ( $5.6 \pm 0.2$ ). There were, however significant difference in pH between the stations ( $P = 0.00146$ ).

Temperature levels differed significantly across the stations ( $p = < 0.0001$ ), with range from 25.33 to 26.7°C.

Conductivity in three stations vary significantly ( $P$  value =  $< 0.0001$ ) and ranged between 140  $\mu\text{S/cm}$  to 416.67  $\mu\text{S/cm}$ . (6) noted that conductivity values greater than 100  $\mu\text{S/cm}$  were indicative of human activity. According to (23), water conductivity of between 150 and 500  $\mu\text{S/cm}$  is ideal for fish culture (23). (26), however, put the desirable range of conductivity for fishponds at between 100 and 2000  $\mu\text{S/cm}$ .

Comment [AP9]: Results presented or discussion?

## 4.3 Length /and weight Relationship

The length-weight relationship in fishes can be affected by a number of factors including season, habitat, gonad maturity, sex, diet, stomach fullness, health and preservation techniques, and differences in the length ranges of the specimen caught. The exact relationship between length and weight differs among species of fish according to their inherited body shape, and within a species according to the condition (robustness) of individual fish. The logarithmic values of length were plotted against their logarithmic respective weight on an arithmetic scale, smooth curve was obtained. Statistical analysis of LWR showed that the regression coefficients obtained from length – weight relationships (LWR) as presented here was a significant correlation between length and weight.

The results from the study showed a strong relationship between the length and weight of the two Tilapia species sampled across stations as shown in Fig 4 to Fig 11. The average  $b$  values were less than 3 for both species which shows that most of the fishes had a negative allometric growth

i.e. most of the Tilapia grows faster in length than in weight. It also indicates when  $b < 3$ , either that the large specimens have changed body shape, i.e., become more elongated, or the small specimens were in better nutritional condition at the time of sampling. These results agree with the studies of (8) and so many other fisheries studies that postulated that there is a strong relationship between the length and the weight of Tilapia species and that generally exhibit a negative allometric growth pattern.

#### 4.4 Condition Factor

The individual fish species condition is determined based on the analysis of length weight data reflecting that the heavier fish at a given length is in better condition (4), hence indicating favourable condition. The Tilapia species found across stations had an average condition factor above two (2). This shows that the fishes are in a favourable condition and agrees with the studies of Puntiuschola , (1), (3) and (15).

Condition sometimes reflects food availability and growth within the weeks prior to sampling. However, condition is variable and dynamic. Individual fish within the same sample vary considerably, and the average condition of each population varies seasonally and yearly.

The differences were attributed to the effect of eutrophication and pollution on growth and other biological aspects of *Oreochromis niloticus* as suggested by, (14) and, (25) , also observed similar results. The study presented the basic information on length – weight relationships and condition factor of *T. mossambica* of study area which would be useful for fishery managers as well as the sustainable management.

Formatted: Font: Italic

Formatted: Font: Italic

5. **Conclusion** the conclusion is the answer to the specific objectives, ie the summary of the result of each specific objective in chronological order

The stomach contents analysis indicated diverse food sources. The major food source was phytoplankton. Other food samples from plant sources found in the stomach analysis were; leaf plants, plant tissue, etc. The animal sources were insect parts, insect larva, protozoa and others. The stomach content analysis of the sampled fishes provided a baseline study of food and feeding habits of Tilapia species in the sampled stations

#### Physico-chemical Parameters ??

Formatted: Font: Not Bold

~~The length weight relationship in fishes can be affected by a number of factors including season, habitat, gonad maturity, sex, diet, stomach fullness, health and preservation techniques, and differences in the length ranges of the specimen caught. The exact relationship between length and weight differs among species of fish according to their inherited body shape, and within a species according to the condition (robustness) of individual fish~~

~~The stomach contents analysis indicated diverse food sources. The major food source was phytoplankton. Other food samples from plant sources found in the stomach analysis were; leaf plants, plant tissue, etc. The animal sources were insect parts, insect larva, protozoa and others. The stomach content analysis of the sampled fishes provided a baseline study of food and feeding habits of Tilapia species in the sampled stations.~~

The condition factor for all the fish samples were greater than one. This result is an indication that all the fishes were in a good condition. The condition factor for the sampled fishes in the three sampled stations agreed with the length and weight relationship

The length and weight relationship in the three sampled stations for the *Tilapia guineensis* and *Sarotherodon melanotheron* showed a negative allometric growth, i.e. they were all  $< 3$ . This is an indication that fishes grow longer in length than in weight. The length-weight relationship in fishes can be affected by a number of factors including season, habitat, gonad maturity, sex, diet, stomach fullness, health and preservation techniques, and differences in the length ranges of the specimen caught. The exact relationship between length and weight differs among species of fish according to

their inherited body shape, and within a species according to the condition (robustness) of individual fish

Comment [AP10]: To be send in the discussion section

~~The condition factor for all the fish samples were greater than one. This result is an indication that all the fishes were in a good condition. The condition factor for the sampled fishes in the three sampled stations agreed with the length and weight relationship~~

## REFERENCES

1. Afroze, S., Hossain, M.A. & Parween, S. (1992). Notes on the size frequency distribution and length-weight relationship of fresh water fish. *Amblyphanygodan mold* (Hamotton) (cyprinfurier cyprinid). *University Journal of Zoology, Rajshatic University*, 10(11), 103 – 104
2. Akintunde, E. A. & Imevbore, A. M. (1979). Aspect of Biology of Cichlid Fishes of Lake Kainji with special references in *Sarotherodengalilacus*. *Nigerian Journal of Natural Science*, 1, 35 – 39.
3. Alam, M. R., Mortuza, M. G., Islam, M. S. & Hossain, M. A. (1994). Notes on the size frequency distribution and length-weight relationship of the freshwater fish *AiliaCoilia* (Hamilton-Buchanan) (Siluriformes:Schibeidea). *University of Journal Zoology, Rajshai University*, 13, 69 – 70.
4. Bolger, T. P. L. & Connolly, P. L. (1989). The suitable indices for the measurement analysis of fish condition. *Journal of Fish Biology*, 34, 171 – 182..

5. Brummett, R. E. (1995). Weed control by adult and juvenile tilapia rendalli in rainfed ponds. NAGA, the ICLARM quarterly.
6. Crane, B. (2006). Results of Water Quality Measurements in Messer Pond. Available at <http://www.messerpond.org/Ecology/WaterSamplingSummary.pdf>. Accessed 23 Aug 2018.
7. Fagade, S. O. & Olaniyan, C. I. O. (1973). Food and feeding inter-relationship of fishes in Lagos Lagoon. *Journal of Fish Biology*, 5, 203 – 225.
8. Froese, R. (2006). Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22, 241 – 253.
9. Fulton, T.W. (1904). The rate of growth fishes. Twenty-second Annual Report, Part 111. Fisheries Board of Scotland, Edinburgh, 141-241.
10. Gwahaba, J. J. (1973). Stomach content analysis. A review of method and application. *Journal of Fish Biology*, 17, 14 – 429.
11. Idodo-Umeh, G (2003). Fresh Water Fishes of Nigeria (Taxomy, Ecological notes, diet and utilization.) Idodo Umeh, Benin City, Nigeria
12. Job, B. E. & Udo, P.J (2002). Food, feeding and condition factor of the Estuaries catfish *chrysiichthysnigrodigitatus* (Lacepe) of the cross River Estuary, Nigeria. *African Journal of Fisheries and Aquaculture*, 3, 43 – 51.
13. Job, B. E. & Nyong, E. A. (2004). Diet composition and condition of the African Rivers Prawn *Macrobrachium vollehoovenii* (Herklot, 1857) Crustacea Decapoda, Palaemonae) of the cross Rivers Estuary, Nigeria. *African Journal of Fisheries and Aquaculture*, 4, 9 – 17.
14. Khallaf, E. A., Galal, M. & Authman, M. (2003). The biology of *Oreochromis niloticus* in a polluted canal. *Ecotoxicology*, 12, 405 – 416.
15. Mortza, M. G. & Mokarrama, N. T. (2000). Notes on the length-weight relationship and condition factor of mud loach *Botialohachata* (Chaudhuri) (Cypriniformes: Cobitidae). *University of Journal of Zoology, Rajshahi University*, 19, 113 – 114.
16. Northcott, M.E., Beveridge, M.C.M. & Ross, L.G. (1991). A laboratory investigation of the filtration and ingestion rates of the Tilapia (*Oreochromis niloticus*) feeding on two species of blue green algae. *Bio Fishes*, 31, 73 – 85.
17. Noble, R.I. (1989). Biological control of Aquatic Weeds Using Fish Water Resources. *Research Institute Proceedings of Workshop on Management of aquatic weeds and mosquito in impoundments*. University of North Carolina at Charlotte, 14-15.
18. Nwafili, S. A. & Lamai, S. L. (2003). An Assessment of Fish species diversity of Shiroro Tropical Man-made Lake after a decade of Post-impoundment. *Journal of Applied Science, Nigeria*, 1, 62 – 63.
19. Nwani, C.D. (2004). Aspects of the biology of mormyrids in Anambra River, Nigeria. ph.D Thesis University of Nigeria Nsukka. 194.
20. Olayide, P. (1992). Food and condition of fresh water fish in Nigerian, Nature conservation preservation and management 1<sup>st</sup> edition published in Great Britain by Chapman and Hall 2 – 6 Boundary Row London, 126 – 127.
21. Pauly, D. (1976). The biology, fishery and potential for aquaculture of *Tilapia melanotheron* in a small West African lagoon. *Aquaculture*, 7, 33 – 49.

22. Pius, M. O. & Benedicta, O. O. (2002). Food and feeding inter-relationship. A preliminary to the formulation of the feed of some Tilapiine fishes. *Tropical Journal of Animal Science* 5(1), 36 – 41.
23. Russell, M., Shuke, R. & Samantha, S. (2011). Effects of Conductivity on Survivorship and Weight of Goldfish (*Carassius auratus*). Available at [http://departments.juniata.edu/biology/eco/documents/Russell\\_et\\_al.pdf](http://departments.juniata.edu/biology/eco/documents/Russell_et_al.pdf). 23 April 2018
24. Saha, S. N. & Dewan, S. (1979). Food and feeding habits of *Oreochromis niloticus* L. Types and amount of food taken by the fish and its size and patterns of feeding. *Bangladesh Journal Zoology*, 7, 53 - 60.
25. Santic, M., Pallaoro, A. & Jardas, I. (2006). Co-variation of gonadosomatic index and parameters of length-weight relationships of Mediterranean horse mackerel, *Trachurus mediterraneus* (Steindachner, 1868). In the eastern Adriatic Sea. *Journal Applied of Ichthyology*, 22, 214 – 217.
26. Stone, N., Shelton, J.L., Haggard, B.E. & Thomforde, H.K. (2013). Interpretation of Water Analysis Reports for Fish Culture. *Southern Regional Aquaculture Center (SRAC) Publication*, 4606(12).

UNDER PEER REVIEW