

Original Research Article

**EFFECTS OF ORGANIC AND INORGANIC FERTILIZER ON THE EARLY
GROWTH RESPONSE OF *Azelia Africana***

ABSTRACT

An experiment was conducted in the nursery of Department of Forestry Technology at the Federal College of Forestry Ibadan, Oyo state, Nigeria to determine the effect of organic and inorganic fertilizers on early growth response of *Azelia africana*. Seedlings were collected from Forestry Research Institute of Nigeria, Ibadan, Oyo State and were transplanted into 2kg perforated polythene pots. The experiment was laid out in a completely randomized design with five treatments replicated three times. The treatments were: 20 t ha⁻¹ water hyacinth compost (T1), 20 t ha⁻¹ water hyacinth + poultry manure compost (T2), 20 t ha⁻¹ poultry manure (T3) and 50 kg N:P:K 20:10:10 (T4) and T5 - control (no fertilizer application). The compost was applied two (2) weeks before planting, while NPK 20:10:10 was applied two 2 weeks after planting. The experiment was monitored for eight (8) weeks after transplanting (WAT), while growth parameters were measured. The results of the study showed that application of fertilizers gave significant (p=0.05) increase in plant height (cm), stem diameter (mm), leaf production, and leaf area (cm²) of *A. africana*. Plant height ranged from 41.43 cm in the control to 47.96 cm in the pots where 20 t ha⁻¹ water hyacinth compost was applied. Stem diameter also increased appreciably across treatments, while leaf production ranged from 9 in the control treatment to 14 in the pots with 20 t ha⁻¹ poultry manure. These results suggested that the incorporation of organic and inorganic fertilizers increased productivity of *A. africana* and hence application of 20 t ha⁻¹ water hyacinth compost may be recommended for the production of *A. africana* especially in the study area.

Comment [H1]: Standardize the way you write NPK.

Formatted: Superscript

Comment [H2]: Standardize the way you write NPK.

Comment [H3]: Avoid these words. These are not technical or scientific terms.

27 Keywords: Water hyacinth compost; *Afzelia africana*; NPK 20:10:10

28

29

INTRODUCTION

30 Forests and trees perform various functions in the ecosystem namely; aesthetics, provision of
31 food and medicine, provision of shelter to wildlife and hygienic purpose (Agbogidi and
32 Eshegbeyi, 2008). It is universally accepted that forests and trees carry out a fundamental role
33 in soil and water resources conservation (Broadhead and Leslie, 2007; Hamilton, 2008). As
34 population density increases and land for food production expands due to agricultural
35 activities and urbanization, natural forests became degraded (Salim and Ullsten, 1999). The
36 degradation has led to the disappearance of most species including agro forest trees and
37 causing difficulty in growing some seedlings including *Afzelia africana* (Nwoboshi, 1985;
38 Keay, 1989; Etukudo, 2000; Ezenwaka *et al.*, 2004). *Afzelia africana* is a leguminous tree
39 found in the humid and dry forest savannah borders or semi-deciduous forest (Keay, 1985). It
40 is used for soil conservation and improvement (Agbogidi and Onomeregbor, 2007). *Afzelia*
41 *africana* is a timber species with high forage, economic and pharmacological values. Its
42 leaves are harvested for grazing during the dry season. The high demand for *A. africana*
43 leaves, seeds, roots and barks for various uses has resulted in corresponding increase in the
44 exploitation at such a rate that sustainability of this natural resource cannot be guaranteed
45 (Mtambalika *et al.*, 2014; Palgreave, 2002). Documented reports on the cultivation and
46 seedling growth of this multipurpose tree known commonly as African mahogany are scarce
47 (Okeke, 1996; Burkill, 1999; Etukudo, 2000; Agbogidi *et al.*, 2008). If the benefits derivable
48 from *A. africana* must continue especially for the future generations, there is the need to
49 stimulate farmers' interest in the cultivation of *A. africana* thereby helping to reduce poverty,
50 helping in conservation role as well as to boost the source of revenue for the government. In

51 | the same vein, information on the domestication of the plant seeds and seedlings are in **piece**
52 | **meal** due mainly to the poor nature of the soil caused by human influences. Consequently, the
53 | need to acquaint farmers with the most successful soil or manures that could enhance the
54 | growth of *A. africana* seedlings cannot be overemphasized. This study was aimed at
55 | investigating the effect of organic and inorganic fertilizers on the seedling establishment of
56 | ~~*Af.zelia*~~ *africana* with a view to recommend the best fertilizer to *A. africana* growers
57 | especially at the nursery stage and to multiply this multi-purpose species ~~that nature has~~
58 | ~~bequeathed to mankind.~~

Formatted: Highlight

59

60 MATERIALS AND METHODS

61 | The experiment was carried out in 2018 at nursery site of the Department of Forestry
62 | Technology, Federal College of Forestry Jericho Ibadan (Latitude 07⁰ 27¹N and longitude 03⁰
63 | 53¹E), Ibadan, Nigeria (FRIN Meteorological Station, 2018). The annual rainfall is 1250 mm
64 | with a bimodal pattern and has a minimum temperature of 21.9 °C and maximum temperature
65 | of 35.5 °C. The experiment was laid out in a Complete Randomized Design with five
66 | treatments replicated three times. The treatments were: 20 t/ha water hyacinth compost (T1);
67 | 20 t ~~4~~¹ha⁻¹ water hyacinth + poultry manure compost (T2); 20 t/ha poultry manure (T3); ~~and~~
68 | 50 kg/ha N:P:K 20:10:10 (T4); and ~~T5~~ control (no fertilizer application) (T5). The compost
69 | was applied two (2) weeks before planting. Eight week old seedlings of *A. africana* were
70 | collected from Forestry Research Institute of Nigeria (FRIN), Ibadan, Oyo State, and potted
71 | into 2kg polythene pots filled with different media treatments, watered and allowed to
72 | stabilize for two weeks before the commencement of growth assessment. Plant height was
73 | measured with a meter rule at the distance from soil level to terminal bud. Leaf production
74 | was determined by counting. Stem diameter ~~at the collar~~ was measured with venier caliper.
75 | Data collected were analysed statistically using Genstat Software Package and were subjected

Formatted: Superscript

76 to analysis of variance. Means were separated using Duncan's multiple range test (DMRT) at
77 5% level of significance.

78

79 RESULTS

80 Table 1: Pre-planting soil physical and chemical properties of the experimental site

| Soil parameters | Content in soil |
|--|-----------------|
| pH (H ₂ O) | 5.8 |
| Organic carbon (g kg ⁻¹) | 9.0 |
| Total nitrogen (g kg ⁻¹) | 1.0 |
| Available phosphorus(mg kg ⁻¹) | 6.0 |
| Exchangeable cations (cmol _e kg ⁻¹) | |
| Ca | 1.32 |
| Mg | 0.25 |
| K | 0.11 |
| Na | 0.31 |
| Extractable micronutrient (mg kg ⁻¹) | |
| Mn | 302.0 |
| Fe | 265.0 |
| Cu | 3.39 |
| Zn | 1.2 |
| Exchangeable Acidity (cmol _e kg ⁻¹) | 0.40 |
| Particle size distribution (g kg ⁻¹) | |
| Sand | 838 |
| Silt | 54 |
| Clay | 108 |
| Textural class | Sandy loam |
| Bulk density (g cm ⁻¹) | 1.62 |
| Saturated hydraulic conductivity (cm hr ⁻¹) | 12.4 |

Formatted: Subscript

Formatted: Subscript

81

82
83
84
85
86
87
88
89
90
91
92
93
94
95
96

Table 2: Chemical properties of poultry manure, water hyacinth compost and water hyacinth + poultry manure compost used.

Comment [H4]:

| Parameter | Poultry manure | Water hyacinth compost | Water hyacinth + Poultry manure compost |
|-------------------------|----------------|------------------------|---|
| pH (H ₂ O) | 6.8 | 5.89 | 5.76 |
| Organic carbon (%) | 22.94 | 31.92 | 32.11 |
| Total Nitrogen (%) | 3.30 | 2.6 | 1.73 |
| C:N | 6.8 | 12.52 | 43.99 |
| Phosphorus <u>P</u> (%) | 0.83 | 1.24 | 1.55 |
| Potassium <u>K</u> (%) | 1.80 | 0.47 | 0.80 |
| Calcium (%) | 2.56 | 1.6 | 1.80 |
| Mgagnesium (%) | 1.58 | 4.30 | 0.92 |

The physical and chemical properties of the soil (0 – 15 cm depth) at the experimental site before planting is as presented in Table 1. The soil is sandy loam, slightly acidic, high bulk density (1.62 g cm⁻³) and has been classified as an Alfisol (Smyth and Montgomery, 1962) with its distinctive characteristics. The data in Table 1 further confirms this assertion and also reveals that the soils are moderate in zincZn, low in potassiumK (0.11 cmol_c kg⁻¹), organic carbon (9.0 g kg⁻¹), total nitrogen (1.0 g kg⁻¹) and phosphorousP (6.0 mg kg⁻¹). Saturated hydraulic conductivity value of 12.4 cm hr⁻¹ indicated a well drainedwell-drained soil.

Formatted: Superscript

Formatted: Subscript


Formatted: Superscript

Formatted: Superscript

Formatted: Superscript


Formatted: Superscript

97 | —The chemical composition of water hyacinth compost, poultry manure and water
 98 | hyacinth + poultry manure compost used is as presented in Table 2. The pH of all the organic
 99 | fertilizers were all acidic with the water hyacinth + poultry manure compost having the most
 100 | acidic pH of 5.76. Organic carbon, C/N and [phosphorus-P](#) were highest in the mixture of
 101 | water hyacinth + poultry manure compost. Poultry manure was highest in [nitrogen-N](#),
 102 | [Kpotassium](#) and [Caalcium](#) content



103 |
 104 | Figure 1: Growth pattern of number of leaves of *Afzelia africana* seedlings as influenced by
 105 | fertilizers
 106 | [T1 \(20 t ha⁻¹ water hyacinth\); T2 \(20 t ha⁻¹ water hyacinth + poultry manure](#)
 107 | [compost\); T3 \(20 t ha⁻¹ poultry manure\); T4 \(50 kg ha⁻¹ NPK 20:10:10\); T5 \(control – no](#)
 108 | [fertilizer application\).](#)

109 |



110 Figure 2: Effects of fertilizers on number of leaves of *Afzelia africana* seedlings

112 T1 (20 t ha⁻¹ water hyacinth); T2 (20 t ha⁻¹ water hyacinth + poultry manure
 113 compost); T3 (20 t ha⁻¹ poultry manure); T4 (50 kg ha⁻¹ NPK 20:10:10); T5 (control – no
 114 fertilizer application).


116
 117
 118
 119 **Effect of organic and inorganic fertilizers on number of leaves of *Afzelia africana***
 120 **seedling**

121 Figure 1 showed the growth pattern of leaf production of *Afzelia africana* seedlings as
 122 influenced by the application of organic and inorganic fertilizers. Results revealed that mean
 123 leaf production of *Afzelia Africana* increased across the study period. T3 (20 t ha⁻¹ +
 124 poultry manure) recorded the highest mean number of leaves across the weeks while T5 (top
 125 soil only) had the least mean number of leaves all through the weeks.

126 The effect of organic and inorganic fertilizers on the number of leaves of *Afzelia africana* is
 127 as presented in Figure 2. Leaf production increased appreciably across treatments and differs
 128 significantly (p=0.05) throughout the experiment. Leaf production ranged from 9 in the


Formatted: Left, Space After: 0 pt, Line spacing: single

129 control medium to 14 in the pots where 50 kg ha^{-1} NPK 20:10:10 was applied. The highest
 130 mean leaf production of *Azelia africana* was observed in T3 with 14 and it was closely
 131 followed by T1, T2, and T4 with 13 and the least was found in T5 (control) with
 132 9.



133
 134 **Figure 3: Growth pattern of height of *Azelia africana* seedlings as influenced by fertilizers**
 135 **Legend:** T1 (20 t ha^{-1} water hyacinth); T2 (20 t ha^{-1} water hyacinth + poultry
 136 manure compost); T3 (20 t ha^{-1} poultry manure); T4 (50 kg ha^{-1} NPK 20:10:10); T5 (control – no
 137 fertilizer application).

144 **Figure 3: Growth pattern of height of *Azelia africana* seedlings as influenced by fertilizers**
 145



146

147 Figure 4: Effects of fertilizers on plant height of *Azelia africana* seedlings

148 [T1 \(20 t ha⁻¹ water hyacinth\);](#) [T2 \(20 t ha⁻¹ water hyacinth + poultry manure](#)
 149 [compost\);](#) [T3 \(20 t ha⁻¹ poultry manure\);](#) [T4 \(50 kg ha⁻¹ NPK 20:10:10\);](#) [T5 \(control – no](#)
 150 [fertilizer application\).](#)

151

152

153

154

155 **Effect of organic and inorganic fertilizers on plant height of *Azelia africana* seedling**

156 Results in Figure 3 showed the growth pattern of plant height of *Azelia africana* seedlings.

157 ~~T~~, the control ~~experiment-treatment~~ followed similar trend with the growth pattern of number

158 of leaves and performed poorly in comparison to others in terms of the number of leaves

159 produced. The highest mean height of *Azelia africana* was observed in T1 (20t/ha water

160 hyacinth compost) with T2 (20 ~~t ha⁻¹ t/ha~~ water hyacinth + poultry manure compost) closely

161 followed and T5 (control) performed least. The effect of organic and inorganic fertilizers on

162 the plant height of *Azelia africana* is as presented in Figure 4. Plant height increased


163 **appreciably** across treatments but did not differ significantly (p=0.05) throughout the

164 experiment. 20 ~~t ha⁻¹t/ha~~ water hyacinth compost (47.96 cm) had highest plant height

Formatted: Highlight

165 | followed by 20 $t\ ha^{-1}$ water hyacinth + poultry manure compost (47.89 cm) and least by
166 | control pot (41.43 cm).


167 |
168 |



169 |

170 | Figure 5: Growth pattern of stem diameter of *Azelia africana* seedlings as influenced by
171 | fertilizers
172 | T1 (20 $t\ ha^{-1}$ water hyacinth); T2 (20 $t\ ha^{-1}$ water hyacinth + poultry manure
173 | compost); T3 (20 $t\ ha^{-1}$ poultry manure); T4 (50 $kg\ ha^{-1}$ NPK 20:10:10); T5 (control – no
174 | fertilizer application).

175 |



176

177 Figure 6: Effects of fertilizers on stem diameter of *Azelia africana* seedlings

178 T1 (20 t ha⁻¹ water hyacinth); T2 (20 t ha⁻¹ water hyacinth + poultry manure
 179 compost); T3 (20 t ha⁻¹ poultry manure); T4 (50 kg ha⁻¹ NPK 20:10:10); T5 (control – no
 180 fertilizer application).

181

182 **Effect of organic and inorganic fertilizers on stem diameter of *Azelia africana* seedling**

183 Results in figure 5 showed the growth pattern of stem diameter of *A. azelia africana* seedlings,
 184 the control experiment performed well at the beginning but poorly at the end of the eighth
 185 week, with T1 (20 t ha⁻¹ water hyacinth) performing best.

186 The effect of organic and inorganic fertilizers on the stem diameter of *A. azelia africana* is
 187 presented in Figure 6. Stem diameter increased appreciably across treatments but did not
 188 differ significantly (p=0.05) throughout the experiment. 20 t ha⁻¹ water hyacinth compost
 189 (0.81 mm) had highest plant height followed by 20 t ha⁻¹ water hyacinth + poultry
 190 manure compost (0.76 mm) and least by control plot (0.62 mm).

Formatted: Highlight

Comment [H5]: It was not the treatment that presented the highest height or diameter ... Was: The plants were submitted to this treatment. This should be neat throughout the text.

191

192

193

194 **DISCUSSION**

195 The major factor affecting plant growth in the tropics is the nutrient deficiency in tropical
196 soils resulting from degraded farmland. The low levels of **nitrogenN**, **phosphorusP**, and
197 **organic carbon** observed in the experimental soil indicated that the soil had a low fertility
198 status. The value obtained for N, P, K and **organic C** is below the critical range (Adeoye and
199 Agboola, 1985, Akinrinde, *et al.*, 2005), thus indicating poor soil fertility not suitable without
200 the addition of external input for planting *A. f. africana*. The chemical composition of
201 water hyacinth compost, poultry manure and water hyacinth + poultry manure compost used
202 in the experiment was relatively high in major elements (N, P, K, Ca and Mg). The **organic**
203 **carbon** content of all the organic fertilizers were less than the values obtained for the
204 composts of *Azadirachta indica*, *Albizia lebbek* and *Khaya senegalensis* by Daldoum and
205 Hammad (2015).

206 The application of the various fertilizers increased the growth of *A. Africana* this agreed with
207 the findings of Uddin (2014) where organic fertilizers enhanced the seedling growth of some
208 leguminous **agroforestry** species. This could result from the nutritional benefits of organic
209 and inorganic fertilizers which include improvement of soil fertility. The result obtained from
210 the plant height showed that water hyacinth compost significantly induced the shoot growth,
211 leaf production and stem diameter of *A. africana* seedlings. This is in support with Razaq, *et*
212 *al.* (2017); Talkah (2015) and Cuesta (2010) that reported that plant height and number of
213 leaves of plants treated with water hyacinth compost had been used and showed showed
214 better performance than control. ~~This result is also in line with the results b~~By Lata (2013)
215 that experimented with water hyacinth manure on *Coriandrum sativum* and revealed positive
216 response with increase in manure rates. ~~This was also s~~upported by study done by Osoro, *et*
217 *al.*, (2014) and Aboul-Enein *et al.* (2011) who advocated that water hyacinth has good N, P,
218 K absorbing capacity from water and thus can be used as a good source of compost material

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Comment [H6]: Organic C or organic carbon?
Arrange throughout the text.

219 to serve as fertilizer in soil with poor amount of N, P, K and C values. Water hyacinth which
220 used to be tagged as waste and nuisance to aquatic environment can be converted to compost
221 for fertilizing plants at the nursery stage in order to improve early growth. This might replace
222 use of expensive, scarce and environmentally hazardous inorganic fertilizers in forest nursery
223 work.

224

225 CONCLUSION

226 Organic and inorganic fertilizer had effect on the growth of *A. f. elia Africana*. It could be
227 observed from the result obtained that there were increases in plant height, stem diameter and
228 number of leaves. However, there was poor performance throughout the assessment period in
229 control treatment when compared to other treatments in terms of the number of leaves
230 produced per plant. Seedlings ~~with in~~ 50 kg ~~ha~~⁻¹ N:P:K 20:10:10 (T4) ~~presented~~ ~~performed~~
231 best at week eight though it was poor at the early stage of the experiment. Seedlings with 20 t
232 ~~ha~~⁻¹~~ha~~ poultry manure (T3) performed well from the beginning of the experiment up to the
233 penultimate week to the end of the experiment. Application of 20 t ~~ha~~⁻¹ ~~t/ha~~ water hyacinth
234 compost gave the highest plant height. Stem diameter also increased appreciably across
235 treatments. Therefore, it can be concluded and recommended that 20 t ~~ha~~⁻¹ ~~t/ha~~ water
236 hyacinth compost can be used by farmers to increase the growth of *A. f. elia africana*.

237

238 REFERENCES

239 Aboul-Enein A, Al-Abd A, Shalaby E, Abul- Ela F, Nasr-Allah A, Mahmoud A. (2011).
240 *Eichhornia crassipes* (Mart) solms, Plant Signaling & Behavior. 2011;6(6):834-836.

241

242 Adeoye, G. O., and Agboola, A. A., (1985). Critical level for soil pH, available P, k, Zn and
243 maize ear leaf content for P, Cu and Mn in sedimentary soils of Southwestern Nigeria. 6: 65 –
244 71.

245

Formatted: Superscript

246 Agbogidi O. M, Onomeregbor V. A (2007). Enhancing germination in seeds of *Afzelia*
 247 *africana* (Sm. Expers.). In: Olufajo, OO, Omokore DF, Akpa GN Sanni SA (ed.).
 248 *Proceedings of the 41st Annual Conference of the Agricultural Society of Nigeria* held in
 249 Zaria, Kaduna State, between 22nd and 26th October, 2007. Pp 260-263.

250

251 Agbogidi OM, Onomeregbor VA, Avwevughwar OI (2008). Eco Physiological effects of
 252 crude oil pollution *Afzelia africana* (Sm. Ex pers.) In: Ezike, K.N.N., Osakwe, I.I., Ekwu,
 253 L.G., Utobo, E. and Mba, C.N. (eds.). *Proceedings of the 42nd Annual Conference of the*
 254 *Agricultural Society of Nigeria* held at the Ebonyi State University, Abakakili, Ebonyi state,
 255 between 19th and 23rd October, 2008. Pp 365-369.

256

257 Agbogidi, O. M. and Eshegbeyi, O. F. (2008). Forestry development for a safe environment.
 258 In: Onyekwelu J.C. Adekunle, V.A.J. and Oke, D.O. (ed.). *Proceedings of the First National*
 259 *Conference of the Forests and Forest Products Society* held at the Federal University of
 260 Technology, Akure, Ondo State between 16th and 18th of April, 2008. Pp 95-98

261

262 Akinrinde, E. A., Obigbesan, J. A., and Onanuga, O. A., (2005). Extractable micronutrient in
 263 relation to physiochemical properties of soil in four ecological zones in Nigeria. *Global*
 264 *Journal of Pure and Applied science* 2. (11), 197-200.

265

266 Broadhead, J., Leslie, R., 2007. Coastal protection in the aftermath of the Indian Ocean
 267 tsunami: what role for forests and trees? In: Proceedings of an FAO Regional Technical
 268 Workshop, Khao Lak, Thailand, 28–31 August 2006, FAO, Thailand.

269

270 Burkill HM (1999). The useful plants of West Africa. Vol. Calesal piniaceae Royal botanic
 271 Gardens. Longman publishing U.K.

272

273 Cuesta B, Villar-Salvador P, Puertolas J, F. Jacobs D, M. Rey Benayas J. (2010). Why do
 274 large, nitrogen rich seedlings better resist stressful transplanting conditions? A physiological
 275 analysis in two functionally contrasting Mediterranean forest species. *Forest ecology and*
 276 *management* [Internet].;260:71-78. Available from:
 277 [http://80.24.165.149/webproduccion/PDFs/10ART 10.PDF](http://80.24.165.149/webproduccion/PDFs/10ART%2010.PDF)

278

279 Daldoum D.M.A. and Hammad G.H.(2015): Performance of *Acacia senegal* (L.) wild
 280 seedlings growth under some tree manures and NPK fertilizers in nursery Site. *Journal of*
 281 *Forest and Environmental Science* Vol. 31, No. 4, pp. 303-311

282

283 Etukudo I. (2000). Forests; our divine treasure. Dorand publishers, Uyo.

284

285 Ezenwaka J, Ekeke B. A., Oyebade B. A. and Anyadiegwe AO (2004). Identification and
286 evolution of some threatened indigenous medical plants of the tropical rainforest of Eastern
287 Nigeria. *J. Conserv.* 2 (122): 131-141. F.P.L.G. Ibadan.

288 Hamilton, L.S. (2008). Forests and water – a thematic study prepared in the framework of the
289 Global Forest Resources Assessment 2005. Forestry Paper #155. Rome, Italy. 78 p.
290
291

292 Key R. W. J. (1989). Trees of Nigeria: a revised version of Nigeria trees vol. 1 & 2.
293

294 Lata, N. (2013) . The impact of water hyacinth manure on growth attributes and yields in
295 Coriandrum. *IOSR Journal Of Enviromental Science, Toxicology and Food*
296 *Technology.*;5(3):04-07.

297
298 Mtambalika K., Munthali C., Gondwe D. and Missanjo E.(2014): Effect of seed size of
299 *Azalia quanzensis* on germination and seedling growth. *International Journal for Forestry*
300 *Research.* 2014;5.

301

302 Nwoboshi L. C. (1985). Teak (*Tectona grandis* Linn. F.) ten year after thinning. *Nig.J.*
303 *Forestry* 15:76-79.

304

305 Okeke R. E. (1996). Comparative anatomy of trees in Nigeria: *Azalia africana* species.
306

307 Osoro N. (2014). Effects of Water Hyacinth (*Eichhornia crassipes* [Mart.] Solms) Compost
308 on Growth and Yield Parameters of Maize (*Zea mays*). *British Journal of Applied Science &*
309 *Technology.*;4(4):617-633.

310
311
312 Palgreave K.C. (2002), Trees of Southern Africa. (3rd Ed). Revised by Coates MC. Struik
313 Publishers, Cape Town.

314

315 Razaq M., Zhang P. and Sheng H. (2017). Influence of nitrogen and phosphorus on the
316 growth and root morphology of *Acer mono*. *PLoS one.* 12(2):8.
317

318 Salim E. and Ullsten O. (1999). Our forest: our future. A paper of the world commission on
319 forest and sustainable development, Cambridge Press, U.K.

320
321 Smyth, A. J., and Montgomery, R. F., (1962). Soil and landuse in central western Nigeria.
322 Government of Western Nigeria, Ibadan, Nigeria.
323

324 Talkah A. (2015). Effect of Organic Fertilizer Water Hyacinth on the Growth and Production
325 Plant Taro (*Colocasia esculenta* L.). *Journal of Environment and Earth Sciences* [Internet].;5
326 (22): 5. Available from
327 :<http://www.iiste.org/Journals/index.php/JESS/article/view/27595/28309>.

328

329 Uddin M.B., Mukul S.A., Arfin Khan M.A.S. and Hossain M.K. (2012). Effects of organic
330 manure on seedlings growth and nodulation capabilities of five popular leguminous
331 agroforestry tree components of Bangladesh. *Journal of Forestry Science* 28 (4): 212 - 219

332

333 .

334

UNDER PEER REVIEW