

## Original Research Article

# EVALUATING EFFICIENCY OF SAMPLING SCHEMES IN TROPICAL NATURAL FORESTS: REVIEW AND SIMULATION EXPERIENCE FROM KENYA

### ABSTRACT

Forest measurements, especially in natural forests are cumbersome and complex. 100% enumeration is costly and inefficient. Thus a good sampling scheme must be used. But different sampling schemes have different levels of reliability and cost associated with them. This study therefore sought to find out a sampling scheme for use in natural forests that would be reliable, efficient and with less effort. Forty-eight sampling schemes (each combining sampling intensity (5, 10, 20, 30%), plot size (25, 50, 100, 400 m<sup>2</sup>) and sampling technique (simple random sampling, systematic sampling along North-South and along East-West orientations) were generated for testing estimates of species diversity, forest regeneration and density measures through simulations using R-software. Sampling error and effort (hours spent per hectare) were used to measure efficiency of each sampling scheme in relation to actual values. Though forest sites differed in composition and structure, cost of sampling increased with decreasing plot size regardless of the attribute. Inventory accuracy increased with decreasing plot size. Optimum plot size to capture inherent population variability for regeneration, density and basal area were found for each forest type. Different sampling schemes were ranked for relative efficiency through simulation techniques, using regeneration density as an example. Generally, both random and systematic sampling-based sampling schemes were found to be effective. Apart from in montane forest, sub-sampling in 1 ha unit gave reliable results. The simulation approach used in this study is potentially useful in developing sampling protocols for complex natural forests. The 1ha-forest inventory method was found inevitable for regeneration assessment in montane forests, but flexible in other forest types where slope gradient was negligible.

**Key words:** sampling efficiency, optimum plot size, inventory protocol, regeneration assessment, R software

### 1. INTRODUCTION

Assessments of tropical natural forests are often constrained by lack of sampling protocols of known reliability. This paper reports findings on evaluation of efficiency of sampling schemes in tropical natural forests based on data from typical tropical forests and woodlands in Kenya, simulated sampling designs to capture such data and existing literature. The study was conceived based on the premises that: (i) Forest assessment studies are complex in the context of tropical mixed natural forests and in the wake of changing roles of forests and tree resources due to dynamic socio-ecological and economic situations; (ii) there are many research initiatives undertaken in forest resources assessment, but studies on efficiency and harmonization of sampling methodologies in forestry are rare; (iii) natural forests and woodlands are today recognized as critical assets for livelihoods sustenance for many people, biodiversity conservation, economic development and climate moderation for which quality information is mandatory in order to guide strategic and management plans; however, there is lack of scientifically tested and locally adapted

**Comment [J1]:** Replace words that are already in the title. The article does not evaluate software R, so substitute as well.

tools to be used in generating the much needed knowledge for those complex and diverse ecosystems; and (iv) based on the existing knowledge in forest sampling techniques, on past practices in forest assessments as well as the current computer technologies, research on efficiency of sampling schemes (accuracy, precision and cost) is achievable. A research was designed seeking to fill the aforementioned gaps by seeking to establish optimum sampling schemes for selected forest attributes with known accuracy and precision. This publication describes a detailed approach used in determining and comparing relative efficiency of various sampling schemes. In this context, a sampling scheme is a framework integrating specific sampling design, intensity and plot size. The approach was deductive, starting from the known situation (true population parameters) to generate scientific approximations (estimated population parameters) through a range of statistical procedures. Various sampling units (plot sizes), sampling intensities (sampling fractions) and spatial distribution of sampling units (sampling designs) were arrived at using R software. Real time data were collected from the field through field surveys by skilled personnel and using approved equipment. Relative accuracy and efficiency of random and systematic sampling designs with four plot sizes (25, 50, 100 and 400 m<sup>2</sup>) and varying intensities (5, 10, 20 and 30%) were investigated with reference to full-cover one-hectare inventory in tropical rain forest, moist lower montane forest and dry woodland forest types in Kenya. It was hypothesized that sampling efficiency (accuracy and precision) for regeneration, species diversity and forest structure differed among individual sampling schemes across forest types, and varied with plot sizes and sampling intensities.

**Comment [J2]:** Begin the introduction by addressing the theory of study. Objectives, assumptions and hypotheses should be removed by the end of the introduction, after placing the reader on the topic and importance of the study.

**Comment [J3]:** It is a manuscript at this time, not yet a publication. To review

**Comment [J4]:** Describe existing sampling methods and protocols, positives and negatives. Characterize in the introduction.

**Comment [J5]:** Methodological aspects should be removed from the introduction to the material and methods section

### 1.1 Review of sampling experiences and efficiency in natural forests

Field studies through sampling are often combined with most commonly used remote sensing technologies to accelerate assessment of forest resources [1, 2, 3, 4, 5, 6, 7]. Commonly used field sampling techniques in tropical forest ecosystems include walk trails, transects and plots to characterize tree species diversity, vegetation types, wildlife richness, forest structure and regeneration [5, 8] as well as to study allometric relationships for modelling forest growth and yield including evaluation of site quality [9, 10]. Sampling strategies in tropical forests are dictated and challenged by such factors as ragged terrain, abundant wildlife, expansiveness of the area and scarcity of baseline data e.g. checklists of indigenous species.

**Comment [J6]:** Clarify whether the methods in question are for vegetation or some other taxonomic group

Different researchers in Kenya have used varied plot sizes, e.g. 20 m x 10 m [5, 11], 10 m x 10 m [12]. In addition, sub-sampling using nested smaller plots within the large units is often applied in assessing forest regeneration and other plot features [4, 13, 14]. Sampling and seedling individuals are counted from different sub-plot sizes e.g. 40 m<sup>2</sup> and 20 m<sup>2</sup>, respectively. The above sampling approach enables the collection of useful information on multiple attributes from forests in a short time. The collected plot-level data reveal actual state of forest conditions e.g. regeneration, recruitment, structure, diversity, disturbances [4, 15, 16, 17, 18, 19, 20]. Integrating use of aerial photographs and field sample plots along altitudinal changes provides data for the description of montane forest vegetation [e.g., 21]. In summary, a mix of different plot shapes, plot sizes and sampling intensities have been applied in different studies in forestry but at the subjective will of different researchers and with no justification nor indication of any possible impact such mix would have on the reliability (accuracy and precision) of the findings. Ecological and socioeconomic factors are increasingly becoming important in contemporary forestry in addition to forest biophysical attributes [22, 23]. The emerging new demands dictate the need to develop tools to collect adequate data efficiently and generate required knowledge to guide sustainable management [14, 24]. To capture quality data from natural forests, different tools and methods commonly used in forest inventory must be well combined [25] and planners and managers of forests and allied resources must have the ability to identify suitable methods to produce the needed data.

## 1.2 Efficiency of inventory methods

Different plot sizes, sampling intensities, sampling techniques have been applied in mixed tropical forests with no indication of the efficiency or quality of methods used. However, multistage sampling techniques are known to increase efficiency in forest inventory [26, 27]. Past studies suggest that a plot size of one hectare is a suitable as a sampling unit [e.g., 28]. Smaller plots have been adopted in some forest vegetation studies [e.g. 29,30]. In vegetation studies, fewer but larger plots are documented to perform better than many but small plots; but there is always need to strike a balance between the cost and precision or accuracy when fixing the required sampling intensity [27]. Although subdividing any forest estate into 1-ha-inventory units is a common and agreeable practice, this study was designed to explore whether or not there could be any opportunity to sub-sample this standard unit to reduce the cost of inventory, and at the same time, achieve statistically similar or higher accuracy of estimates on-per-hectare basis.

## 2. MATERIALS AND METHODS

### 2.1 Study Sites

Three selected tropical natural forests in Kenya: Kakamega tropical rain forest (TRF), Mount Elgon moist montane forest (MMF) and Loruk dry woodland forest (DWF) were used to develop evaluation protocol for evaluation of sampling efficiency in complex forests. Figure 1 shows studied forests and sites. These forests reflect environmental gradients (climatic, topographic and anthropogenic disturbance); from low rainfall dry vegetation zone to high rainfall humid zone and lower montane moist forest zone (Table 1).

**Comment [J7]:** Add geographic coordinates of each area

**Table 1: Location, Elevation and climate characterising Study Sites in Kenya**

**Comment [J8]:** Add information about the duration of the study, in all titles

Features	TRF – Kakamega forest site	MMF - Mt. Elgon forest site	DWF –Loruk Dry woodland Site
i. Mean rainfall (mm yr <sup>-1</sup> )	1971-2000	1460–1622	629
ii. Wettest month (mm)	January (61)	May (231)	May (92)
iii. Driest month (mm)	May (273)	January (41)	February (21)
iv. Altitude (m a.s.l.)	1580	2000-2060	987
v. Mean annual temperature	20.4 °C	15.2–18.0 °C	23.7 °C
vi. Average warmest month	February (21.3°C)		March (24.8°C)
vii. Average coldest month	July (19.3°C)		August (22.5 °C).
viii. Disturbance history	Moderate logging	Extensive Logging	Livestock grazing
ix. Climate type	<i>Tropical Humid and warm</i>	<i>Temperate Moist and warm</i>	<i>Dry Tropical climate t</i>

**Sources:** [4, 17, 31, 32, 33, 34].

### 2.2 Field methods

#### 2.2.1 Forest unit of reference

From inside each forest, a one-hectare (100 m x 100 m) forest unit was selected, at least 500 m from forest edge. It was referred to as forest unit of reference and represented the “studied populations”

for different forest attributes of interest. Field data from these units were used as “controls” against which relative efficiency of each of the sampling schemes was compared and evaluated. The one-hectare unit was subdivided into smaller units during field data collection as illustrated in Figure 2.

### 2.2.2 Field work organization

Data collection over 1-ha forest unit of reference was done in the field to determine “true” values of forest attributes. It was achieved by establishing 10m x 10m temporal field plots to be subdivided into four 5 m x 5 m subplots to ensure accurate field observations of forest attributes from seedling stage were made and recorded. Complete enumeration of forest attributes was carefully and systematically done in four hundred 5 m x 5 m smallest units of data compilation. To enhance accuracy of observations on small sized individuals of the regeneration (seedlings), search and counts were done within 1 m x 1 m subplots, one after the other, within the 5 m x 5 m plot. Pre-prepared field data collection sheets were used and filled manually by trained field assistant. Field measurements and observations were later entered in MS Excel spreadsheets. The labelling was done for each 5 m x 5 m plot with an identification number for easy retrieval (see illustration in Figure 2). The largest plot size we tested in the sampling study was 20 m x 20 m. Each data entry was linked to a uniquely coded 5 m x 5 m plot. Data for plots larger than 5 m x 5 m were obtained through computer simulation using R Software by collapsing boundaries and merging adjacent smaller plots as applicable: from 5 m x 5 m plots, 5 m x 10 m and 10 m x 10m plots were formed. Merging adjacent 10 m x 10 m plots formed 20 m x 20 m plots. Merging of smaller plots was automatically associated with collating records they contained. Sums, averages and other computations were done for different plot sizes using R software modules. Similar data would otherwise be obtained in practice from field activity.

**Comment [J9]:** Are the areas private or governmental? This must be informed and the authorizations for the research must be presented. As scientists we can not invade any space, public or private, to conduct our studies.

**Comment [J10]:** Describe the environmental attributes measured. Has any forest attribute been assessed? Richness, abundance, spatial distribution? Inform and detail.




Figure 1: Location of Study Sites in Kenya




Figure 2: Example of field arrangement for four hundred 5m by 5m subplots in the 100 m x 100 m forest unit (Horizontal: E-W; Vertical: N-S).

Each cell with a number represent a coded 5 m x 5 m plot for easy data set identification, entry, storage, retrieval and use in sampling simulation activity. First three digits denote the forest site (122-Kakamega; 111- Mt Elgon; 131- Loruk). The subsequent digits represent serial plot number within the 100 m x 100 m frame.

### 2.2.3 Sampling designs

Sampling design or method is the pattern of distribution of sampling units over the sampling frame (population). Three basic designs were tested in each forest type: Simple random sampling (abbreviated as SRS), systematic sampling along vertical transect facing North – South direction (systematic sampling design abbreviated as SSV), and systematic sampling along horizontal transect facing East – West direction (systematic sampling design abbreviated as SSH). A fourth design was uniquely applied in the montane forest to assess the effect of diagonal transect across the slope gradient. Systematic sampling along diagonal transect was abbreviated as SSD. Figure 3 illustrates the three different transect directions along which plots can be systematically located. The number of plots selected and used in each design varied depending on the plot size and sampling intensity (Table 2).

**Table 2. Sample sizes and distribution of sample plots among different sampling schemes**

Sampling Design	Sampling intensity (n /N %) <sup>1</sup>	Plot sizes			
		5 m x5 m (25 m <sup>2</sup> )	10 m x 5 m (50 m <sup>2</sup> )	10 m x10 m (100 m <sup>2</sup> )	20 m x 20 m (400 m <sup>2</sup> )
SRS	5	20	10	5	1
	10	40	20	10	2
	20	80	40	20	5
	30	120	60	30	7
SSH	5	20	10	5	1
	10	40	20	10	2
	20	80	40	20	5
	30	120	60	30	7
SSV	5	20	10	5	1
	10	40	20	10	2
	20	80	40	20	5
	30	120	60	30	7

SRS = Simple random sampling; SSH = Systematic plot sampling along horizontal transect;

SSV = Systematic plot sampling along vertical transect;

<sup>1</sup>n = sample size (no. of sample plots selected from one-hectare forest ecosystem);

N = population size (total number of plots in a one-hectare forest)

A factorial combination of sampling design (3 levels), sampling intensity (4 levels) and plot sizes (4 levels) defined the 48 sampling schemes that were tested and compared for their efficiency.

#### 2.2.4. Sampling frame, sampling schemes designing and administration

The sampling frame was made of the sampling units i.e. plots in the one-hectare forest unit of reference. Population size (N) varied between 25 and 400 depending on the plot size: 400, 200, 100 and 25 units for 5 m x 5 m, 10 m x 5 m, 10 m x 10 m and 20 m x 20 m plot size, respectively. A sampling scheme was defined by the combination of three elements: sampling design (D), sampling intensity (I) and plot size (S). Each scheme was applied and evaluated for efficiency (combining accuracy, precision and cost) on each forest type. Sampling was performed on each population of selected attributes in the forest unit of reference, applying 48 different schemes in a simulated framework in R Software (Table 2).

#### 2.3. Assessed forest variables and derived attributes

Key attributes of interest included components of forest structure, composition and regeneration which are of high ecological, silvicultural and conservation significance [354, 36]. Forest canopy height was measured to the nearest m from each 5 m x 5 m unit using suunto hypsometer. Tree diameters at breast height [10, 37] were measured using callipers to the nearest mm and cm for saplings and trees, respectively. Light screening efficiency in the forest was determined at the plot centre, using a 1m x 1m transparent polythene fixed on a wooden frame and subdivided in 100 square grid, [38]. A canopy gap unit was any space measuring 5 m x 5 m or more, devoid of tree canopy cover. Tree species were identified based on dendrology documents [e.g. 39], herbarium specimens or local names. Inventory effort to complete work within a plot (duration; [40] was recorded in minutes, using a watch chronometer. For each 5 m x 5 m plot, tree seedling counts were done systematically and tallied progressively from 1 m x 1 m subplots. A field team of 4-people (1 supervisor, 1 skilled technical staff and 2 field assistants) was used.

**Comment [J11]:** Was any material collected for identification? where was the identification made? in some laboratory? in some research institution? To describe



### 3. RESULTS AND DISCUSSION

#### 3.1 Evaluating efficiency of sampling schemes for forest structure studies

Screening schemes aimed at finding sampling protocols that reduce sampling error and enhance accuracy [41]. Accuracy levels of different sampling schemes were compared for each assessed attribute, and a checklist of best schemes was provided for each forest type. A desired precision level (expressed as a percentage of the mean; Error %) was set to be  $\leq 25\%$ . This error was adequate for inventories targeting multiple attributes, e.g. in tropical forests. Assessments of simpler ecosystems e.g. forest plantations, use acceptable error  $< 25\%$  [42]. Similarly, diagnostic inventories focusing on one or two forest species also require use of acceptable error  $< 25\%$  [41].

##### 3.1.1 Reference population inherent variability

Population mean variance was based on 100 % intensity for each plot size (Eq. 1). The smallest mean variance for each attribute was identified for each forest type.

$$\text{Population mean variance} = \frac{\sigma^2}{N} \dots\dots\dots(\text{Eq. 1})$$

where,  $\sigma^2$  = population variance; N = Total number of plots per ha (varied with plot size).


**Comment [J12]:** This is method. Data Analysis, Remove for Methods Section

The plot size that produced smallest mean variance per hectare (Table 3) was considered as the most precise and accurate. The number of trees  $\text{ha}^{-1}$  exhibited low inherent variability with 5 m x 5 m plot size for all the three forest types. However, the measure of variability in the tropical rain forest was highest (moist montane forest: 13%, dry woodland forest: 0.2%). Basal area mean variance was minimized with different plot sizes across the forest types: largest (20mx20m) for the montane forest, 10mx10m in tropical rain forest and smallest (5mx5m) in dry woodland forest (Table 3). It implies that large diameter trees are more scattered in Mt Elgon moist montane forest than in other forest types (TRF and DWF), thus requiring larger plot size to capture inherent variability. Based on the findings in Table 3 above, simultaneous assessment of the three attributes would have the sub-plots designed within 1-ha-forest unit as illustrated in Figure 3. Each forest type requires a distinct design.



**Table 3: Mean population variance for different forest attributes with complete inventory (100 % intensity) of a one-hectare forest unit of reference**

Forest attribute	Forest	Smallest $\sigma^2/N$	Inventory cost (ha hr <sup>-1</sup> )	Hours ha <sup>-1</sup>	Basic unit of data compilation (m x m)
Seedlings ha <sup>-1</sup>	TRF	1,726,559.16	0.02	50	5x5
	MMF	223,003.91	0.02	50	5x5
	DWF	3,033.49	0.04	25	5x5
Stand density ha <sup>-1</sup>	TRF	5,239.46	0.02	50	5x5
	MMF	2,848.56	0.02	50	5x5
	DWF	2,427.01	0.04	25	5x5
Basal area / ha	TRF	46.967955	0.08	12.5	10x10
	MMF	7.148552	0.41	2.4	20x20
	DWF	0.050566	0.04	25	5x5



**Figure 3. Plot sizes and lay outs for simultaneous inventory of seedlings, trees and basal area in 1 ha – forest unit with minimization of inherent population variability as a controlling factor for each attribute: S = seedlings counts (< 1 cm dbh); T = trees counts (> 1 cm dbh); and BA = Basal area (m<sup>2</sup> ha<sup>-1</sup>) in Tropical rain forest, Moist montane forest and Dry woodland forest in Kenya.**

### 3.1.2 Cost and precision of different sampling schemes: case of regeneration assessment

Sample variance and standard error of mean (SE) for each sampling scheme (Eq. 2) were computed before calculating the sampling error as a percentage of the mean (Eq.3) [2, 41, 43, 44]. The sampling error percent, also referred to as uncertainty level [2], measured precision of sampling schemes. The smaller the uncertainty around the sample mean, the more precise was the sampling scheme. Figure 4 shows that precision in assessing seedlings per hectare increased with decreasing plot size.


$$SE = \frac{s}{\sqrt{n}} \dots\dots\dots(Eq. 2)$$

Where s = sample variance for the sampling scheme; n = sample size

$$\text{Sampling error \%} = \text{uncertainty \%} = \frac{SE \times t}{\bar{x}} \times 100 \dots\dots\dots(Eq. 3)$$

Where t = Student's t value obtained for each sample size from t-table with  $\alpha = 5\%$ .


**Comment [J13]:** This is method. Data Analysis, Remove for Methods Section



**Figure 4:** Effect of plot size on precision level in forest inventory (e.g. seedlings at 100 % intensity)

**Comment [J14]:** Format. Delete external border and format points

Cost-efficiency of sampling was measured through sampling effort on-per-hectare basis. Sampling effort increased with decreasing plot size (Figure 5). With 100% sampling intensity, larger plot sizes led to cheaper inventory in each forest type.



**Figure 5. Relationship between sampling effort and plot size with complete forest inventory in different forest types (Tropical rainforest –TRF, Moist montane forest – MMF and Dry woodland forest – DWF) – case of seedlings ha<sup>-1</sup>**

**Comment [J15]:** Format. Delete external border and format lines

**Comment [J16]:** Richness sampled, describe

Results on uncertainty levels and cost-efficiency associated with various sampling schemes for seedling assessment in three forest types are summarized below (Table 5):

- Most reliable sampling design in TRF was systematic sampling along transects facing north-south with 25 to 50 m<sup>2</sup> plot sizes, and 30 % sampling intensity (95% CI uncertainty level < ± 25%). However, the cost of the larger plot size was 50% lower that of smaller plot. The most optimum sampling scheme (protocol) in tropical rain forest is therefore SSV-5 mx10 m – 30%.
- In the dry woodland forest, sub-sampling one hectare unit was found possible for both random and systematic sampling (uncertainty levels were between ± 15% and ± 25%). Overall, systematic sampling with 30% intensity and 5 m x 10 m plot is most preferred due to practical field advantages over random sampling.
- There was no reliable sampling scheme for montane forest (all uncertainty levels < 25% required 100 % intensity). In this context, complete inventory over 1-ha is inevitable. Minimum plot size is 100m x 100m. Compiling seedlings data in sub-plots influenced uncertainty levels between < ± 14% and < ± 25%. Though it appears cheaper to use 20 m x 20 m sub-plot as data compilation unit, 10mx10m sub-plot size strikes a better balance between precision and cost.

**Table 4: Forest inventory schemes for seedlings in selected natural forests in Kenya**

Forest	Sampling design	Plot size	Sampling intensity	Sampling effort		Regeneration sampling error
				hours/ha	ha/hour	
TRF	SRS	25	100	50.42	0.02	15.36
TRF	SRS	50	100	25.28	0.04	16.17
TRF	SRS	100	100	12.47	0.08	18.40
TRF	SSH (E-W)	50	5	26.00	0.04	22.51
TRF	SSH (E-W)	25	100	50.42	0.02	15.36
TRF	SSH (E-W)	50	100	25.28	0.04	16.17
TRF	SSH (E-W)	100	100	12.47	0.08	18.40
TRF	SSV (N-S)	25	30	50.89	0.02	20.92
TRF	SSV (N-S)	50	30	25.17	0.04	23.53
TRF	SSV (N-S)	25	100	50.42	0.02	15.36
TRF	SSV (N-S)	50	100	25.28	0.04	16.17
TRF	SSV (N-S)	100	100	12.47	0.08	18.40
DWF	SRS	400	10	1.17	0.86	-
DWF	SRS	25	20	21.75	0.05	24.40
DWF	SRS	50	20	11.00	0.09	21.42
DWF	SRS	25	30	21.83	0.05	18.39
DWF	SRS	50	30	11.11	0.09	20.62
DWF	SRS	100	30	5.72	0.17	21.18
DWF	SRS	25	100	22.38	0.04	9.87
DWF	SRS	50	100	11.35	0.09	10.67
DWF	SRS	100	100	5.65	0.18	12.12
DWF	SRS	400	100	1.45	0.69	18.64
DWF	SSH (E-W)	50	10	11.33	0.09	21.09
DWF	SSH (E-W)	25	20	22.17	0.05	21.31
DWF	SSH (E-W)	50	20	11.25	0.09	21.94
DWF	SSH (E-W)	25	30	22.00	0.05	18.76
DWF	SSH (E-W)	50	30	11.50	0.09	19.12
DWF	SSH (E-W)	25	100	22.38	0.04	9.87
DWF	SSH (E-W)	50	100	11.35	0.09	10.67
DWF	SSH (E-W)	100	100	5.65	0.18	12.12
DWF	SSH (E-W)	400	100	1.45	0.69	18.64
DWF	SSV (N-S)	25	20	22.00	0.05	22.82
DWF	SSV (N-S)	50	20	10.83	0.09	24.39
DWF	SSV (N-S)	25	30	21.89	0.05	17.97
DWF	SSV (N-S)	50	30	10.89	0.09	18.39
DWF	SSV (N-S)	100	30	5.44	0.18	19.76
DWF	SSV (N-S)	400	30	1.28	0.78	19.73
DWF	SSV (N-S)	25	100	22.38	0.04	9.87
DWF	SSV (N-S)	50	100	11.35	0.09	10.67
DWF	SSV (N-S)	100	100	5.65	0.18	12.12
DWF	SSV (N-S)	400	100	1.45	0.69	18.64

Table 4 (continued)

Forest	Sampling design	Plot size	Sampling intensity	Sampling effort		Regeneration sampling error
				hours/ha	ha/hour	
MMF	SRS	25	100	40.68	0.02	14.74
MMF	SRS	50	100	20.10	0.05	15.62
MMF	SRS	100	100	9.98	0.10	18.22
MMF	SRS	400	100	2.43	0.41	24.04
MMF	SSH (E-W)	25	100	40.68	0.02	14.74
MMF	SSH (E-W)	50	100	20.10	0.05	15.62
MMF	SSH (E-W)	100	100	9.98	0.10	18.22
MMF	SSH (E-W)	400	100	2.43	0.41	24.04
MMF	SSV (N-S)	25	100	40.68	0.02	14.74
MMF	SSV (N-S)	50	100	20.10	0.05	15.62
MMF	SSV (N-S)	100	100	9.98	0.10	18.22
MMF	SSV (N-S)	400	100	2.43	0.41	24.04

### 3.1.3 Relative efficiency of different sampling schemes based on cost, precision and accuracy: case of regeneration assessment

Equation 4 was applied to each applicable sampling scheme as shown below to find the best performing ones (Figure 6).

$$\text{Efficiency \%} = \frac{\sigma_1^2 \times C_1}{s_1^2 \times C_1 / n_1} \times 100 \dots \dots \dots \text{(Equation 1)}$$

Where  $s_1^2$  = sample variance for the sampling scheme;  $n_1$  = sample size (i.e. no. plots);  $C_1$  = time (hours) spent on measuring variables (ie sampling effort or cost  $ha^{-1}$ );  $\sigma_1^2$  = population variance for reference for the variable;  $C_1$  = actual total cost of measuring variables in one-hectare forest unit of reference with the selected plot size;  $N_1$  the population size (number of plots per ha, varying with plot size).

**Comment [J17]:** Method

Results indicate that systematic counting of seedlings in 5 m x 5 m plot sizes over 30% of the entire one hectare (SSV-5mx5m-30%) was the most efficient scheme for Tropical rain forest regeneration inventory among other schemes (Figure 6). Using measures of sampling efficiency, best schemes were identified as **SSV-5mx5m-30%** (83% efficiency) and **SSH-5mx10m-5%** (80%) for TRF. In DWF, best performing schemes were **SRS-10mx10m-30%** (91 % efficient), **SSV-10mx10m-30%** (75 %) and **SSV-5mx10m-30%** (74%). For MMF, all evaluated sampling schemes with intensity < 100% were not efficient enough (<50%). Seedling surveys should be done over the entire 1 ha, subdivided into 5m x 5 m sub-plots (systematic inventory along transects using 5mx5m -100% design).


## 4. CONCLUSION

This study demonstrated that screening sampling schemes and developing sampling protocols are achievable using good empirical field data and computer programming software such as R. Evaluation of possibilities to sub-sample one hectare area or not, optimum sampling unit, intensity and design between random and systematic sampling were achieved in two closed canopy natural forests and open dry woodland forest in Kenya; thereby guiding development of sampling protocols for regeneration, density and basal area studies. The 1ha-forest inventory method was found inevitable for regeneration

**Comment [J18]:** The study does not clearly indicate which indicators were used. More important than talking about design and software, it is important to address environmental and ecological indicators and how they respond to the model being tested. More importance was given to the method than to the object studied.

assessment in montane forest; but flexible in other forest types where slope gradient was negligible. Slope gradient is suspected to influence inventory methods for regeneration.

UNDER PEER REVIEW






Figure 6: Relative efficiencies (cost, precision and accuracy combined) of candidate sampling schemes in assessing forest regeneration (no. seedlings ha<sup>-1</sup>) in tropical rain forest (TRF), moist montane forest (MMF) and dry woodland forest (DWF), Kenya



## REFERENCES

1. Nilson M, Folving S, Kennedy P, Puumalainen J, Chirici G, Corona P, *et al.* Combining remote sensing and field data for deriving unbiased estimates of forest parameter over large regions. In, Corana, P., Kohl, M., & Marchetti, M. (Eds). *Advances in Forest Inventory for sustainable Forest Management and Biodiversity Conservation*. 2003.
2. Lackmann S. *Good practice in designing a forest inventory*. Lesson 8. Regional Workshop: "Capacity Development for Sustainable National Greenhouse Gas Inventories – AFOLU sector (CD-REDD II) Programme, Quito, Ecuador. 2011.
3. Asthana DK, Asthana M. *Environment: problems and solutions*. Ramnagar, New Dheli: S. Chand and Company Ltd. 1998.
4. Hitimana J, Kiyiapi J, Njunge J. Forest structure characteristics in disturbed and undisturbed sites of Mt Elgon moist lower montane forest, western Kenya. *Forest Ecology and Management*. 2004; 194, 269-291.
5. Hitimana J, Kiyiapi J, Kisioh H, Warinwa F, Lenaiyasa P, Mayienda R, *et al.* Participatory ecological assessment of Kirisia forest reserve, Samburu, Kenya. *AWF Technical Paper Series*, April 2009. Nairobi: African Wildlife Foundation. 2009.
6. Ucar Z, Bettinger P, Merry K, Siry J, Bowker JM, Andakbuluta R. A comparison of two sampling approaches for assessing the urban forest canopy cover from aerial photography. *Urban Forestry & Urban Greening*. 2016; 16, 221–230.
7. Nowak DJ, Crane DE, Stevens JC, Hoehn RE, Walton JT, Bond J. A ground-based method of assessing urban forest structure and ecosystem services. *Arboric. Urban For*. 2008; 34, 347–358.
8. Mutiso FM, Cheboiwo J, Kiyiapi JL, Sang FK, Hitimana J. Comparative Study of Sampling Methods for Efficient Diagnosis of Health Status of Selected Natural Forest Ecosystems in Kenya. *Journal of Natural Sciences Research*. 2016; 5(2).
9. Weiskittel AR., Hann DW, Kershaw Jr, JA, Vanclay J. *Forest growth and yield modeling*. Wiley-Blackwell, Oxford, UK. 2011.
10. Burkhart HE, Tome M. *Modeling forest trees and stands*. Springer Science, London. 2012.
11. Warinwa F, Mwaura F, Kiringe JW, Ndubi AO. Land Cover Dynamics in the Kirisia Forest Ecosystem, Samburu County, Kenya. *Advances in Remote Sensing*. 2016; 5: 168-182.
12. Ronoh DK. *Anthropogenic disturbance indicators and population structure of *Zanthoxylum gillettii* (De Wild Waterman) and *Prunus Africana* (Hook. f.) Kalkman in South West Mau Forest, Kenya*. Unpublished master's thesis, University of Kabianga. 2016
13. Ronoh DK, Sirmah PK, Hitimana J, Mullah CJA. Variation in Regeneration Density and Population Structure of *Prunus africana* Across Human Disturbance Gradient in South West Mau Forest, Kenya. *International Journal of Natural Resource Ecology and Management*. 2018; 3 (1): 1-8.
14. Kiruki HM, van der Zanden EH, Gikuma-Njuru P, Verburg PH. The effect of charcoal production and other land uses on diversity, structure and regeneration of woodlands in a semi-arid area in Kenya. *Forest Ecology and Management*. 2017; 391, 282-295.
15. Clausnitzer V, Churchfield S, Hutterer R. Habitat occurrence and feeding ecology of *Crocodylus montis* and *Lophuromys flavopunctatus* on Mt. Elgon, Uganda. *African Journal of Ecology*. 2003; 41(1), 1–8.
16. Bulafu CE, Mucunguzi P, Kakudidi EK. Diversity and distribution of wild terrestrial orchids of Mt Elgon Forest National Park, eastern Uganda. *African Journal of Ecology*. 2007; 45 (3), 21–28.
17. Hitimana J, Kiyiapi J, Njunge J, Bargerei R. Disturbance indicators and population decline of logged species in Mt Elgon Forest, Kenya. *African Journal of Ecology*. 2010; 48, 699-708.
18. Hitimana J, Kiyiapi JL, Kibugi PW, Kisioh H, Mayienda R, Warinwa F, *et al.* Challenges of linking socio-economic significance and conservation value of forests In Drylands of Kenya: Case study of Kirisia Forest- Samburu pastoralists coexistence. In: Grillo, O. & Venora, G. (Eds). *Biological Diversity and Sustainable Resource Use*. 2011. Intech. Chapter 1, Pp. 1-34.

**Comment [J19]:** Less than 25% of references are from the last five years. Update.

**Formatted:** Font: Portuguese (Brazil)

19. Kiringe JW, Mwaura F, Warinwa F. Characterization of Water Source Types and Uses in Kirisia Forest Watershed, Samburu County, Kenya. *Environment and Natural Resources Research*. 2016; 6 (3).
20. Reed MS, Cokie MRJ. Effects of grazing and cultivation on forest plant communities in Mount Elgon National Park, Uganda. *African Journal of Ecology*. 2000; 38 (2), 154–162.
21. Hamilton AC, Perrott RA. A study of altitudinal zonation in the Montane Forest Belt of Mt. Elgon, East Africa. *Vegetatio*. 1981; 45, 107-125.
22. Vanclay JK. Realizing opportunities in forest growth modelling. *Canadian Journal of Forest Research*. 2003; 33, 536 – 541.
23. Pretzsch H. *Forest dynamics, growth and yield, from measurement to model*. Springer-Verlag, Berlin. 2009.
24. Nacoulma BMI, Schumann K, Traore S, Bernhardt-Roermann M, Hahn K, Wittig R, *et al*. Impacts of land-use on West African Savanna vegetation: a comparison between protected and communal area in Burkina Faso. *Biodiversity Conservation*. 2011; 20, 3341-3362.
25. Bellefontaine R, Petit S, Pain-Orcet M, Deleporte P, Bertault J. *Trees outside forests: towards better awareness*. FAO Conservation guide 35, Rome. 2002.
26. Alder D, Synnott TJ. *Permanent sample plot techniques for mixed tropical forest*. Tropical Forestry Papers, Commonwealth Forestry Institute, University of Oxford. 1992. No.25.
27. Gerwing JJ, Schnitzer SA, Burnham RJ, Bongers F, Chave J, DeWalt SJ, *et al*. Standard Protocol for Liana Censuses. *Biotropica*. 2006; 38(2), 256–261.
28. Synnott TJ. *A manual of permanent plot procedures for tropical rainforests*. Tropical Forestry Papers. Commonwealth Forestry Institute, University of Oxford. 1979; No. 14.
29. Gebreselasse AG. *Plant Communities, Species Diversity, Seedling Bank and Resprouting in Nandi Forests, Kenya*. Unpublished PhD Thesis, University of Koblenz-Landau. 2011.
30. Lee HS, Davies SJ, LaFrankie JV, Tan S, Yamakura T, Itoh A, *et al*. Floristic and structural diversity of mixed dipterocarp forest in Lambir Hills National Park, Sarawak, Malaysia. *Journal of Tropical Forest Science*. 2002; 14(3), 379-400.
31. Fashing, P., Forrestel, A., Scully, C., & Cords, M. (2003). Long-term tree population dynamics and their implications for the conservation of the Kakamega forest, Kenya. *Biodiversity and Conservation*, 13, 753–771.
32. Joetzoald R, Schmidt H. *Farm Management Handbook of Kenya, West Kenya*. II/A. 1983; pp. 309–324.
33. Kokwaro J. Conservation status of the Kakamega forest in Kenya: the easternmost relic of the equatorial rain forest of Africa. *Monographs in Systematic Botany Missouri Botanical Garden*. 1988; 25, 471–489.
34. Mutiso FM, Hitimana J, Kiyiapi JL, Sang FK, Eboh E. Recovery of Kakamega tropical rainforest from anthropogenic disturbances. *Journal of Tropical Forest Science*. 2013; 25(4), 566–576.
35. Feld CK, da Silva PM, Sousa JP, de Bello F, Bugter R, Grandin U, *et al*. Indicators of biodiversity and ecosystem services: a synthesis across ecosystems and spatial scales. 2009.
36. Hamilton AJ. Species diversity or biodiversity? *Journal of Environmental Management*. 2005; 75, 89–92.
37. Avery TE, Burkhardt HE. *Forest Measurements*, 5th Ed. Illinois: Waveland Press. 2015.
38. Brower JE, Zar JH, Von Ende. *Field and Laboratory Methods for General Ecology*, 3<sup>rd</sup> Ed. Duburques (USA): Wm. C. Brown Publishers. 1990.
39. Beentje HJ. *Kenya Trees, Shrubs and Lianas*. Nairobi : National Museums of Kenya. 1994.
40. Muchiri MN, Ngugi J, Kinyanjui M, Balazi K, Ojuang F, Nduati P, *et al*. *Improving Capacity in Forest Resources Assessment in Kenya (IC-FRA): Field Manual for Biophysical Forest Resources Assessment in Kenya*. KEFRI, Muguga. 2016.
41. Klauberg C, Vidal E, Silva CA, de M, Bentes M, Hudak AT. Sampling methods for titica vine (*Heteropsis* spp.) inventory in a tropical forest. *Annals of Forest Science* 2016; 73, 757–764.

Formatted: Portuguese (Brazil)

Formatted: Font: Portuguese (Brazil)

42. Cavalcanti FJB, Machado AS, Osokawa RT, Cunha US. Comparison of the estimated values by sampling in the characterization of the structure of a forest area in the Amazon with the information recorded in forest census. *Revista Árvore*. 2011; 35, 1061–1068.
43. Morais R, Sheuber M. Statistical precision in phytosociological surveys. In: *Proceedings of the International Symposium on Assessment and Monitoring of Forests in Tropical Dry Regions with special reference to Gallery Forests*, Nov. 1996, Brasília. 1997; 135-147.
44. Wong JLG, Thornber K, Baker N. *Resource assessment of non-wood forest products: experience and biometric principles*. FAO, Rome. 2001.

Formatted: Font: Portuguese (Brazil)

UNDER PEER REVIEW