

Original Research Article

EFFECT OF AIR-DRIED MUCUNA (*Mucuna pruriens*) LEAF MEAL IN THE DIETS OF GROWER RABBIT BUCKS ON THE PERFORMANCE, NUTRIENT DIGESTIBILITY AND ECONOMIC COST OF PRODUCTION

Comment [u1]: Performance, Nutrient digestibility and Economic cost of production of Weaner Rabbit Bucks fed diets containing dried *Mucuna pruriens* leaf meal

ABSTRACT

A twelve week feeding trial was conducted to investigate the effect of air-dried mucuna (*Mucuna pruriens*) leaf meal (MLM) in the diets of grower rabbit bucks. Mucuna leaf meal was included at 0%, 5%, 10%, 15% and 20% for T₁, T₂, T₃, T₄ and T₅ respectively. Thirty grower rabbits bucks were assigned to five experimental dietary treatments denoted T₁, T₂, T₃, T₄ and T₅ respectively in a completely Randomized Design (CRD). Each treatment had six rabbits with each serving as a replicate. Fibre analysis results showed that mucuna leaf meal had 22% cellulose, 11% acid detergent lignin, high level of 38% hemicellulose, 49% neutral detergent fibre and 33% acid detergent fibre. Results of performance showed that there were no significant differences in final body weight, daily weight gain, daily feed and protein intake. Protein efficiency ratio and feed conversion ratio were significantly higher at T₁. Nutrient digestibility by rabbit indicates significant different ($p < 0.05$) in ether extract (EE), crude fibre (CF), and crude protein (CP). EE and CF were significantly higher at T₄, and CP was highest at T₁. There was no significant difference ($P > 0.05$) in NFE across the treatments. There was also no significant difference ($P > 0.05$) for cost per kg gain. Cost of feed consumed, total cost and percentage feed cost were significantly higher ($P < 0.05$) at T₁, T₂, and T₃. Percentage drug cost was significantly better ($P < 0.05$) at T₄ and T₅, percentage cage cost and animal were statistically depressed ($P < 0.05$) at T₁, T₂ and T₃ and highest at T₅. It was concluded that mucuna (*Mucuna pruriens*) leaf meal has potentials as a valuable unconventional leguminous feedstuff. The use of mucuna *pruriens* leaf meal in the diets of rabbit bucks promoted growth best at 5% and thus carcass yield. Rabbit farmer can incorporate mucuna *pruriens* leaf meal at 5% for growth performance.

Key words: mucuna, rabbit bucks, performance, nutrient digestibility and economic of production

INTRODUCTION

Small-livestock such as rabbits have advantages of fast growth, large litter size, and short generation interval and good quality consumable and non-consumable animal products, but are faced with the challenge of feedstuff availability and affordability. In Nigeria today, the cost of feeding livestock intensively is over 70%²⁴. Based on their findings, unconventional feedstuffs

37 could play a vital role in alleviating this enormous challenge posed by intensive feeding in
38 livestock production, because they are mostly cost-free and are readily available.

39
40 The cost of feeding rabbits is high, a condition that also prevails for other Nigerian livestock
41 species¹. Less developed countries, like Nigeria, are facing serious competition between human
42 and livestock (especially, the monogastric animals) for available conventional feedstuffs³⁰. This
43 increased competition for available conventional feeds and scarcity of food have both
44 encouraged nutritionists, scientists and agriculturists to research into the use of unconventional
45 feedstuffs that are cheap, readily available and are possible substitute for more expensive protein
46 sources (groundnut cake and soybean meal) and energy sources such as maize²².

47 Forages offer a considerable potential as major source of energy, protein, minerals and vitamins
48 for herbivorous animals and are readily available¹⁹. Forages not only serve as a source of fibre
49 for rabbit, they are essential for normal functioning of the gut health and mobility; caecotrophy
50 and appetite stimulation⁹.

51
52 *Mucuna pruriens*, widely known as “velvet bean,” is a vigorous annual climbing legume
53 originally from Southern China and eastern India, where it is at one time widely cultivated as a
54 green crop¹⁰. It is one of the most popular green crops currently known in the tropics. Velvet
55 beans have great potential as both food and feed as suggested by experiences worldwide. The
56 velvet beans has been traditionally used as a food source by certain ethnic groups in a number of
57 countries. It is cultivated in Asia, America, Africa, and Pacific Islands, where its pods are used as
58 a vegetable for human consumption, and its young leaves are used for animal feed.

59
60 *Mucuna utilis* (velvet bean), a tropics legume, is little known and has a low human preference as
61 an energy/protein source but high preferences in livestock feed¹¹. It is comparable to soya bean
62 in terms of amino acid and mineral profile¹⁷. However, the use of velvet beans as a source of
63 protein for monogastrics is limited by the presence of antinutritional factors like trypsin
64 inhibitors, haemagglutinins, phytic acids, hydrocyanic acid and tannins¹².

65
66 Little has been reported on the use of mucuna leaves in the diets of rabbits. Therefore, mucuna
67 leaf meal, a potential feedstuff, could play a vital role in livestock production, hence, its effect on
68 the performance of growing rabbit bucks was investigated in this study.

69

70 MATERIALS AND METHODS

71 **Location**

72 The study was conducted at the Rabbitry Unit of the Livestock Teaching and Research Farm,
73 University of Agriculture, Makurdi, Benue State. Makurdi lies between Latitude 7⁰44'N and
74 longitude 8⁰21'E in the Southern Guinea savanna Zone, Benue State. The area has an annual
75 rainfall of 6-7 months in duration (i.e. March – October) and ranging from 508 to 1016 mm with
76 a minimum temperature range of 24.20 +1.4⁰C and maximum temperature range of 36.33
77 +3.70⁰C, respectively. The relative humidity ranges between 39.50 + 2.20% and 64.00 +
78 4.80%²⁹

Comment [u2]: Experimental site or Site of experiment should be used

79 **Processing of Mucuna Leaf Meal and Diet Preparation**

80 Mucuna leaves were sourced from within Makurdi metropolis at the back of Rabbitry Unit of the
81 Livestock Teaching and Research Farm, University of Agriculture, Makurdi, Benue State. The
82 leaves were harvested from the wild, such that yellow leaves were discarded, leaving the
83 greenish leaves for use. Harvested leaves were air-dried, turned frequently to enhance uniform
84 drying until the material became crispy. Air-dried mucuna leaves were milled using a roller
85 miller to obtain mucuna leaf meal (MLM) which was subjected to proximate analysis ⁵ and the
86 proximate composition determined. Five iso-caloric and iso-nitrogenous experimental diets were
87 formulated which contained 0%, 5%, 10%, 15%, and 20% mucuna leaf meal and 0%MLM
88 served as the control diet (Table 1).

Comment [u3]: Which period was the experiment conducted example May to August, 2018

Comment [u4]: Collection, processing and preparation of experimental diets

89

90 **Experimental Design**

91 The experimental design was completely randomized design (CRD). Thirty rabbits were grouped
92 into five with each group containing six animals and replicated six times; each animal serving as
93 a replicate. The treatment groups were balanced of their weights and randomly exposed to the
94 five dietary treatments.

95 **Experimental Animals and Management**

96 Thirty healthy grower rabbit bucks of about four to five (4-5) weeks of age were purchased from
97 farms within Makurdi metropolis for the study. Before the arrival of the rabbits, the cages,
98 feeders and drinkers were properly cleaned and disinfected using izal. On arrival, the rabbits
99 were housed individually in 40 x 60 x 40 cm³ cages having wire mesh floor, 1m above the

100 ground and acclimatized for 10 days. Standard rabbit husbandry practices including medications,
101 recommended sanitary space measures and other health practices were strictly observed
102 throughout the experimental period ¹⁵. The animals were served feed and water ad-libitum daily
103 for 84 days

104

105

UNDER PEER REVIEW

106 Table 1. Feed Composition of Grower Rabbits Diets Containing mucuna leaf meal (MLM)

Ingredients (Kcal/kg)	Experimental Diets				
	T ₁ (0%MLM)	T ₂ (5%MLM)	T ₃ (10%MLM)	T ₄ (15%MLM)	T ₅ (20%MLM)
Maize	38	38	37	34	33
Maize offal	12.15	10.15	7.15	8.15	8.15
Brewer dry grain	10	10	10	10	10
Soya bean meal	16	12	12	9	7
Rice offal	19	19	19	19	17
Mucuna leaf meal	0	5	10	15	20
Blood	2	3	3	2	2
Bone ash	2	2	2	2	2
Salt	0.3	0.3	0.3	0.3	0.3
Lysine	0.1	0.1	0.1	0.1	0.1
Methionine	0.2	0.2	0.2	0.2	0.2
Premix	0.25	0.25	0.25	0.25	0.25
Total	100	100	100	100	100
Calculated	Analysis (%)				
Crude protein	17.23	17.25	17.33	17.10	17.29
Crude fibre	13.02	13.17	13.38	13.81	13.45
ME (Kcal/kg)	2560.41	2570.24	2580.00	2567.76	2587.01
Ether extract	2.2	2.30	2.46	2.53	2.66
Calcium	1.2	1.22	1.21	1.20	1.16
Phosphorus	0.8	0.84	0.82	0.80	0.76
Methionine	0.25	0.23	0.22	0.21	0.19
Lysine	1.56	1.21	1.21	1.37	1.32

107 MLM= mucuna leaf meal, ME= metabolizable energy, T1-5= treatment 1,2,3,4 and 5.

Comment [u5]: Remove

108 **Table 2: Proximate composition of Experimental Diets (Analysed)**

Constituents	T ₁	T ₂	T ₃	T ₄	T ₅
Moisture	11.88	11.97	11.77	10.89	12.00
Ash	9.50	5.42	10.14	17.47	7.79
Ether extract	5.18	5.52	6.39	6.87	6.24
Fibre	12.81	15.01	11.19	20.80	15.41
Protein	16.94	16.39	15.42	18.56	18.23
NFE	43.60	45.69	45.12	25.43	40.34

Comment [u6]: Moisture (%), Ash (%),.....

109 NFE=Nitrogen Free Extract, T₁= 0%MLM, T₂= 5%MLM, T₃= 10%MLM, T₄= 15%MLM, and

110 T₅= 20%MLM, %= percentage, MLM= Mucuna leaf meal

111 **Proximate Composition**

112 Proximate composition of mucuna leaf meal, treatment diets as well as fecal samples collected
 113 during digestibility trial were determined using the standard methods ⁵ of Purity Laboratory Jos,
 114 Plateau State.

115
 116

Fibre Fraction

117 Fibre fraction of mucuna leaf meal was determined at Animal Nutrition laboratory Department
 118 of Animal Nutrition, Federal University of Agriculture, Abeokuta, Ogun State.

119
 120

Growth Performance Indices

121 Feed intake (FI): a known quantity of feed offered (FO) to each rabbit and the leftover feed
 122 (LOF) at the end of every week were weighed and the feed consumed was calculated by
 123 difference. That is, FI= FO – LOF

124 Body weight gain (BWG): Individual rabbits were weighed at the commencement of the trial,
 125 weekly thereafter, and at the end of the feeding trial. Total weight gain was obtained by
 126 difference between final live body weight (FW) and initial body weight (IW). That is, FW – IW.

127 Daily weight gain was determined as the total weight gain divided the number of the days the
 128 experiment lasted.

129 Feed Conversion Ratio (FCR): This is a measure of an animal's efficiency in converting feed
130 into desired output. It was calculated as the ratio of feed intake to body weight gain

131

132
$$FCR = \frac{FI}{BWG} \dots\dots\dots (1)$$

133

134 Protein Intake (PI) was calculated as follows; $FI \times \% \text{ crude protein in feed}$.

135 Protein Efficiency Ratio (PER); this expresses numerically the growth promoting value of
136 protein, it involves using the weight of protein intake in a test diet to divide the weight gain by
137 the animal on the test diet. It was calculated as follows;

138
$$PER = \frac{BWG}{PI} \dots\dots\dots (2)$$

139 **Digestibility Trial**

140 At the end of the 11th week of the feeding trial, three (3) rabbits, with live weights closed to their
141 treatment average were selected from each treatment and used for the digestibility trial. Faecal
142 collection lasted for five (5) days. During this period, nylon net were tied under individual rabbit
143 cages for daily faecal collection. Before the commencement of faecal collection, the rabbits were
144 deprived of feed for 18 hours to ensure that faecal collection corresponded to the feed offered.
145 The fresh collected faeces were weighed and oven dried at 80⁰C for 24 hours, the oven-dried
146 faeces per replicate was also weighed. At the end of the digestibility study, collected faeces from
147 each replicate were bulked, thoroughly mixed together and milled. Samples of the milled faeces
148 were stored in air tight containers for proximate analysis. Also sample of feed from each
149 treatment were taken for proximate analysis. Proximate composition was determined as outlined
150 by⁵

151

152 Digestibility coefficients were calculated using the following equation by²⁰.

153
$$\text{Apparent digestibility} = \frac{\text{Nutrient in feed} - \text{Nutrient in faeces}}{\text{Nutrient in feed}} \times 100 \dots\dots\dots (3)$$

154

155

156 **Economics of production**

157 The cost of feed ingredients and other services such as transportation and processing (milling of
158 mucuna leaf) were used to get the actual cost of each ingredient during the study. The
159 formulation for each diet was used to determine the cost of feed by multiplying unit cost (₦) of
160 each ingredient by its proportion in the diet to determine its cost contribution to the diet. The sum
161 of all the cost contribution from all the ingredients that made up each diet gave the units cost of
162 (₦) diet. Value of rabbit per kg multiply by the final weight (kg) were used to get revenue.
163 Benefit per rabbit was gotten by subtracting total cost of production from revenue. Cost benefit
164 ratio was calculated by dividing total cost of production by benefits per rabbit. Feed cost divided
165 by total cost multiplied by one hundred gives percentage feed cost. Percentage drug cost equals
166 to drug cost divided by total cost multiplied by one hundred. Cage cost divided by total cost
167 multiplied by one hundred equals to percentage cage cost and percentage animal cost was
168 determine as cost of animal divided by total cost multiplied by one hundred.

169

170 **Statistical Analysis**

171 All the data generated were subjected to the analysis of variance (ANOVA) using Statistical
172 Software (SPSS version 16) and significant differences were separated using its Duncan New
173 Multiple Range Test⁽²⁸⁾.

174

175

176 **RESULTS**

177 **Table 3:** Proximate composition and fibre fraction of mucuna leaf meal (MLM)

Composition	Percentages (%)
Dry matter	88.97
Ash	10.25
Ether extract	4.91
Crude fibre	26.54
Crude protein	26.09
Nitrogen free extract (NFE)	21.18
Fraction	%
Cellulose	22
Hemicellulose	38
Neutral detergent fibre	49
Acid detergent fibre	33
Acid detergent lignin	11

178
179 Table 3 shows proximate composition and fibre fraction of mucuna leaf meal (MLM). The value
180 11.03 for moisture in this study is lower than 12.50%¹³ but similar to 11.37%³¹. The crude
181 protein of 26.09% recorded for MLM was higher than the values of 22.94%¹³ but lower than
182 31.91% by³¹. The value of 4.91% ether extract in MLM was lower than 8.50%¹⁴ and¹³. Crude
183 fibre (26.54%) obtained is higher than 12.50% and 14.80%³¹. Ash 10.25% obtained in this study
184 was higher than 5.80% recorded by¹⁶. NFE (21.18%) recorded in this study was lower than
185 47.51%¹⁴. From the results and reports, it is noted that differences exist among report with
186 composition and these differences could be attributed to variations in processing method and
187 varieties. However, on average, the best material (MLM) is a fibrous protein source.

188
189
190

191 **Fibre Fraction of Mucuna Leaf Meal**

192 The fibre fractions are presented in table 3. Higher cellulose and acid detergent lignin (ADL)
193 lead to low degradation by the microbes. Higher percentages of hemicellulose, neutral detergent
194 fibre (NDF) and acid detergent fibre (ADF) indicates high nutrient degradation by microbes
195 which shows that mucuna leaf meal is saved for rabbit. The value of NDF 49% in this present
196 study is higher than 43% reported by ⁴, 33% ADF in this research is lower compare to 38% as
197 quoted by ⁴. The value for ADL in this work is higher than 7.1% as reported by the earlier
198 author.

199

UNDER PEER REVIEW

200 **Table 4: Growth Performance of Rabbit Bucks Fed Diets containing Different levels of Mucuna Leaf Meal (MLM)**

Parameters	Experimental Diets					SEM	P-value
	T ₁ (0%MLM)	T ₂ (5%MLM)	T ₃ (10%MLM)	T ₄ (15%MLM)	T ₅ (20%MLM)		
Final Weight (g)	1719.0	1645.8	1583.7	1459.2	1467.0	155.51	.062
Total Weight Gain (g)	1263.0	1143.6	1095.8	978.83	983.00	171.95	.083
Daily Weight Gain (g)	15.04	13.05	13.05	11.59	11.73	2.03	.073
Total Feed Intake (g)	5201.50	5737.40	5464.33	5336.67	5364.33	516.94	.590
Daily Feed Intake (g)	62.67	69.13	65.83	64.28	64.63	6.23	.589
Protein Intake (g)	10.79	11.92	11.41	10.99	11.18	1.07	.542
Protein Efficiency Ratio	1.39 ^a	1.09 ^b	1.14 ^b	1.05 ^b	1.05 ^b	0.12 [*]	.023
Feed Conversion Ratio	4.17 ^a	5.04 ^{ab}	5.09 ^{ab}	5.76 ^b	5.60 ^b	0.72 [*]	.023
Mortality Rate (%)	33.3	16.67	0.00	0.00	0.00	–	–

201

Means on the same row with different superscript are significantly different (P<0.05), SED= Standard error of mean, *=significant, T₁ = 0%MLM, T₂ = 5%MLM, T₃ = 10%MLM, T₄ = 15%MLM, T₅ = 20%ML

204 Table 4 contained growth performance of rabbit bucks fed diets containing mucuna leaf meal
205 (MLM). There were no significant differences in the final body weight (FBW), daily weight gain
206 (DWG), daily feed intake (DFI), and protein intake. Animals on T₁ (controlled diet) had the highest
207 FBW (1719.0g) and lowest in T₄ (1459.2g). The daily weight gain of 11.59-15.04g per rabbit per
208 day were lower than 17.65-18.57g/day³ and 18.00-20.00g², but was higher than 4.94-14.80g/day⁸
209 and 8.70-9.91 g/day¹⁸ who fed rabbits on different levels of groundnut haulms. The average daily
210 weight gain was better at T₁ (0% MLM), T₂ and T₃ but decreased numerically at T₄ and T₅ which
211 had the lowest figures. This implied that mucuna leaf meal had some growth suppressing effect in
212 the diets of rabbits on the weight gain. The quantity of feed consumed per rabbit per day observed
213 in this study was however quite higher than 48.83 – 52.13 g²⁵, 44.73 to 57.90 g²¹ but lower than
214 77.64 – 87.59 g and 63.89 -82.46 g⁶. Similarities (p>0.05) observed in feed intake suggest that all
215 the diets were palatable and thus accepted by the rabbits as the inclusion of MLM at varying levels
216 did not depressed feed consumption. There were no significant differences (P<0.05) in protein
217 intake and the values ranged from 10.79 to 11.92g.

218
219 Significant differences (P<0.05) occurred in the feed conversion ratio and protein efficiency ratio.
220 The values of feed conversion ratio and protein efficiency ratio (Table 3) obtained in this study
221 showed that the control diet (T₁) was better. The significant difference between the control (T₁), T₄
222 and T₅ indicate a decline in absorption with the level of mucuna in the body beyond 10%. This has
223 also manifested in the final weight. Though the final weight is statistically seminar on analysis of
224 variance, the trendy decrease in final weight clearly shows that it is not a chance occurrence but diet
225 effect.

226

227 **Table 5: Digestibility of Rabbit Bucks fed Diets containing different levels of Mucuna Leaf**
 228 **Meal (MLM)**

Parameters	Experimental Diets					SEM	P-value
	T ₁	T ₂	T ₃	T ₄	T ₅		
Ether extract	67.83 ^c	63.50 ^c	73.13 ^b	75.37 ^a	61.71 ^c	5.25 [*]	.039
Crude Fibre	27.03 ^b	21.68 ^b	10.33 ^b	56.16 ^a	21.44 ^b	8.97 [*]	.001
Crude Protein	80.44 ^a	72.77 ^b	71.07 ^c	78.82 ^b	71.26 ^c	4.11 [*]	.049
NFE	80.23	79.31	80.49	75.66	76.12	5.46	.715

229 abc Means on the same row with different superscript are significantly different (P<0.05), SEM=
 230 Standard error of mean, T₁ = 0%MLM, T₂ = 5%MLM, T₃ = 10%MLM, T₄ = 15%MLM, T₅ =
 231 20%MLM, NFE= nitrogen free extract.
 232

233
 234 Table 5 showed coefficient digestibility of nutrients by rabbit bucks fed diets containing different
 235 levels of mucuna leaf meal (MLM)
 236 There were significant (P<0.05) differences in nutrient digestibility by rabbit. Ether extract and
 237 crude fibre varied without pattern while crude protein varied first from 0 to 10%, became elevated
 238 at 15% and fell again at 20%. This was almost a trend of declined except 15% which cannot be
 239 explained. NFE was not significantly affected meaning that energy utilization was similar across the
 240 treatment. The coefficient of digestibility of crude protein which ranged from 71.07%-80.44%, was
 241 higher than 63.30 to 76.28%²³ but comparable with 70.56 to 81.31% and 72.25 to 82.88%²⁷ and²⁶
 242 respectively. This suggested good availability of crude protein in the diet with MLM up to 20%
 243 inclusion levels. High values indicated efficient utilization of protein for tissue accretion while
 244 lower values are indications of poor crude protein utilization for tissue synthesis²⁶. The crude fibre
 245 digestibility was low in T₃ (10.33%) and high in T₄ (56.16%). Coefficient digestibility of crude
 246 fibre (10.33%-56.16%) was low compared to values 27.54 to 56.36 21 and 71.00 to 82.29%²⁷

247 Ether extract ranged from 61.71% - 75.37 % and were lower than 71.12% - 78.43% as reported by
248 ²⁶. High digestibility value of ether extract in this study attests to the rabbit ability to utilize dietary
249 fat ²⁶.

250 The high digestibility of NFE represents the readily available carbohydrates. This could be an
251 indication that readily available carbohydrates were well utilized by the rabbits across the
252 treatments.

UNDER PEER REVIEW

253 **Table 6: Economic cost of production of Rabbit Bucks fed Diets containing Mucuna Leaf Meal (MLM)**
 254

Parameters	Experimental Diets					SEM	P-value
	T ₁	T ₂	T ₃	T ₄	T ₅		
Cost of weaner rabbit (₦)	1300	1300	1300	1300	1300	-	-
Cost per Kg	90.95	86.74	85.05	74.98	70.83	-	-
Cost of feed consumed (₦)	473.08 ^a	497.66 ^a	464.74 ^a	400.14 ^b	379.96 ^b	43.76*	.001
Cost per kg gain	379.23	437.17	432.90	431.51	396.77	55.29	.399
Total cost (₦)	1.853 ^a	1.878 ^a	1.845 ^a	1.780 ^b	1.7600 ^b	43.76*	.001
Revenue (₦)	3.438	3.292	3.167	2.918	2.934	311.02	.062
Benefit per Rabbit (₦)	1.585	1.414	1.323	1.138	1.173	282.09	.125
Cost Benefit ratio	1.190	1.344	1.415	1.667	1.580	1.354	.158
% feed cost	25.397 ^a	26.478 ^a	25.180 ^a	22.363 ^b	21.562 ^b	1.766*	.000
% Drug cost	8.108 ^b	7.992 ^b	8.132 ^b	8.427 ^a	8.527 ^a	0.189*	.000
% Cage Cost	1.193 ^c	1.720 ^c	1.192 ^b	1.225 ^{ab}	1.258 ^a	0.031*	.000
% Animal Cost	66.307 ^b	63.921 ^d	65.100 ^b	67.421 ^{ab}	68.200 ^a	1.059*	.000

255

256 ab Means on the same row with different superscript are significantly different (P<0.05), SEM= Standard error of mean, *=significant,

257 T₁ = 0%MLM, T₂ = 5%MLM, T₃ = 10%MLM, T₄ = 15%MLM, T₅ = 20%MLM, %= percentage, ₦361= \$1.

258 Table 6 economic of production of rabbit bucks fed different levels of mucuna leaf
259 meal (MLM)

260 The result of economics of production in this feeding trial showed that cost/kg of the diet was
261 higher in T₁ (₦90.95) and progressively reduced to T₅ (₦70.83) as the level of MLM was
262 increasing in the diets due to the minimal cost of MLM. The cost/kg of the diets, agrees with the
263 findings of ⁷ who observed that the cost/kg feed was reduced generally with increasing dietary
264 yam peel meal. There was no significant difference among the treatments for cost per kg gain,
265 revenue, benefit per rabbit, and cost benefits ratio. However, the cost of feed consumed, total
266 cost of production, percentage feed cost, percentage drug cost, percentage cage cost and
267 percentage animal cost were significantly different (P<0.05). Cost of feed consumed, total cost of
268 production and percentage feed cost were trendy, they were higher at T₁, T₂ and T₃ and declined
269 at T₄ and T₅, meaning the cost reduced as the test ingredient increased to 15 and 20%
270 respectively and these were in agreement with the report of ⁷ who worked on Effects of
271 Replacing Maize with Sun-Dried Yam Peel Meal on Growth Performance Carcass
272 Characteristics and Economics of Production of Meat Type Rabbit. The percentage drug cost,
273 percentage cage cost and percentage animal does not follow any particular trend and as such the
274 variations cannot be attributed to the effect of MLM.

275 The significant differences observed in this study could be attributed to variations in the body
276 weight gain of the animal, disease infestations and different quantities of feed consumed among
277 other things.

278 Base on the findings from this study, the following conclusions have been drawn:

279 This study has revealed that *Mucna pruriens* leaf meal has potentials as a valuable
280 unconventional leguminous feedstuff.

281 The use of *Mucuna pruriens* leaf meal in the diets of rabbit bucks promoted growth best at 5%
282 and thus carcass yield.

283 The economic analysis revealed that with MLM, benefit can be maximized at 20% level of
284 inclusion and also it costs less to produce 1kg live weight of rabbit as level of MLM increased.

285 Based on the results obtained from this study it could be recommended that:

286 Rabbit farmers can incorporate *Mucuna pruriens* leaf meal at 5% in the diets of rabbit bucks for
287 growth performance.

References

- 289 1. Adeyemi, O.A., R. A. Sobayo, F. A. Aluko and D. B. Oke. (2008). Utilisation of rumen
290 filtrate fermented corn-cobs by weaner rabbits. *Nigerian Journal of Animal Production*
291 35(1):69-75.
- 292 2. Aduku, O. A., Dima, N. I., & Aganga, A. A. (1988). A note on comparative evaluation of
293 palm kernel meal, pea-nut meal and sunflower meal in diets of weaned rabbits. *Journal of*
294 *Applied Rabbit Research*, 11, 264-265.
- 295 3. Agunbiade, J. A., Adeyemi, O. A., Fasina, O. E., Ashorobi, B. O., Adebajo, M. O., &
296 Waide, O. A. (1999). Cassava peels and leaves in the diets of rabbits: Effect on
297 performance and carcass characteristics. *Nigerian Journal of Animal Production*, 26, 29-
298 34
- 299 4. Andrea, M. M., and Pablo E. C. (1999). A tropical forage solution to poor quality
300 ruminant diets: A review of *Lablab purpurens*. *Livestock Research for Rural*
301 *Development* (11) 2 1999.
- 302 5. AOAC. (2002). Official of analysis of AOAC International. 17th edition. 1st revision.
303 Gaithersburg, MD, USA, Association of Analytical Communities.
- 304 6. Attah, S., Ortsgera, D. D., & Anugwa, F. O. I. (2011). Effect of replacement of rice offal
305 levels of melon (*Citrullus vulgaris*) seed offal on performance of growing rabbits.
306 *Nigerian Journal of Animal Production*, 38(2), 67-73.
- 307 7. Ayoola, M.A., and Akinbani, A.S. (2011). Effect of replacing maize with sun-dried yam
308 peel meal on growth performance, carcass characteristics and economics of production of
309 meat type rabbit. *Journal of Agricultural Research*. 2(4):70-73
- 310 8. Bawa, G. S., Orunmuyi, M., & Onabanjo, O. A. (2005). Effect of dietary inclusion levels
311 of tresmechanically extracted neem seed cake on performance of young rabbits. *Nigerian*
312 *Journal of Animal Production*, 32(2), 233-239.
- 313 9. De Blas, C. and Wiseman, J. (2003). The nutrition of the rabbit. Cabi Publishing, Oxon,
314 UK.
- 315 10. Duke J.A. New York, NY, USA: Plenum press; 1981. Handbook of legumes of world
316 economic importance.
- 317 11. Emenalom, O. O. and Nwachukwu, I. C. (2006). Effect of calcium hydroxide soaked and
318 cooked velvet beans (*Mucuna pruriens*) on the performance of finisher broilers. *Nigerian*
319 *Journal of Animal Production*, 33(1):53-57
- 320 12. Emiola, I. A., Ologhobo, A. D., Akinlade, J., Adedeji, O. S. and Bamigbade, O. M.
321 (2003). Effects of inclusion of differently processed mucuna seed meal on performance
322 characteristics of broilers. *Tropical Animal Production Investigation* 6:13-21
- 323 13. Ezeokonkwo, M. A. and Okafor, Sunday, N. (2015). Proximate Composition of *Mucuna*
324 *utilis* (Velvet Bean). *IOSR Journal of Applied Chemistry (IOSR-Jac)*. e-ISSN: 2278-
325 5736. Vol. 8, Issue 10 Ver. 1(Oct. 2015), PP 42-45 www.iosrjournals.org
- 326 14. Fathima K. R., P. Tresina Soris, and V.R. Mohan (2010). Nutritional and Antinutritional
327 Assessment of *Mucuna pruriens* (L.) DC. Var. *pruriens*; an Underutilized Tribal Pulse.
328 *Advances in Bio Research*, Vol 1(2) December 2010:79-89.
- 329 15. Fielding, D. (1991). Rabbits. *The Tropical Agriculturalist*. CTA. Macmillan Education
330 Ltd. Macmillan Publishers London, UK, pp: 16–17.
- 331 16. Ifemeje, J. C. (2016). Chemical and Phytochemical Compositions of *Mucuna pruriens*
332 Leaves. *African Journal of Science and Research*, (5) 2:14-17.

- 333 17. Iyayi E. A., Taiwo V. O. (2003). The effect of diet incorporating mucuna (*Mucuna*
334 *Pruriens*) seed meal on the performance of laying hens and broiler. Trop. Subtrop.
335 Agroecosyst., 1: 239-246.
- 336 18. Iyeghe-Erakpotobor, G. T., Tudunwada, I. K., Tijani, A., Muhammad, I. R., & Bello, N.
337 (2012). Influence of level of groundnut haulms and feeding system on performance of
338 growing rabbits. Nigerian Journal of Animal Production, 38(1), 64-73.
- 339 19. Kubkomawa, H, Olaweuye, H. U., Etuk, E. B. and Okoli J. C. (2015). Nutrient
340 Requirements and feed resource availability for pastoral cattle in the tropical Africa; A
341 review. J Agric. and Crop Res. 3(7): 100-116.
- 342 20. McDonald, P., Edward, R. A., Greenlagh, J. F. D. and Morgan, C. A. (1998). Animal
343 Nutrition, 5th Edition. Chapter 10: Evaluation of Foods (A) Digestibility, pp: 221-229.
- 344 21. Mufwa B. J., Kibon A., Mafindi M., Yakubu J. A. (2011). Effect of graded levels of
345 brewers dried grain on the performance of growing rabbits: 1. Growth performance and
346 economic of production. J. Sci. Multidisc. Res. Volume (3) 47-54.
- 347 22. Obun, C.O. and O.A. Adeyemi. (2012). Effects of raw and toasted *Daniellia oliveri* Rolfe
348 seed meal as replacement for groundnut meal on the performance of broiler chickens.
349 Revista Científica UDO Agrícola 12 (4): 947-954.
- 350 23. Ojebiyi, O.O., Oladunjoye, I.O., Aboderin, O.J. and Okelade A.A. (2013). Synergetic
351 Effect of Cassava Sievate/Soybean Milk Residue Mixtures as Replacement for Maize in
352 Growing Rabbits Diet. *Journal of Natural Sciences Research*.3, No.7, 2013. Pp. 126-135.
- 353 24. Oluremi, O. I. A., Mou, P. M. and Adenkola, A. Y. (2008). Effect of fermentation of
354 sweet orange (*Citrus sinensis*) fruit peel on its maize replacement value in broiler diet.
355 Livestock Research for Rural Development. 26(5).
- 356 25. Oluremi, O.I.A., Igyu, A.D. and Abu, F.T. (2005). Response of growing rabbits to dietary
357 replacement of maize with sweet orange (*Citrus sinensis*) rind. *Production Agriculture*
358 *and Technology* 1(1): 130-136.
- 359 26. Onifade, A. A. and Tewe, O. O. (1993), Alternative tropical energy feed resources in
360 rabbit diets: Growth performance, diet digestibility and blood composition. *World*
361 *rabbit Science*. 1:17-24.
- 362 27. Oso, A. O., Bamgbose, A. M., Isah, O. A., Olatunji, J. E. N., Mabadeje, A. T., Alade, A.
363 A and Oni, A. O. (2006). Performance of weaner rabbits fed rice millings waste based
364 diets. *Journal of Animal and Veterinary Advances*, 5(10): 836-838.
- 365 28. Statistical Package for Social Sciences (SPSS, 2007). Version 16.0. Polar Engineering
366 and Consulting <http://www.winwrap.com>
- 367 29. TAC. (2009). Tactical Air Command Meteorological Data. Nigeria Air Force, Makurdi.
- 368 30. Teguia, A. and A.C. Beynen. (2005). Alternative feedstuffs for broilers in Cameroon.
369 Livestock Res Rural Dev. 17(3). Art. 34. <http://www.lrrd.org/lrrd17/3/tegu17034.htm>.
- 370 31. Ujowundu CO, Kalu FN, Emejulu AA, Okafor OE, Nkwonta CG, Nwosunjoku CE
371 (2010). Evaluation of the chemical composition of *Mucuna utilis* leaves used in Herbal
372 Medicine in South-Eastern Nigeria. Afr. J. Pharm. Pharmacol., 4(11): 811-816.

373