

# COMPARATIVE EFFECT OF FUNGICIDES AGAINST BLAST DISEASE OF RICE

---

## ABSTRACT

Keeping in view the importance of rice blast disease, an experiment was conducted in the Laboratory of the Department of Plant Pathology & Seed Science, Sylhet Agricultural University, Sylhet Bangladesh and at the field of regional BADC Seed Production farm, Khadimnagar, Sylhet, Bangladesh, to evaluate seed health status of the collected samples and effectiveness of fungicides against the blast disease of rice. Treatments viz T<sub>1</sub>: Edifen 50 EC (Edifenphos), T<sub>2</sub>: Karisma 28 SC (Azoxystrobin 20% + Cyproconazole 8%), T<sub>3</sub>: Nativo 75 WP (Tebuconazole 50% + Trifloxystrobin 25%), T<sub>4</sub>: Trooper 75WP (Tricyclazole), T<sub>5</sub>: Stanza 75WP (Imidazole), T<sub>6</sub>: Amister top (Azoxystrobin 20% + Difenconazole 12.5%), T<sub>7</sub>: Control (water) were used both in laboratory and field condition. In laboratory, different seed borne fungi, like *Aspergillus*, *Fusarium*, *Curvularia*, *Penicillium*, *Pyricularia*, *Bipolaris*, *Alternaria*, were detected from the collected seed sample by blotter method. In the field, treatments were applied as foliar sprays for three times with ten days interval. The lowest blast disease incidence (34.0%), lowest severity (31.6%) was found in T<sub>2</sub>: Karisma 28 SC treated plots, and gave best result in term of yield (6.3 ton/ha) in comparison to other treatments. The results of the present studies suggested that use of T<sub>2</sub> = Karisma 28 SC is the best choice against rice blast with lowest disease incidence and highest yield.

*Keywords: Rice blast, Pyricularia grisea, seed borne pathogens, fungicidal control*

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is an important crop to provide staple food and food security to millions population of the world and is one of the main foodstuffs in Asia. It is central to Bangladesh's economy, accounting for nearly 20 percent of gross domestic product (GDP) and providing about one-sixth of the national income of Bangladesh [18]. Every year production of rice is affected by different factors of which diseases play a vital role. In Bangladesh, 43 diseases are known to occur on the rice crop, among these diseases 27 are seed borne, of which 14 are of major importance. Of the seed borne diseases of rice, 22 are caused by fungi [3]. Among all the seed borne diseases of rice, blast caused by *Pyricularia grisea* is a major and one of the most devastating diseases. Outbreaks of rice blast are a serious and recurrent problem in all rice growing regions of the world. Rice blast is the most harmful fungal disease in Bangladesh, which can lead to losses in rice yield up to 70-80% [13, 14]. Blast is known to attack nearly all above ground parts as well as during all growth stages of plant. Incidence and severity of blast disease is increasing especially in the Boro season. In recent years, in Bangladesh, frequency of blast occurrence has increased with invasion into new areas (north and northwest parts of the country). The most popular and

33 mega varieties BRRI dhan29 and BRRI dhan28 are recognized to be highly susceptible to  
34 blast disease [1]. For blast disease management at field level chemical control is mainly  
35 practiced, while and other options, particularly water management, are mostly difficult to  
36 practice more problematic [9, 11]. Due to non availability of location specific resistant  
37 varieties for blast disease, the chemical control is the alternate only strategy for the farmers  
38 to harvest obtain economic yield. Moreover, poor bio-efficacy of the biocontrol agents under  
39 the severe epidemic conditions makes the chemical control is an inevitable and ultimate  
40 solution for blast disease management.

41 To combat with against this most devastating and recurrent disease, efforts have been made  
42 to find out the efficacy of various fungicides on the management of rice blast disease and  
43 their impact on grain yield.

44

## 45 2. MATERIAL AND METHODS

46

### 47 2.1 Description of the Study Area

48 The experiment was conducted in the laboratory of dept. Department of Plant Pathology and  
49 Seed Science, Sylhet Agricultural University, Sylhet, Bangladesh, and in the field of regional  
50 BADC farm, Sylhet, Bangladesh during Boro season from December 2016 to May 2017. The  
51 experimental site falls under the Agroecological zone-22 named Northern and Eastern  
52 Piedmont Plains. The climate of the area is subtropical, in the month of December and  
53 January the weather is cold, but from February to June it is characterized by heavy rainfalls,  
54 high temperature and high humidity, while it is and scanty during rest of the year.

### 55 2.2 Experimental Material and Design

56 A very commonly used mega rice variety BRRI dhan28 was used as the experimental unit.  
57 This study was conducted in Randomized Complete Block Design (RCBD) with three  
58 replications. There were 21 unit plots altogether in the field experiment having plot size 2 m<sup>2</sup>.  
59 In the laboratory Completely Randomized Design (CRD) with four replications was followed  
60 where 28 experimental plates were used.

### 61 2.3 Determination of moisture content

62 Moisture content of the seeds of each sample was determined by an electric digital moisture  
63 meter immediately after seed collection.

### 64 2.4 Purity Test

65 Rice seed (40 g) was taken from each original farmer's seed sample for conducting purity  
66 test. Accordingly the seed was grouped into three categories following International Rules for  
67 Testing Seeds (ISTA 2001) as a) pure seed b) other crop seed c) inert matter.

### 68 2.5 Detection of seed borne pathogens by standard blotter method

69 For the experiment, seeds were collected from farmers of different upazilas of Sunamganj  
70 district. The farmer's stored seeds were subjected to blotter incubation test for detection and  
71 identification of seed borne pathogens.

### 72 2.6 Seed treatment with fungicide

73 After blotter incubation test samples having highest *Pyricularia grisea* association were  
74 treated with the **treatment selected** fungicides.

## 75 **2.7 Land preparation and Transplanting**

76 40 days-old seedlings were uprooted from the seedbed and transplanted in the main field.  
77 The selected experimental plot was opened in third week of November 2016. Before  
78 transplanting, harrowing, ploughing, cross ploughing, followed by laddering **were** done to  
79 obtain a good tilth. Two seedlings per hill were transplanted, where hill to hill and row to row  
80 distance was **200 cm x 200 cm**. Transplanting was done **on December 21, 2016**.

## 81 **2.8 Intercultural operations**

82 Different intercultural operations such as weeding, irrigation, fertilization were done as per  
83 requirements.

## 84 **2.9 Procedure of application of treatments in the field**

85 Treatments were applied as foliar spray for 3 times at 10 days intervals in the field.

## 86 **2.10 Assessment of disease incidence**

87 The experiment plots were being monitored after 10 days of interval for the first appearance  
88 of blast disease. The incidence of disease was recorded for three times (35, 45 and 55  
89 DAT). Percent disease incidence was measured **with** the following formula:

$$90 \text{ Disease incidence (\%)} = \frac{\text{Number of infected plant}}{\text{Total number of plant}} \times 100$$

## 91 **2.11 Assessment of disease severity**

92 The observations were recorded and scored at 35, 45 and 55 DAT according to disease  
93 severity score (0-9) from IRRI 1996; [5]. Five infected plants were selected randomly from  
94 each plot.

- 95 ➤ 0 = Leaf free from spot
- 96 ➤ 1 = Small brown specks of pin point size.
- 97 ➤ 2 = Small roundish to slightly elongated, necrotic gray spots, about 1-2 mm in  
98 diameter, with a distinct brown margin, **lesions are mostly (phrase to be completed?)**
- 99 ➤ 3 = Lesion type is the same as in 2, but **a** significant number of lesions **s** are on the  
100 upper leaves.
- 101 ➤ 4 = Typical susceptible blast lesions, 3 mm or longer, infecting less than 4% of the  
102 leaf area.
- 103 ➤ 5 = Typical susceptible blast lesions, 3 mm or longer, infecting **less than** 4-10% of  
104 the leaf area.

- 105 ➤ 6 = Typical susceptible blast lesions, 3 mm or longer, infecting less than 11-25% of  
106 the leaf area.
- 107 ➤ 7 = Typical susceptible blast lesions, 3 mm or longer, infecting less than 26-50% of  
108 the leaf area.
- 109 ➤ 8 = Typical susceptible blast lesions, 3 mm or longer, infecting less than 51-75% of  
110 the leaf area, many leaves dead.
- 111 ➤ 9 = Typical susceptible blast lesions, 3 mm or longer, infecting more than 75% of the  
112 leaf area



113

114

**Plate 1: disease severity scale of rice blast**

115 Disease severity was determined by using following formula [7].

116 Disease severity (%) =  $\frac{\text{Sum of total rating}}{\text{Total no. of observation} \times \text{highest grade in the scale}} \times 100$

## 117 2.12 Harvesting and recording of data

118 The crop was harvested at full ripening stage. The following parameters were recorded from  
119 laboratory and each unit plot and analyzed statistically.

- 120 I. Purity (%)
- 121 II. Germination (%)
- 122 III. Moisture (%)
- 123 IV. Pathogen association with seeds
- 124 V. Disease incidence (%)
- 125 VI. Disease severity (%)
- 126 VII. Yield and yield contributing attributes
- 127

## 128 2.13 Statistical analysis


129 The data obtained for different parameters were statistically analyzed to find out the  
130 significant difference among the treatment. The analysis of variance was performed by using  
131 r program. The difference among the treatment means was estimated by lsd (least  
132 significance difference) test at 5% level of probability [6] (gomez and gomez, 1984).

## 134 3. RESULTS AND DISCUSSIONS

135

### 136 3.1 Determination of moisture content

137 The moisture content of the seed samples varied from 11.88% to 15.09%. The average  
138 moisture content of the seed was 13.23%. Only two samples had moisture content less than  
139 12% but remaining 8 samples had more than 12% moisture content (figure 1).



140

141

Figure 1: Moisture percentage of farmer's stored seed

142 Farmers are not aware enough about the role of accurate moisture percentage on the  
143 storage quality of the seed.

### 144 3.2 Purity analysis

145 In purity analysis, according to ISTA (International Seed Testing Association) rules, seeds  
146 were categorized into three components such as pure seed, other seed and inert matter.  
147 The percentage of pure seeds, other seeds and inert matter ranged 91.25- 95.75%, 2.96 –  
148 7.5% and 1 - 2.36%, and the averages were 93.91%, 4.58% and 1.50%, respectively. Four  
149 samples had more than 95% purity while the rest showed less than 95% (Table 1).  
150  
151

Table 1. Purity percentage of farmers stored seeds


No. of Farmers	Pure seed (%)	Other crop seed (%)	Inert matter (%)
1	95.75	2.96	1.29
2	95.00	3.50	1.5
3	94.20	3.70	2.1
4	91.25	7.50	1.25
5	93.75	4.38	1.87
6	92.13	6.63	1.24
7	95.00	3.72	1.28
8	93.90	5.10	1.00
9	95.45	3.40	1.15
10	92.70	4.94	2.36
Average	93.91	4.58	1.5

152

### 153 3.3 Detection of fungal genera by blotter incubation method

154 After incubation of the sample seeds on blotter paper a total of 7 fungal genera were found  
155 to be associated namely *Aspergillus*, *Fusarium*, *Curvularia*, *Penicillium*, *Pyricularia*, *Bipolaris*,  
156 *Alternaria*. The fungi were detected through germinated conidia observation from sample  
157 seed.






158 **Plate 2: Conidia of the detected fungi under stereo and compound microscope**

159 Seed borne fungal pathogens of rice are detected by many researchers through blotter  
 160 incubation and agar plate method. Ibiom *et al.* (2008) found that *Fusarium moniliforme*,  
 161 *Bipolaris oryzae*, *Fusarium oxysporum*, *Chaetomium globosum*, *Curvularia lunata*,  
 162 *Aspergillus niger*, *Aspergillus flavus*, *Aspergillus terreus*, *Alternaria tenuis* and *Penicillium*  
 163 *spp.* were prevalent in storage [9].

164 **3.4 Effect of Treatments on Pyricularia grisea association with sample seeds**

165 A composite sample was made having highest *Pyricularia grisea* association. ~~Seeds were~~  
 166 ~~treated with the treatments and results observed. From the results it was revealed that~~  
 167 ~~Treatments showed significant effect on the pathogen, with T2 (Karisma 28 SC) providing~~  
 168 ~~the best results resulted minimum association of the pathogen and T7 (Control) resulted~~  
 169 ~~maximum association of the pathogen with the seeds-(Figure 2).~~

170



171  
172

173 **Figure 2: Effect of Treatments on *Pyricularia grisea* association with seed samples**  
 174

175 Manandhar *et al.* (1998) and Hajimo (2001) reported that *P. oryzae* is one of the most  
 176 important fungal pathogen of rice because of its widespread occurrence and destructive  
 177 nature [6, 12]. They also suggested systemic transmission of the fungus from seeds to  
 178 seedlings.

179 **3.5 Evaluation of different treatments on disease incidence of rice blast (leaf**  
 180 **and neck) in field condition**

181 The results of field efficacy of different treatments on disease incidence of rice blast at 35, 45  
 182 and 55 days after transplanting (DAT) were observed and are presented in the table 2. At 35  
 183 DAT, the maximum disease incidence (40.66%) was recorded in T<sub>7</sub> (Control) which was  
 184 statistically similar (38%) to T<sub>5</sub> (Stanza 75 WP) and also similar to T<sub>6</sub> (Amister Top; 37.66%).  
 185 Minimum blast incidence (20.66%) was recorded in T<sub>2</sub> (Karisma 28 SC) followed by T<sub>3</sub>  
 186 (Nativo 75 WP; 26%), then T<sub>4</sub> (Trooper 75WP; 29.33%). At 45 DAT highest incidence was  
 187 recorded in T<sub>7</sub> (Control 56.66%) and lowest incidence was recorded in T<sub>2</sub> (Karisma 28 SC  
 188 25.66%). Both the treatments were statistically different from all other treatments applied. At  
 189 55 DAT disease incidence was minimum in T<sub>2</sub> (Karisma 28 SC 30.33%) followed by T<sub>3</sub>  
 190 (Nativo WP 36%). Maximum incidence was found in T<sub>7</sub> (control 62.33%).

191 The results revealed that in all parameters of incidence status at all times Karisma 28 SC  
 192 could significantly reduce the incidence of the disease.

193  
 194 **Table 2: Effect of different treatments on Rice Blast (leaf and neck) disease incidence**  
 195 **in the field**

Treatments	Disease Incidence (%)		
	35 DAT	45 DAT	55 DAT
T <sub>1</sub> (Edifen 50 EC )	34 b	42.66 c	47.33 cd
T <sub>2</sub> (Karisma 28 SC)	20.66 e	25.66 e	30.33 f
T <sub>3</sub> (Nativo 75 WP)	26 d	36.33 d	36 e
T <sub>4</sub> (Trooper 75WP)	29.33 c	47.33 b	49.33 c
T <sub>5</sub> (Stanza 75 WP)	38 a	46.66 bc	55.66 b
T <sub>6</sub> (Amister top)	37.66 a	43.33 bc	42.66 d
T <sub>7</sub> (Control)	40.66 a	56.66 a	62.33 a
LSD(0.05)	3.12	4.60	5.02
CV (%)	5.44	6.07	6.11

196 Note: Different letter (s) in the same column showed the significant difference at 0.05 level of probability

197  
 198 From reviewed from previous records in the literature it was observed that the effect of  
 199 different fungicides on leaf blast disease under field conditions was significantly less higher  
 200 (15.56%) in tricyclazole sprayed plots followed by kitazin (17.63%) and edifenphos (18.03%)  
 201 [4]. In this experiment we can see that percent disease incidence was comparatively lower in  
 202 T<sub>2</sub> (= Karisma 28 SC; Azoxystrobin 20% + Cyproconazole 8%) than T<sub>4</sub> (= Trooper 75WP;  
 203 Tricyclazole).

204 **3.6 Evaluation of different treatments on disease severity of rice blast (leaf**  
 205 **and neck) in field condition**

206 Disease severity of rice blast at three different days after transplanting (DAT) under different  
 207 treatments was observed. All the treatments resulted significant effect on blast disease



208 control. At 35 DAT, the maximum disease severity (32.66%) was recorded in T<sub>7</sub> (Control)  
 209 which was statistically similar to T<sub>1</sub> (Edifen 50 EC; 31.66%), T<sub>5</sub> (Stanza 75 WP; 33.33%) and  
 210 T<sub>6</sub> (Amister Top; 30.66%). Minimum blast severity (21.66%) was recorded in T<sub>2</sub> (Karisma 28  
 211 SC). At 45 DAT highest severity (40%) was recorded in T<sub>7</sub> (Control) and lowest severity  
 212 (24.33%) was recorded in T<sub>2</sub> (Karisma 28 SC). After T<sub>2</sub> (Karisma 28 SC), T<sub>3</sub> (Nativo 75 WP),  
 213 T<sub>1</sub> (Edifen 50 EC), and T<sub>6</sub> (Amister top) significantly reduced blast severity at 45 DAT. At 55  
 214 DAT, disease severity was minimum (27.66%) in T<sub>2</sub> (Karisma 28 SC). T<sub>3</sub> (Nativo WP;  
 215 30.66%) showed statistically similar result with T<sub>4</sub> (Trooper 75WP; 34%) and T<sub>2</sub> (Karisma 28  
 216 SC; 27.66%), whereas T<sub>2</sub> (Trooper 75WP; 27.66%) and T<sub>4</sub> (Karisma 28 SC; 30.66%) were  
 217 statistically different. Maximum severity was found in T<sub>7</sub> (control; 42.33 %).  
 218

219 **Table 3: Effect of different treatments on Rice Blast (leaf and neck) disease severity in**  
 220 **the field**

Treatments	Disease Severity (%)		
	35 DAT	45 DAT	55 DAT
T <sub>1</sub> (Edifen 50 EC )	31.66 a	31.66 c	36 cd
T <sub>2</sub> (Karisma 28 SC)	21.66 c	24.33 d	27.66 f
T <sub>3</sub> (Nativo 75 WP)	25.66 bc	29.33 c	30.66 ef
T <sub>4</sub> (Trooper 75WP)	29 ab	36 b	34 de
T <sub>5</sub> (Stanza 75 WP)	33.33 a	37.66 ab	38.33
T <sub>6</sub> (Amister top)	30.66 a	30.66 c	bc
T <sub>7</sub> (Control)	32.66 a	40 a	40 ab
			42.33 a
LSD(0.05)	4.33	3.79	3.93
CV (%)	8.33	6.50	6.21

221 Note: Different letter (s) in the same column showed the significant difference at 0.05 level of probability

222

223 Researchers found that application of isoprothiolane and tricyclazole significantly reduced  
 224 the blast severity by 19.5% and 20.06% respectively [2]. Sood and Kapoor (1997) found  
 225 similar result where tricyclazole was the most effective in reducing leaf and neck blast [19].  
 226 Now, In recent years the pathogen *P. grisea* is showing resistance against Trooper 75 WP  
 227 (tricyclazole). In this experiment a new fungicide (Karisma 28 SC; Azoxystrobin 20% +  
 228 Cyproconazole 8%) was used against the disease. The new fungicide which showed the  
 229 best results in controlling the blast disease among all the fungicides used.  
 230

### 231 3.7 Performance of different treatments on yield and yield contributing 232 attributes

233 Along with the all yield other contributing characters, yield was assessed and compared  
 234 within the treatments.

#### 235 3.7.1 Plant Height (cm)

236 The effect of the different treatments on plant height were observed and is presented in the  
 237 (Table 4). From the experiment we found that no any No treatment significantly affected the  
 238 height of the rice plant.

#### 239 3.7.2 Spikelet per panicle

240 Spikelet is the main yield contributing attribute of rice plant. If the number of spikelets is  
 241 higher in each panicle, yield will be increased maximum. In this case T<sub>2</sub> (Karisma 28 SC)  
 242 resulted produced the highest number of spikelets per panicle. It is also statistically similar to  
 243 T<sub>3</sub> (Nativo 75 WP). The lowest number of spikelets was recorded in T<sub>7</sub> (Control) (Table 4).

244 **3.7.3 Unfilled grain per panicle**

245 Rice blast, especially panicle and node blast, causes severe damage to the grain and  
 246 panicle of rice. In severe node blast total panicle breaks down at the base point of the  
 247 panicle. Panicle blast causes unfilled grains, resulting in poor yield. In the present study it  
 248 was found that treatment seven (T<sub>7</sub>) was unsuccessful to control the disease. the number of  
 249 unfilled grains was maximum in T<sub>7</sub> (control), while the lowest number of unfilled grain was  
 250 found in T<sub>2</sub> (Karisma 28 SC). All the other treatments showed significantly better results in  
 251 comparison to than the control plot (Table 4).

252 **3.7.4 Number of effective tiller per hill**

253 Tiller number was not significantly different among the treatments. Not a single treatment  
 254 showed any effect on the difference of number of effective tiller.

255 **3.7.5 Yield**

256 The grain yield was statistically different from one treatment to another among the  
 257 treatments. The minimum and maximum yields were respectively recorded in T<sub>7</sub> (control)  
 258 and maximum was found in T<sub>2</sub> (Karisma 28 SC). Yield differed among the treatments due to  
 259 disease severity, lower number of spikelet per panicle, weather factors (Table 4).  
 260

261 **Table 4: Effect of different treatments on yield and yield contributing attributes**

Treatments	Plant height (cm)	No. of spikelet/panicle	No. of unfilled grain/panicle	No. of effective tiller/hill	Yield/plot (kg/plot)	Yield (ton/ha)
T <sub>1</sub> (Edifen 50 EC)	69.00 a	123 b	20 bc	15.66 ab	2.06 bc	5.15
T <sub>2</sub> (Karisma 28 SC)	72.13 a	134 a	15.33 d	16.33 ab	2.52 a	6.3
T <sub>3</sub> (Nativo 75 WP)	69.83 a	131 a	18 cd	17.33 a	2.18 b	5.45
T <sub>4</sub> (Trooper 75WP)	68.40 a	118.66 bc	21.33 bc	14.66 b	2.02 c	5.05
T <sub>5</sub> (Stanza 75 WP)	73.66 a	114.33 c	19 cd	17.33 a	2.02 c	5.05
T <sub>6</sub> (Amister top)	69.23 a	122 b	23.66 b	16.66 ab	2.05 bc	5.12
T <sub>7</sub> (Control)	73.50 a	105.66 d	34 a	17 ab	1.81 d	4.52
LSD(0.05)	6.78	6.14	4.14	2.66	0.14	
CV (%)	5.38	2.84	10.76	9.10	3.76	

262 Note: Different letter (s) in the same column showed the significant difference at 0.05 level of probability

263  
 264 Prabhu *et al.* (2003) reported that fungicides application increased the yield of rice [15].  
 265 Kumbhar, (2005) found that a maximum increase of 60.99% in grain yield was achieved with  
 266 tricyclazole 75 WP [12]. Similar results were also shown by Prajapati *et al.*, who (2004) and  
 267 they concluded that tricyclazole was significantly superior in decreasing the leaf blast and  
 268 neck blast by 62.9 and 64.1 percent, respectively, with a corresponding increase of 72.3  
 269 percent in grain yield [18]. Here in the In our experiment Karisma 28 SC (combination of  
 270 Azoxystrobin 20% + Cyproconazole 8%), a being newly introduced fungicide, resulted in  
 271 higher grain yield (6.3 ton/ha) in comparison with the very commonly used fungicide Trooper  
 272 75 WP (Tricyclazole) (5.05 ton/ha).

273  
274

#### 4. CONCLUSION

275 Considering the overall findings it was revealed that the seed health status of farmer's stored  
276 boro rice seeds of BRR1 dhan28 is not at satisfactory level. Farmers are therefore may be  
277 advised to collect the seeds from a reliable source, and check the seed health status before  
278 sowing in the main field. Since fungal diseases are most devastating on rice worldwide,  
279 fungicides are important tools to control them. The trial on management of rice blast disease  
280 by the use of several chemical fungicides revealed that Karisma 28 SC (Azoxystrobin 20% +  
281 Cyproconazole 8%) provides the most effective control of leaf and neck blast of rice.

282

#### COMPETING INTERESTS

283 The author has declared no competing interest.

284

285

#### REFERENCES

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

1. Anonymous. Annual research review report for 2010- 2011. Bangladesh Rice Research Institute, Gazipur 1701, Bangladesh; 2011.
2. Arun KS, Sachin U, Ajay S. Field evaluation of insecticides and fungicides for the control of whorl maggot, *Hydrellia philippina* and rice blast caused by *Pyricularia grisea*. *Oryzae* 2011;48 (3) 280-281.
3. Fakir GA. *An annotated list of seed-borne disease in Bangladesh*. Seed Pathology centre, Dept. of Plant Pathology, Bangladesh Agricultural University, Mymensing;2000.
4. Ganesh NR, Gangadhara NB, Basavaraja NT, Krishna NR. Fungicidal management of leaf blast disease in rice *Global Journal of Bio-Science and BioTechnology* 2012; 1 (1) 18-21.
5. Ghazanfar MU, Wakas W, Sahi ST, Saleem Y. Influence of various fungicides on the management of rice blast disease. *Mycopathology* 2009; 7(1) 29-34.
6. Gomez and Gomez. *Statistical procedures for agricultural research*. John Wiley and Sons, New York 1984.
7. Hajimo K. Rice Blast Disease. *Pesticide Outlook* 2001;pp 23-25.
8. Haque MS, Rahman ML, and Malek MA. Effect of fungicides and number of spray on cercospora leaf spot of cowpea. *Bangladesh Journal of Plant Pathology* 1994;10 (1&2) 3-4.
9. Ibiam OFA, Umechuruba CI. A survey of seed-borne fungi associated with seeds of rice (*Oryzae sativa* L. var. 12, 15, and 29) in storage and the field in afikpo north local government area of ebonyi state. *Scientia Africana* 2008; 7 (2) 1-4.
10. IRRI Rice knowledge bank. International Rice Research Institute. Manila, the Philippines 2013. available <<http://www.knowledgebank.irri.org/ipm/riceblast.html>>.
11. ISTA International Rules for Seed Testing. *Seed Science and Technology* 2001; 24 39-42.
12. Kumbhar CT. Evaluation of new fungicide formulations against blast disease of rice. *Karnataka Journal of Agricultural Sciences* 2005;18 (1) 184-185.
13. Lee FN, Singh MP, Counce PA, Gibbons JH. The Mediation Mechanism for flood-induced rice blast field resistance. BR Wells Rice Research Studies. AAES Research Series 2003;pp 517
14. Manandhar HK, Jorgensen HJL, Mathur SB, Petersen VS. Seed borne infection of rice by *Pyricularia oryzae* and its transmission to seedlings. *Plant Disease* 1998; 82(10) 1094-1099.

- 323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339
15. Miah G, Rafii MY, Ismail MR, Puteh AB, Rahim HA, Asfaliza R, Latif MA. Blast resistance in rice: A review of conventional breeding to molecular approaches. *Molecular Biology Reports* 2013; **40** 2369–2388.
  16. Nasruddin A, N Amin. Effects of cultivar, planting period, and fungicide usage on rice blast infection levels and crop yield. *Journal of Agricultural Science* 2013; **5**(1) 160-167.
  17. Prabhu AS, Filippi MC, Zimmermann FJP. Cultivar response to fungicide application in relation to rice blast control, productivity and sustainability. *Pesquisa Agropecuária Brasileira* 2003; **38** 11-17.
  18. Prajapati KS, Patil RC, Pathak AR. Field evaluation of new fungicides against blast of rice. *Pesticides Research Journal* 2004; **16** (2) 26-28.
  19. Sood GK Kapoor AS. Efficacy of new fungicides in the management of rice blast. *Plant Disease Research* 1997; **12** 140-142.
  20. Timothy ST, Khandaker M, Catherine C, Anwarul H, Nazria I, Saad Q, Yan S. Agriculture and Adaptation in Bangladesh: Current and Projected Impacts of Climate Change. IFPRI (International Food Policy Research Institute) 2013; Discussion Paper 01281, pp. 76.

340

UNDER PEER REVIEW