

## Original Research Article

# COMPARATIVE EFFECT OF FUNGICIDES AGAINST BLAST DISEASE OF RICE

### ABSTRACT

Keeping in view the importance of rice blast disease, an experiment was conducted in the Laboratory of dept. of Plant Pathology & Seed Science, Sylhet Agricultural University, Sylhet Bangladesh and at the field of regional BADC Seed Production farm, Khadimnagar, Sylhet, Bangladesh to evaluate seed health status of the collected samples and effectiveness of fungicides against blast disease of rice. Treatments viz T<sub>1</sub>: Edifen 50 EC (Edifenphos), T<sub>2</sub>: Karisma 28 SC (Azoxystrobin 20% + Cyproconazole 8%), T<sub>3</sub>: Nativio 75 WP (Tebuconazole 50% + Trifloxystrobin 25%), T<sub>4</sub>: Trooper 75WP (Tricyclazole), T<sub>5</sub>: Stanza 75WP (Imidazole), T<sub>6</sub>: Amister top (Azoxystrobin 20% + Difenconazole 12.5%), T<sub>7</sub>: Control (water) were used both in laboratory and field condition. In laboratory, different seed borne fungi like *Aspergillus*, *Fusarium*, *Curvularia*, *Penicillium*, *Pyricularia*, *Bipolaris*, *Alternaria*, were detected from the collected seed sample by blotter method. In the field, treatments were applied as foliar spray for three times with ten days interval. The lowest blast disease incidence (34.0%), lowest severity (31.6%) was found in T<sub>2</sub>: Karisma 28 SC treated plots and gave best result in term of yield (6.3 ton/ha) in comparison to other treatments. The results of the present studies suggested that use of T<sub>2</sub> = Karisma 28 SC is the best choice against rice blast with lowest disease incidence and highest yield.

**Keywords:** Rice blast, *Pyricularia grisea*, seed borne pathogens, fungicidal control

### 1. INTRODUCTION

Rice (*Oryza sativa* L.) is an important crop to provide staple food and food security to millions population of the world and is one of the main foodstuffs in Asia. It is central to Bangladesh's economy, accounting for nearly 20 percent of gross domestic product (GDP) and providing about one-sixth of the national income of Bangladesh [18]. Every year production of rice is affected by different factors of which disease play a vital role. In Bangladesh, 43 diseases are known to occur on the rice crop, among these diseases 27 are seed borne of which 14 are of major importance. Of the seed borne diseases of rice, 22 are caused by fungi [3]. Among all the seed borne diseases of rice, blast is a major and one of the most devastating diseases caused by *Pyricularia grisea*. Outbreaks of rice blast are a serious and recurrent problem in all rice growing regions of the world. Rice blast is the most harmful fungal disease in Bangladesh, which can lead to losses in rice yield up to 70 to 80% [13, 14]. Blast is known to attack nearly all above ground parts as well as during all growth stages of plant. Incidence and severity of blast disease is increasing especially in the Boro season. In recent years, in Bangladesh, frequency of blast occurrence has increased with invasion into new areas (north and northwest parts of the country). The most popular and

33 mega varieties BRRI dhan29 and BRRI dhan28 are recognized highly susceptible to blast  
34 disease [1]. For blast disease management at field level chemical control is mainly practiced  
35 and other options particularly water management is mostly difficult to practice [9, 11]. Due to  
36 non availability of location specific resistant varieties for blast disease, the chemical control  
37 is the alternate strategy for the farmers to harvest economic yield. Moreover, poor bio-  
38 efficacy of the bio control agents under the severe epidemic condition makes the chemical  
39 control is an inevitable and ultimate solution for blast disease management.

40 To combat with this most devastating and recurrent disease, efforts have been made to find  
41 out the efficacy of various fungicides on the management of rice blast disease and their  
42 impact on grain yield.

43

## 44 **2. MATERIAL AND METHODS**

45

### 46 **2.1 Description of the Study Area**

47 The experiment was conducted in the laboratory of dept. of Plant Pathology and Seed  
48 Science, Sylhet Agricultural University, Sylhet, Bangladesh and in the field of regional BADC  
49 farm, Sylhet, Bangladesh during Boro season from December 2016 to May 2017. The  
50 experimental site falls under the Agro ecological zone-22 named Northern and Eastern  
51 Piedmont Plains. The climate of the area is subtropical in the month of December and  
52 January the weather is cold but from February to June it is characterized by heavy rainfall,  
53 high temperature and high humidity and scanty during rest of the year.

### 54 **2.2 Experimental Material and Design**

55 A very commonly used mega rice variety BRRI dhan28 was used as the experimental unit.  
56 This study was conducted in Randomized Complete Block Design (RCBD) with three  
57 replications. There were 21 unit plots altogether in the field experiment having plot size 2m<sup>2</sup>.  
58 In the laboratory Completely Randomized Design (CRD) with four replications was followed  
59 where 28 experimental plates were used.

### 60 **2.3 Determination of moisture content**

61 Moisture content of the seeds of each sample was determined by an electric digital moisture  
62 meter immediately after seed collection.

### 63 **2.4 Purity Test**

64 Rice seed (40g) was taken from each original farmer's seed sample for conducting purity  
65 test. Accordingly the seed was grouped into three categories following International Rules for  
66 Testing Seeds (ISTA 2001) as a) pure seed b) other crop seed c) inert matter.

### 67 **2.5 Detection of seed borne pathogens by standard blotter method**

68 For the experiment, seeds were collected from farmers of different upazillas of sunamgonj  
69 district. The farmer's stored seeds were subjected to blotter incubation test for detection and  
70 identification of seed borne pathogens.

### 71 **2.6 Seed treatment with fungicide**

72 After blotter incubation test samples having highest *Pyricularia grisea* association were  
73 treated with the treatment fungicides.

#### 74 **2.7 Land preparation and Transplanting**

75 40 days old seedlings were uprooted from the seedbed and transplanted in the main field.  
76 The selected experimental plot was opened in third week of November 2016. Before  
77 transplanting harrowing, ploughing, cross ploughing, followed by laddering was done to  
78 obtain a good tilth. Two seedlings per hill were transplanted where hill to hill and row to row  
79 distance was 200cmx200cm. Transplanting was done at 21<sup>st</sup> December 2016.

#### 80 **2.8 Intercultural operations**

81 Different intercultural operations such as weeding, irrigation, fertilization were done as per  
82 requirements.

#### 83 **2.9 Procedure of application of treatments in the field**

84 Treatments were applied as foliar spray for 3 times at 10 days intervals in the field.

#### 85 **2.10 Assessment of disease incidence**

86 The experiment plots were being monitored after 10 days of interval for the first appearance  
87 of blast disease. The incidence of disease was recorded for three times (35, 45 and 55  
88 DAT). Percent disease incidence was measured by the following formula:

$$89 \text{ Disease incidence (\%)} = \frac{\text{Number of infected plant}}{\text{Total number of plant}} \times 100$$

#### 90 **2.11 Assessment of disease severity**

91 The observations were recorded and scored at 35, 45 and 55 DAT according to disease  
92 severity score (0-9) from IRRRI 1996; [5]. Five infected plants were selected randomly from  
93 each plot.

- 94 > 0 = Leaf free from spot
- 95 > 1 = Small brown specks of pin point size.
- 96 > 2 = Small roundish to slightly elongated, necrotic gray spots, about 1-2 mm in  
97 diameter, with a distinct brown margin, lesions are mostly
- 98 > 3 = Lesion type is the same as in 2, but significant number of lesion are on the upper  
99 leaves.
- 100 > 4 = Typical susceptible blast lesions, 3 mm or longer, infecting less than 4% of the  
101 leaf area.
- 102 > 5 = Typical susceptible blast lesions, 3 mm or longer, infecting less than 4-10% of  
103 the leaf area.

- 104 > 6 = Typical susceptible blast lesions, 3 mm or longer, infecting less than 11-25% of  
105 the leaf area.
- 106 > 7 = Typical susceptible blast lesions, 3 mm or longer, infecting less than 26-50% of  
107 the leaf area.
- 108 > 8 = Typical susceptible blast lesions, 3 mm or longer, infecting less than 51-75% of  
109 the leaf area, many leaves dead.
- 110 > 9 = Typical susceptible blast lesions, 3 mm or longer, infecting more than 75% of the  
111 leaf area



112

113

**Plate 1: disease severity scale of rice blast**

114 Disease severity was determined by using following formula [7].  
 115 
$$\text{Disease severity (\%)} = \frac{\text{Sum of total rating}}{\text{Total no.of observation} \times \text{highest grade in the scale}} \times 100$$

116 **2.12 Harvesting and recording of data**

117 The crop was harvested at full ripening stage. Following parameters were recorded from  
 118 laboratory and each unit plot and analyzed statistically.

- 119 I. Purity (%)
- 120 II. Germination (%)
- 121 III. Moisture (%)
- 122 IV. Pathogen association with seeds
- 123 V. Disease incidence (%)
- 124 VI. Disease severity (%)
- 125 VII. Yield and yield contributing attributes
- 126


127 **2.13 Statistical analysis**

128 The data obtained for different parameters were statistically analyzed to find out the  
 129 significant difference among the treatment. The analysis of variance was performed by using  
 130 r program. The difference among the treatment means was estimated by lsd (least  
 131 significance difference) test at 5% level of probability (gomez and gomez, 1984).  
 132

133 **3. RESULTS AND DISCUSSIONS**

134 **3.1 Determination of moisture content**

135 The moisture content of the seed samples varied from 11.88% to 15.09%. The average  
 136 moisture content of the seed was 13.23 %. Only two samples had moisture content less than  
 137 12% but remaining 8 samples had more than 12% moisture content (figure 1).



138 **Figure 1: Moisture percentage of farmer's stored seed**

140 Farmers are not aware enough about the role of accurate moisture percentage on the  
141 storage quality of the seed.

### 142 3.2 Purity analysis

143 In purity analysis, according to ISTA (International Seed Testing Association) rules, seeds  
144 were categorized into three components such as pure seed, other seed and inert matter.  
145 The percentage of pure seeds, other seeds and inert matter ranged 91.25- 95.75%, 2.96 –  
146 7.5% and 1 - 2.36%, and the averages were 93.91%, 4.58% and 1.50%, respectively. Four  
147 samples had more than 95% purity while the rest showed less than 95% (Table 2).

148 **Table 1. Purity percentage of farmers stored seeds**

No. of Farmers	Pure seed (%)	Other crop seed (%)	Inert matter (%)
1	95.75	2.96	1.29
2	95.00	3.50	1.5
3	94.20	3.70	2.1
4	91.25	7.50	1.25
5	93.75	4.38	1.87
6	92.13	6.63	1.24
7	95.00	3.72	1.28
8	93.90	5.10	1.00
9	95.45	3.40	1.15
10	92.70	4.94	2.36
Average	93.91	4.58	1.5

149

### 150 3.3 Detection of fungal genera by blotter incubation method


151 After incubation of the sample seeds on blotter paper a total of 7 fungal genera were found  
152 to be associated namely *Aspergillus*, *Fusarium*, *Curvularia*, *Penicillium*, *Pyricularia*, *Bipolaris*,  
153 *Alternaria*. The fungi were detected through germinated conidia observation from sample  
154 seed.



*Alternaria sp.*

*Fusarium sp.*

*Aspergillus sp.*



155 **Plate 2: Conidia of the detected fungi under stereo and compound microscope**


156 Seed borne fungal pathogens of rice are detected by many researchers through blotter  
 157 incubation and agar plate method. Ibiam et al. (2008) found that *Fusarium moniliforme*,  
 158 *Bipolaris oryzae*, *Fusarium oxysporum*, *Chaetomium globosum*, *Curvularia lunata*,  
 159 *Aspergillus niger*, *Aspergillus flavus*, *Aspergillus terreus*, *Alternaria tenuis* and *Penicillium sp.*  
 160 were prevalent in storage [8].

**Comment [m1]:** italics

161 **3.4 Effect of Treatments on *Pyricularia grisea* association with sample seeds**

**Comment [m2]:** no need numbers

162 A composite sample was made having highest *Pyricularia grisea* association. Seeds were  
 163 treated with the treatments and results observed. From the results it was revealed that  
 164 treatments showed significant effect on the pathogen, T2 (Karisma 28 SC) resulted minimum  
 165 association of the pathogen and T7 (Control) resulted maximum association of the pathogen  
 166 with the seeds (Figure 2).



**Comment [m3]:** remove the percentage symbol insidevthe graph

167  
 168 **Figure 2: Effect of Treatments on *Pyricularia grisea* association with seed samples**  
 169

170 Manandhar *et al.* (1998) and Hajimo (2001) reported that *P. oryzae* is one of the most  
 171 important fungal pathogen of rice because of its widespread occurrence and destructive  
 172 nature[6, 12]. They also suggested systemic transmission of the fungus from seeds to  
 173 seedlings.

174 **3.5 Evaluation of different treatments on disease incidence of rice blast (leaf  
 175 and neck) in field condition**

176 The results of field efficacy of different treatments on disease incidence of rice blast at 35, 45  
 177 and 55 days after transplanting (DAT) were observed and presented in the table 1. At 35  
 178 DAT, the maximum disease incidence (40.66%) was recorded in T<sub>7</sub> (Control) which was  
 179 statistically similar (38%) with T<sub>5</sub> (Stanza 75 WP) and also similar to T<sub>6</sub> (Amister Top;  
 180 37.66%). Minimum blast incidence (20.66%) was recorded at T<sub>2</sub> (Karisma 28 SC) followed  
 181 by T<sub>3</sub> (Nativo 75 WP; 26%) then T<sub>4</sub> (Trooper 75WP; 29.33%). At 45 DAT highest incidence  
 182 was recorded in T<sub>7</sub> (Control 56.66%) and lowest incidence was recorded in T<sub>2</sub> (Karisma 28  
 183 SC 25.66%). Both the treatments were statistically different to all other treatments applied.  
 184 At 55 DAT disease incidence was minimum in T<sub>2</sub> (Karisma 28 SC 30.33%) followed by T<sub>3</sub>  
 185 (Nativo WP 36%). Maximum incidence was found in T<sub>7</sub> (control 62.33%).

186 The results revealed that in all parameters of incidence status Karisma 28 SC could  
 187 significantly reduce the incidence of the disease.

188 **Table 2: Effect of different treatments on Rice Blast (leaf and neck) disease incidence  
 189 in the field**

Treatments	Disease Incidence (%)		
	35 DAT	45 DAT	55 DAT
T <sub>1</sub> (Edifen 50 EC )	34 b	42.66 c	47.33 cd
T <sub>2</sub> (Karisma 28 SC)	20.66 e	25.66 e	30.33 f
T <sub>3</sub> (Nativo 75 WP)	26 d	36.33 d	36 e
T <sub>4</sub> (Trooper 75WP)	29.33 c	47.33 b	49.33 c
T <sub>5</sub> (Stanza 75 WP)	38 a	46.66 bc	55.66 b
T <sub>6</sub> (Amister top)	37.66 a	43.33 bc	42.66 d
T <sub>7</sub> (Control)	40.66 a	56.66 a	62.33 a
LSD(0.05)	3.12	4.60	5.02
CV (%)	5.44	6.07	6.11

190 Note: Different letter (s) in the same column showed the significant difference at 0.05 level of probability

191  
 192 From the literature reviewed from previous records it was observed that the effect of different  
 193 fungicides on leaf blast disease under field conditions was significantly less (15.56%) in  
 194 tricyclazole sprayed plots followed by kitazin (17.63%) and edifenphos (18.03%) [4]. In this  
 195 experiment we can see that percent disease incidence was comparatively lower in T<sub>2</sub>  
 196 =Karisma 28 SC (Azoxytrobino 20% + Cyproconazole 8%) than T<sub>4</sub>= Trooper 75WP  
 197 (Tricyclazole).

198 **3.6 Evaluation of different treatments on disease severity of rice blast (leaf  
 199 and neck) in field condition**

200 Disease severity of rice blast at three different days after transplanting (DAT) under different  
 201 treatments was observed. All the treatments resulted significant effect on blast disease  
 202 control. At 35 DAT, the maximum disease severity (32.66%) was recorded in T<sub>7</sub> (Control)  
 203 which was statistically similar to T<sub>1</sub> (Edifen 50 EC 31.66%), T<sub>5</sub> (Stanza 75 WP 33.33%) and  
 204 T<sub>6</sub> (Amister Top 30.66%). Minimum blast severity (21.66%) was recorded at T<sub>2</sub> (Karisma 28  
 205 SC). At 45 DAT highest severity (40%) was recorded in T<sub>7</sub> (Control) and lowest severity  
 206 (24.33%) was recorded in T<sub>2</sub> (Karisma 28 SC). After T<sub>2</sub> (Karisma 28 SC), T<sub>3</sub> (Nativo 75 WP),



207 T<sub>1</sub> (Edifen 50 EC), and T<sub>6</sub> (Amister top) significantly reduced blast severity at 45DAT. At 55  
 208 DAT, disease severity was minimum (27.66%) in T<sub>2</sub> (Karisma 28 SC). T<sub>3</sub> (Nativo WP  
 209 30.66%) showed statistically similar result with T<sub>4</sub> (Trooper 75WP 34%) and T<sub>2</sub> (Karisma 28  
 210 SC 27.66%) whereas T<sub>2</sub> (Trooper 75WP 27.66%) and T<sub>4</sub> (Karisma 28 SC 30.66%) are  
 211 statistically different. Maximum severity was found in T<sub>7</sub> (control 42.33 %).

212 **Table 3: Effect of different treatments on Rice Blast (leaf and neck) disease severity in**  
 213 **the field**

Treatments	Disease Severity (%)		
	35 DAT	45 DAT	55 DAT
T <sub>1</sub> (Edifen 50 EC )	31.66 a	31.66 c	36 cd
T <sub>2</sub> (Karisma 28 SC)	21.66 c	24.33 d	27.66 f
T <sub>3</sub> (Nativo 75 WP)	25.66 bc	29.33 c	30.66 ef
T <sub>4</sub> (Trooper 75WP)	29 ab	36 b	34 de
T <sub>5</sub> (Stanza 75 WP)	33.33 a	37.66 ab	38.33
T <sub>6</sub> (Amister top)	30.66 a	30.66 c	bc
T <sub>7</sub> (Control)	32.66 a	40 a	40 ab
LSD(0.05)	4.33	3.79	3.93
CV (%)	8.33	6.50	6.21

Note: Different letter (s) in the same column showed the significant difference at 0.05 level of probability

214

215

216

217

218

219

220

221

222

223

224

### 3.7 Performance of different treatments on yield and yield contributing attributes

225

226

227

228

#### 3.7.1 Plant Height (cm)

229

230

231

232

#### 3.7.2 Spikelet per panicle

233

234

235

236

237

#### 3.7.3 Unfilled grain per panicle

238

239

240

241

242

243

244

245

246

Rice blast specially panicle and node blast causes severe damage to the grain and panicle of rice. In severe node blast total panicle breaks down at the base point of the panicle. Panicle blast causes unfilled grain resulting poor yield. In the present study it is found that treatment seven (T<sub>7</sub>) was unsuccessful to control the disease. Number of unfilled grain was maximum in T<sub>7</sub> (control). The lowest no. of unfilled grain was found in T<sub>2</sub> (Karisma 28 SC). All the other treatments showed significantly better result in comparison to control plot (Table 7).

247 **3.7.4 No. of effective tiller per hill**

248 Tiller number was not significantly differed among the treatments. Not a single treatment  
249 showed any effect on the difference of number of effective tiller.

250 **3.7.5 Yield**

251 The grain yield was statistically different from one treatment to another treatment. The  
252 minimum yield was recorded in T<sub>7</sub> (control) and maximum was found in T<sub>2</sub> (Karisma 28 SC).  
253 Yield differed among the treatments due to disease severity, lower number of spikelet per  
254 panicle, weather factors (table 7).  
255

256 **Table 4: Effect of different treatments on yield and yield contributing attributes**

Treatments	Plant height (cm)	No. of spikelet/panicle	No. of unfilled grain/panicle	No. of effective tiller/hill	Yield/plot (kg/plot)	Yield (ton/ha)
T <sub>1</sub> (Edifen 50 EC)	69.00 a	123 b	20 bc	15.66 ab	2.06 bc	5.15
T <sub>2</sub> (Karisma 28 SC)	72.13 a	134 a	15.33 d	16.33 ab	2.52 a	6.3
T <sub>3</sub> (Nativo 75 WP)	69.83 a	131 a	18 cd	17.33 a	2.18 b	5.45
T <sub>4</sub> (Trooper 75WP)	68.40 a	118.66 bc	21.33 bc	14.66 b	2.02 c	5.05
T <sub>5</sub> (Stanza 75 WP)	73.66 a	114.33 c	19 cd	17.33 a	2.02 c	5.05
T <sub>6</sub> (Amister top)	69.23 a	122 b	23.66 b	16.66 ab	2.05 bc	5.12
T <sub>7</sub> (Control)	73.50 a	105.66 d	34 a	17 ab	1.81 d	4.52
LSD(0.05)	6.78	6.14	4.14	2.66	0.14	
CV (%)	5.38	2.84	10.76	9.10	3.76	

257 Note: Different letter (s) in the same column showed the significant difference at 0.05 level of probability

257

258

259 Prabhu *et al.* (2003) reported that fungicides application increased the yield of rice [15].  
260 Kumbhar, (2005) found a maximum increase of 60.99% in grain yield was achieved with  
261 tricyclazole 75 WP [10]. Similar results also shown by Prajapati *et al.* (2004) and they  
262 concluded that tricyclazole was significantly superior in decreasing the leaf blast and neck  
263 blast by 62.9 and 64.1 percent, respectively with corresponding increase of 72.3 percent in  
264 grain yield [16]. Here in the experiment Karisma 28 SC (combination of Azoxystrobin 20% +  
265 Cyproconazole 8%) being newly introduced fungicide resulted higher grain yield (6.3 ton/ha)  
266 in comparison to very commonly used fungicide Trooper 75 WP (Tricyclazole) (5.05ton/ha).  
267

267

268

268 **4. CONCLUSION**

269

270

271

272

273

274

275

276

277

278

278 **COMPETING INTERESTS**

279

279 The author has declared no competing interest.

280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
  
324  
325  
326  
327  
328  
329  
330

## REFERENCES

1. Anonymous. Annual research review report for 2010- 2011. Bangladesh Rice Research Institute, Gazipur 1701, Bangladesh; 2011.
2. Arun KS, Sachin U, Ajay S. Field evaluation of insecticides and fungicides for the control of whorl maggot, *Hydrellia philippina* and rice blast caused by *Pyricularia grisea*. *Oryzae* 2011; **48** (3) 280-281.
3. Fakir GA. *An annotated list of seed-borne disease in Bangladesh*. Seed Pathology centre, Dept. of Plant Pathology, Bangladesh Agricultural University, Mymensing;2000.
4. Ganesh NR, Gangadhara NB, Basavaraja NT, Krishna NR. Fungicidal management of leaf blast disease in rice *Global Journal of Bio-Science and BioTechnology* 2012; **1** (1) 18-21.
5. Ghazanfar MU, Wakas W, Sahi ST, Saleem Y. Influence of various fungicides on the management of rice blast disease. *Mycopathology* 2009; **7**(1) 29-34.
6. Gomez and Gomez. *Statistical procedures for agricultural research*. John Wiley and Sons, New York 1984.
7. Hajimo K. Rice Blast Disease. *Pesticide Outlook* 2001;pp 23-25.
8. Haque MS, Rahman ML, and Malek MA. Effect of fungicides and number of spray on cercospora leaf spot of cowpea. *Bangladesh Journal of Plant Pathology* 1994; **10** (1&2) 3-4.
9. Ibiam OFA, Umechuruba CI. A survey of seed-borne fungi associated with seeds of rice (*Oryzae sativa* L) faro12, 15, and 29) in storage and the field in afikpo north local government area of ebonyi state. *Scientia Africana*2008; **7** (2) 1-4.
10. IRRI Rice knowledge bank. International Rice Research Institute. Manila, the Philippines 2013. available <<http://www.knowledgebank.irri.org/ipm/riceblast.html>>.
11. ISTA International Rules for Seed Testing. *Seed Science and Technology*2001; **24** 39-42.
12. Kumbhar CT. Evaluation of new fungicide formulations against blast disease of rice. *Karnataka Journal of Agricultural Sciences* 2005; **18** (1) 184-185.
13. Lee FN, Singh MP, Counce PA, Gibbons JH. The Mediation Mechanism for flood-induced rice blast field resistance. BR Wells Rice Research Studies. AAES Research Series 2003;pp 517
14. Manandhar HK, Jorgensen HJL, Mathur SB, Petersen VS. Seed borne infection of rice by *Pyricularia oryzae* and its transmission to seedlings. *Plant Disease* 1998; **82**(10) 1094-1099.
15. Miah G, Rafii MY, Ismail MR, Puteh AB, Rahim HA, Asfaliza R, Latif MA. Blast resistance in rice: A review of conventional breeding to molecular approaches. *Molecular Biology Reports* 2013; **40** 2369–2388.
16. Nasruddin A, N Amin. Effects of cultivar, planting period, and fungicide usage on rice blast infection levels and crop yield. *Journal of Agricultural Science* 2013; **5**(1) 160-167.
17. Prabhu AS, Filippi MC, Zimmermann FJP. Cultivar response to fungicide application in relation to rice blast control, productivity and sustainability. *Pesquisa Agropecuária Brasileira* 2003; **38** 11-17.
18. Prajapati KS, Patil RC, Pathak AR. Field evaluation of new fungicides against blast of rice. *Pesticides Research Journal* 2004; **16** (2) 26-28.
19. Sood GK Kapoor AS. Efficacy of new fungicides in the management of rice blast. *Plant Disease Research* 1997; **12** 140-142.

331  
332  
333  
334  
335

20. Timothy ST, Khandaker M, Catherine C, Anwarul H, Nazria I, Saad Q, Yan S. Agriculture and Adaptation in Bangladesh: Current and Projected Impacts of Climate Change. IFPRI (International Food Policy Research Institute) 2013; Discussion Paper 01281, pp. 76.

336

UNDER PEER REVIEW