

2 **EFFECT OF POWDERED AND COMPOSTED MEAT BONES ON THE**
3 **GROWTH AND YIELD OF WATER SPINACH (*Ipomoea aquatica*)**

4
5
6 • **Abstract:**

7 For agriculture, the most important is the soil's function as a base and medium for plant
8 growth. Soil health and soil condition is important, and the management of soil organic
9 matter is an important part of managing soil health and maintaining soil condition. The
10 experiment was carried out in the field lab (Net House) of Soil, Water and Environment
11 discipline, Khulna University, Bangladesh. Total 21 plastic pots were used to continue the
12 experiment with 7 treatments. For the experiment, the meat bones were collected and used in
13 the experimental pot along with soil in different doses. This research has revealed that meat
14 bones both powdered and composted, has significantly increased the growth parameters of
15 water spinach (*Ipomoea aquatica*) such as a number of leaves, shoot length, fresh weight, dry
16 weight and moisture content.

17 **Keywords:** Soil Organic Matter, Meat bones, Water spinach (*Ipomoea aquatica*), Pot
18 experiment

19 **1. Introduction**

20 Agriculture is the most important sector of the economy of Bangladesh, contributing about
21 23% of the country's GDP and employing about 62% of the total labour force. Bangladesh,
22 after independence, has adopted various measures to raise the vegetable production for
23 tackling the malnutrition problem in the country. Nevertheless, vegetable production has
24 marginally increased from 7.4 lakh tons in 1974/75 to 9.2 lakh tons in 1987/88 with year to
25 year fluctuations (GOB 1985 to GOB 1989).

26 Water spinach (*Ipomoea aquatica*) also known as 'kalmi shak' (in Bengali) is one of the
27 popular vegetables that is promoted to grow in Bangladesh due to its good nutritive value,
28 antioxidant properties, high fiber content and many other health health-related benefits
29 (Hongfei, 2011; Kala and Prakash, 2004; Faruq *et al.*, 2002; and Ogle *et al.*, 2001). It is a
30 vascular semi-aquatic herbaceous perennial plant belonging to *Convolvulaceae* (USDA,
31 2005). It has a hollow and viny stem, grows prostrate or floating, and roots are coming from
32 the nodes that penetrate the soil. Water spinach is native to the tropics and subtropics of
33 Southeast Asia, Southern China and India (Gothberg, *et al.*, 2005 and Chen *et al.*, 1991). It is
34 a green vegetable and is ranked high among the world's healthiest foods, and there are plenty
35 of reasons for it being so. It is one of the ideal options to manage weight and to lower the
36 cholesterol levels. It consists of rich amounts of iron that is required by the red blood cells

37 during process of hemoglobin formation. So, people who are suffering with anemia should
38 include their diet with this iron rich leafy vegetable (Gupta *et al.*, 2005).



39

40

Figure 1: Water spinach

41 For agriculture, the most important is the soil's function as a base and medium for plant
42 growth. Soil health and soil condition is important, and the management of soil organic
43 matter is an important part of managing soil health and maintaining soil condition. Most of
44 the soils of Bangladesh have low organic matter content, usually less than 2% (Bhuiyan,
45 1994). A good soil should have at least 2.5% organic matter, but in Bangladesh, most soils
46 have less than 1.5%, and some soils have even less than 1% organic matter (BARC, 2005).
47 The pressure on the management of soil organic matter is increasing as costs of inputs for
48 agriculture increase and the capacity and ability to overcome soils in poor condition by
49 adding more fertilizer, adding one more cultivation, adding one more irrigation or adding
50 another input are diminished (MacEwan, 2007). The ultimate source of organic matter for
51 most soils is through the fixation of carbon dioxide from the atmosphere through
52 photosynthetic reactions by plants. There is also a very small input from autotrophic bacteria.
53 However, in some instances, there may also be some input from industrial and mining
54 products derived from petroleum or coal. At the broad scale, these sources of soil carbon are
55 insignificant. Soil organic matter is derived from organic materials that are added to the soil,
56 and the majority of soil organic matter derives from the breakdown of residues remaining
57 after plants have died. These residues can take the form of root residues located in the soil
58 matrix or leaves, stems and stubble existing as litter on the soil surface. Animals also provide
59 a proportion of the soil organic matter to varying degrees depending on management and the
60 ecosystem (Tate, 1987).

61 Additions of large amounts of organic materials as composts or as biochar can increase the
62 levels of soil organic matter in soils (Gibson *et al.*, 2002). Soils are used as a filter or sink for
63 effluent and waste materials. The wastes produced from meat bones can be used as such types
64 of organic materials as the replacement of compost or fertilizers.

65 Maximum people eat various types of meat daily. There are some particular occasions such
66 as wedding ceremony, some religious festivals like Eid, Puja etc which is incomplete without
67 meat. A lot of meat bones can be found after these ceremonies. If we use these bones as

68 fertilizers for plant growth, then the waste load will be controlled. Usually, beef bones,
69 however, other types of a creature like chicken or pigeon bone meal or emulsion can be used
70 to prepare meat bone fertilizer. The processed bone meal goes through to become a powder or
71 compost will quickly kill off many pathogens.

72 Meat and bone meal (MBM) can be a viable alternative to mineral fertilizers because it
73 contains about 8% Nitrogen (N), 5 % Phosphorus (P), 1 % Potassium (K) and 10 % Calcium
74 (Ca) (Ylivainio *et al.*, 2007), which makes it a valuable source of nutrients for plant
75 production. It has about 50% protein, 35% ash, 8-12% fat, and 4-7% moisture, and contains a
76 big amount of nutrients. Chemical properties of Meat Bone Materials (MBM) vary a lot from
77 different raw materials. On average, the pH tends to be acidic, about 6.5. Organic matter in
78 content is about 50% (Jeng *et al.*, 2006). Besides a great deal of phosphorous, bone meal also
79 provides smaller amounts of essential nitrogen, potassium and calcium. As the bone meal
80 breaks down over the season, it will steadily and slowly release these nutrients for plant
81 uptake. The advantageous aspects of slow release nutrition are that plant roots will not burn,
82 and that soil vitality is increased for extended periods (Cayuela *et al.*, 2009).

83

84 **1.1. Objectives**

- 85 ▪ To minimize the rate of waste load by using the meat bones as organic fertilizer which
86 is environmental friendly as well as to improve soil health
- 87 ▪ For determining the effectiveness of different types of meat bones in water spinach
88 cultivation
- 89 ▪ To assess the effect of composted and powdered meat bones on the growth and yield
90 of water spinach (*Ipomoea aquatica*)

91

92 **2. Methods and Materials**

93 **2.1. Location of the study area**

94 The experiment was carried out in the field lab (Net House) of Soil, Water and Environment
95 discipline, Khulna University, Bangladesh.

96 **2.2. Collection and preparation of the soil**

97 The soil for our research was collected from the research field of Soil, Water and
98 Environment Discipline, Khulna University. Spade, plastic bag were used for collecting
99 the soil. Then the soil was air dried at room temperature, and all the plant debris were
100 removed manually. Then the massive aggregates were broken by gentle crushing by a
101 hammer. Then the soil was sieved by 2mm sieve.

102 **2.3. Preparation of pot**

103 After sieving, the soil was taken into pots for sowing the seeds. Total 21 plastic pots were
104 used to continue the experiment. The pots were cleaned and labeled. Each Pot was filled

105 with 3kg sieved soil. Pots had no pore in its bottom to protect the leaching of meat bone
106 fertilizers from the soil. The experiment was started on 24th June, 2018.

107 **2.4. Powdered and composted meat bones preparation**

108 The meat bones were collected after eating. Then some of the bones were crushed into
109 powder form with the help of crusher (Haman dista). To prepare meat bone compost, bones
110 (pigeon, chicken, and beef) were crushed and transferred it into a plastic bag. Sufficient
111 amount of water was added into it to degrade the meat bones and kept it by digging soil into
112 30cm depth. After 60 days (4th June – 4th August), the compost was ready to be applied into
113 my experiment. 10g powdered and composted meat bones were applied into each pot. 5g was
114 mixed up with the soil before sowing the seeds, and the rest 5g was added when the seeds
115 were grown up to plants.

116

117 **2.5. Treatments of investigation**

118 Seven treatments were used in the experiment.

119 T₀ = Control

120 T₁ = 500 kg ha⁻¹ powdered beef bone

121 T₂ = 500 kg ha⁻¹ powdered chicken bone

122 T₃ = 500 kg ha⁻¹ powdered pigeon bone

123 T₄ = 500 kg ha⁻¹ composted beef bone

124 T₅ = 500 kg ha⁻¹ composted chicken bone

125 T₆ = 500 kg ha⁻¹ composted pigeon bone

126

127 **2.6. Sowing of the seeds**

128 The seeds were sown on 24th June, 2018. The seeds were sown thoroughly as it was possible
129 to keep uniformity and then the seeds were covered by soils. 0.01 g seeds (5 kg ha⁻¹ as
130 recommended by BARI, 2005 for trial experiment) were sown in each pot and maximum
131 seeds germinated within 5 days. After germination only five plants were kept in each pot.



Figure 2: Sowing the seeds of water spinach

2.7. Intercultural operations

2.7.1. Watering

For normal plant growth, sufficient amount of water was added regularly to each pot for maintaining optimum moisture.

2.7.2 General observation

The pots under experiment were frequently observed to note any change in the crop growth and other characteristics. The crop growth was very satisfactory in some treatments. But some treatments showed a lower number of plants.

2.7.3 Harvesting

The experimental crops were harvested after 40 days of germination. The harvested plants were tagged separately, weighted, oven dried at 65⁰C temperature for 24 hours until moisture content reached to a minimum condition. The dried material of plants per pot from each treatment was collected.

2.8. Morphological attributes of water spinach

2.8.1. Number of leaves

The number of leaves of five plants of each pot was counted and average value was taken.

2.8.2. Shoot length (cm)

Shoot length was measured using a measuring scale from the root level to the tip of the plant. From each pot, five plants were measured and averaged.

2.8.3. Root length (cm)

Root length was measured using a measuring scale from root level to the tip of the longest root at harvest, and their average value was taken as the root length in cm.

157 **2.8.4. Fresh weight per plant (gm)**

158 Harvest of five plants from each pot, fresh weight of the whole plant was taken by an
159 electrical balance_ and their mean value was calculated as fresh weight expressed in gm/plant.

160 **2.8.5. Dry weight per plant (gm)**

161 Five plants from each pot were collected and oven dried at 65⁰C for 48 hours, weighed in
162 gm/plant by an electrical balance and average value was recorded.

163

164 **2.8.6. Moisture content (%)**

165 Percent moisture was calculated by using the formula:

166 Moisture content (%) = $\frac{W_f - W_o}{W_f} \times 100$

167 Where,

168 W_f = Fresh weight of the plant sample

169 W_o = Oven dry weight of the plant sample

170

171 **2.9. Statistical analysis**

172 The collected data on different parameters were represented in bar diagram by Microsoft
173 office excel program 16.0 and statistically analyzed following the analysis of variance
174 (ANOVA) technique. Statistical analysis was performed by using MINITAB 18 statistical
175 package.

176

177 **3. Result and Discussion**

178 Effect of powdered and composted meat bone on the growth and yield of Water spinach
179 (*Ipomoea aquatica*) was studied following pot experiment. The number of leaves per plant,
180 shoot length per plant, root length per pant, fresh weight per plant, dry weight per plant and
181 percent moisture content were measured for the plants treated with 10g powdered and
182 composted meat bone and compared with plants grown with no meat bone which was control
183 experiment. The data of morphological attributes are presented in Table 1.

184


185 **Table 1:** Observed growth factors of the experimented water spinach for different treatments

Treatments	No. of leaves /plant	Root length /plant (cm)	Shoot length /plant (cm)	Fresh weight /plant (g)	Dry weight /plant (g)	Moisture content (%)
T ₀	8	8.81	17.64	1.09	0.08	64.36
T ₁	9	9.31	24.32	1.56	0.37	76.49
T ₂	9	9.98	24.25	1.23	0.32	73.85
T ₃	8	10.93	18.65	1.17	0.35	70.25
T ₄	9	12.65	22.01	2.67	0.40	84.97
T ₅	11	13.51	24.96	3.47	0.52	85.05
T ₆	10	13.91	23.93	3.30	0.48	85.39

186

187 **3.1. Number of leaves per plant**

188 The number of leaves of water spinach was significantly influenced by different treatments in
 189 maximum cases. The highest number of leaves was found in T₅ (10.73) (Table 1). The results
 190 exhibited that there was the insignificant difference in T₁, T₂, T₄ and T₆ compared to the
 191 control experiment T₀ and a significant difference was found in T₃ and T₅ (Figure 3).
 192 Application of composted chicken bone showed a significant increment of the number of
 193 leaves in water spinach, whereas application of other meet bone fertilizers did not show any
 194 positive effect on the number of leaves of water spinach. On the basis of the number of leaves
 195 of plants, application of composted chicken bone is the best for using, and it is recommended
 196 to use as a soil amendment for the growth of such type of leafy vegetables like water spinach.



197


198 **Figure 3:** Effect of powdered and composted meat bone on leaves number of Water spinach

199

200 **3.2. Root length per plant (cm)**

201 Length of root per plant ranged from 8.81cm to 13.91cm and highest length of root found in
 202 T₆ (13.91cm) (Table 1). Length of root varied significantly in T₆ compared with control
 203 treatment T₀ and there was the insignificant difference among T₁, T₂, T₃, T₄ and T₅ (Figure 4).

204 Distinct A distinct difference was found between T₀ and T₆. So, application of compost of
 205 pigeon bone showed a significant increment of length of root in water spinach whereas
 206 application of powdered chicken bone, beef bones did not show any positive effect on the
 207 root length of water spinach. On the basis of the length of the root of plants application of
 208 composted pigeon bone is the best for uses, and it can be recommended for the plants.




209

210 **Figure 4:** Effect of powdered and composted meat bone on root length of water spinach

211

212 3.3. Shoot length per plant (cm)

213 Length of shoot per plant ranged from 17.64 cm to 24.96 cm. and highest length of shoot was
214 found in T₅ (24.96cm) (Table 1). There was insignificant relationship between T₁, T₂ and T₆
215 but significant difference was found in T₅ compared with the control treatment T₀ (Figure 5).
216 So, application of composted chicken bone showed significant increment of length of shoot in
217 water spinach whereas application of other meat bone fertilizers did not show any positive
218 effect on the shoot length of water spinach. On the basis of the length of shoot of plants
219 application of composted chicken bone is the best for uses and it is recommended for the
220 plants.



221


222 **Figure 5:** Effect of powdered and composted meat bone on shoot length of water spinach

223

224 3.4. Fresh weight per plant (g)

225 Fresh weight per plant ranged from 1.09g to 3.47g, and highest fresh weight of plant was
226 found in T₅ (3.47g) (Table 1). Fresh weight varied significantly in T₄, T₅, T₆ compared with
227 control treatment T₀ but there was the insignificant difference between T₀, T₁, T₂ and T₃ (Fig.
228 4.4.) The observation was T₅ > T₆ > T₄ > T₁ > T₂ > T₃ > T₀. So, application of compost of chicken
229 bones showed a significant increment of fresh weight in water spinach, whereas application
230 of meat bones and pigeon bones did not show any positive effect on the fresh weight. On the
231 basis of the fresh weight of plants application of composted chicken bone is the best for uses,
232 and it is recommended for the leafy vegetables.

233




234

235 **Figure 6:** Effect of powdered and composted meat bone on the fresh weight of water spinach

236

237 3.5. Dry weight per plant (g)

238 Dry weight per plant ranged from 0.08g to 0.52g and highest dry weight was found in T₅
 239 (0.52g) (Table 1). Dry weight varied significantly in all treatments compared to the control
 240 treatment T₀ and there was insignificant difference between T₂ and T₃ as well as T₁ and T₄
 241 (Figure 7). So, application of compost of chicken bone showed significant increment of the
 242 dry weight in water spinach whereas application of other meat bone fertilizers did not show
 243 any positive effect on the dry weight. On the basis of dry weight of plants application of
 244 composted chicken bone is the best for using and it is recommended for the plants.



245


246 **Figure 7:** Effect of powdered and composted meat bone on the dry weight of water spinach

247

248 3.6. Moisture content per plant (%)

249 Moisture content per plant ranged from 64.36% to 85.39% and highest moisture content
250 found in T₆ (85.39%) and the lowest moisture content observed in T₀ (Table 1). The
251 observations are T₆ > T₅ > T₄ > T₁ > T₂ > T₃ > T₀. The following figure shows significant
252 differences when control experiment is compared with other treatment and significant
253 differences are found in T₁ and T₃ but insignificant relationship was found among T₄, T₅ and
254 T₆ treatments (Figure 8). So, application of compost of pigeon bone showed a significant
255 increment of moisture content of water spinach whereas application of chicken bone, beef
256 bones both powdered and composted did not show any positive effect on the moisture content
257 of water spinach. On the basis of the moisture content of plants application of composted
258 pigeon bone is the best for using and it can be recommended for the plants.

259



260

261 | **Figure 8:** Effect of powdered and composted meat bone on the moisture content of water
262 spinach

263

264 4. Summary and Conclusion

265 Powdered and composted meat bones contain sufficient nutrients such as nitrogen,
266 phosphorus, potassium, calcium which are capable to enhance the growth rate of plant. This
267 research has revealed that meat bones both powdered and composted, has significantly
268 increased the growth parameters of water spinach (*Ipomoea aquatica*) such as number of
269 leaves, shoot length, fresh weight, dry weight and moisture content. The result showed that
270 these parameters are mostly increased by using composted chicken bones compared to that of
271 other treatments. Only root length and moisture content of water spinach is increased by
272 using composted pigeon bones. The highest vegetative growth was found by using composted
273 chicken bones, and lowest vegetative growth was found at control experiment. So, the use of

274 | composted chicken bone has the potential to show the highest response on the growth and
275 | yield of water spinach and other types of leafy vegetables.

276 | So, it is recommended for the formers to use composted meat bones, specially composted
277 | chicken bones for the improvement of soil quality and to improve its productivity. On the
278 | contrary, the a load of waste from meat bones will be reduced and will help in waste
279 | management. In the same time, the use of costly and harmful chemical fertilizers will be
280 | reduced and will introduce organic farming, which is environment friendly and beneficial for
281 | human health.

282

283 | **5. References**

284 | BARC (Bangladesh Agricultural Research Council), 2005. Fertilizer Recommendation
285 | Guide. pp.22, 48, 174.

286 | Bhuiyan, N.I. 1994. Crop production trend and need of sustainability in agriculture. A paper
287 | presented in a three-day long workshop on integrated nutrients management for
288 | sustainable agriculture held at SRDI, Dhaka, June, 26-28.

289 | Cayuela, M.L.; Sinicco, T. and Mondini, C. 2009. Mineralization dynamic and biochemical
290 | properties during initial decomposition of plant and animal residues in soil. *Applied*
291 | *Soil Ecology Journal*. 41: 118–127.

292 | Chen, B.H.; Yang, S.H. and Han, H. 1991. Characterization of major carotenoids in water
293 | Convolvulus (*Ipomoea aquatica*) by open-column, thin layer and high-performance
294 | Liquid chromatography. *Journal of Chromatography*, 534: 147-155.

295 | Faruq, U.Z.; Sani, A. and Hassan, L.G. 2002. Proximate composition of sickle pod (*Senna*
296 | *obtusifolia*) leaves. *Nigerian Journal of Applied Science*, 11:157-164

297 | GOB (1985): Yearbook of Agricultural Statistics of Bangladesh 1984-1985, Dhaka:
298 | Bangladesh Bureau of Statistics.

299 | GOB (1989): Yearbook of Agricultural Statistics of Bangladesh 1987-1988, Dhaka:
300 | Bangladesh Bureau of Statistics.

301 | Gibson, T.S.; Chan, K.Y.; Sharma, G. and Shearman, R. (2002). Soil Carbon Sequestration
302 | Utilizing Recycled Organics - A review of the scientific literature. Project 00/01R-
303 | 3.2.6A. The Organic Waste Recycling Unit, NSW Agriculture. Report prepared for
304 | Resource NSW.

305 | Gothberg, A.; Greger, M. and Bengtsson, B. 2005. Accumulation of heavy metals in water
306 | Spinach (*Ipomoea aquatica*) cultivated in Bangkok region, Thailand. *Environmental*
307 | *Toxicology and Chemistry*, 21: 1934-1939.

308 | Gupta, S.; Lakshmi, A.J.; Manjunath, M.N. and Prakash, J. 2005. Analysis of nutrient and
309 | antinutrient content of underutilized green leafy vegetables. *LWT Food Science*.43:
310 | 11–21.

- 311 Hongfei, F. 2011. Evaluation of antioxidant activities of principal carotenoids available in
312 water spinach (*Ipomoea aquatica*). *Journal of Food Composition and Analysis*, 24(2):
313 288-297.
- 314 Jeng, A.S.; Haraldsen, T.K.; Gronlund, A. and Pedersen, P.A. 2006. Meat and bone meal as
315 nitrogen and phosphorus fertilizer to cereals and rye grass. *Nutrient Cycling in*
316 *Agroecosystems*, 76: 183-191.
- 317 Kala. A. and Prakash, J. 2004. Nutrient composition and sensory profile of differently cooked
318 green leafy vegetables. *International Journal of Food Properties*, 7: 659-669.
- 319 MacEwan R.J. 2007. Soil Health for Victoria's Agriculture- Context, Terminology and
320 Concepts. MIS 07898 Final Report. Department of Primary Industries Research
321 Victoria Bendigo, State of Victoria.
- 322 Ogle, B.M.; Ha-Thi, A.G.; Mulokozi, G. and Hambraeus, L. 2001. Micronutrient composition
323 and nutritional importance of gathered vegetables. *Vietnam International Journal of*
324 *Food Science and Nutrition*, 52: 485-499.
- 325 Tate, Robert L. 1987. Soil Organic Matter – Biological and Ecological Effects. John Wiley
326 and Sons, New York.
- 327 USDA (The United States Department of Agriculture), 2005. *Ipomoea aquatica* Forssk.
- 328 Ylivainio, K.; Uusitalo, R. and Turtola, E. 2007. Meat bone meal and fox manure as P
329 sources for ryegrass (*Lolium multiflorum*) grown on a limed soil. *Nutrient Cycling in*
330 *Agroecosystems*, 81: 267-278.
- 331