

1
2
3
4
5
6

Taxonomical investigation on some species of genus *Allium* based on the pollen morphology grain micromorphology

ABSTRACT

Aims: The main aim of this research was to investigate the micromorphological characteristics of the pollen grains in seven species from genus *Allium* belong to three subgenera including *Melanocrommyum*, *Polyprason* and *Cepa* (seven sections).

Methodology: The pollen grains were examined by using Light Microscopy (LM), Scanning Electron Microscopy (SEM) and the pollen grains of one species under TEM (Transmission Electron Microscopy).

Results: The pollen grains were oblate and medium in shape and size. The pollen ornamentation of exine surface, exine ornamentation on sulcus edge, number of exine surface lumina and the state of pollen grain apex in the examined species were different. Semitectate and columellate ectexine with discontinuous endexine were seen in the pollen wall structure (sporoderm).

Conclusion: The dendrogram obtained from the pollen characters in SEM observations by using the numerical taxonomy system (NTSYS) software confirmed phylogram of the studied species obtained from recent phylogenetic research. Our palynological dendrogram can be used for segregation the sections and subgenera taxonomical levels in the studied species of genus *Allium*.

Keywords: *Allium*, Amaryllidaceae, Monocotyledons, Palynology, Phylogeny, Taxonomy

7
8

9
10

11

1. INTRODUCTION

The genus *Allium* is one of the largest Monocotyledons with a wide dispersion in central and south-east Asia, where the species of this genus constitutes a great part of herbaceous societies [1, 2]. Approximately, fifty species of this genus are planted extensively or in local level that economically have great importance. Some of the wild species of this genus are used as edible, medicinal and even decorative plants. Moreover, the wild species of this genus have valuable potential for supplying of human consuming [3].

The taxonomical position of *Allium* as the polymorph genus is sophisticated [4, 5]. Lately, the phylogenetical and biogeographical examination on this genus has been ~~done been~~ done on the endemic species on China [1] and confirmed monophyly in *Allium* by using the phylogenetic analysis of molecular data (ITS nuclear marker and *rps16* chloroplast marker); but, the obtained phylogram from the phylogenetic analysis wasn't confirmed for some subgenera. In their research, three main monophyletic groups were specified: the first clade includes subgenera *Nectaroscordum*, *Amerallium* and *Microscordum*; the second clade includes subgenera *Caloscordum*, *Anguinum*, *Vvedenskya*, *Porphyroprason* and *Melanocrommyum* and the third clade includes subgenera *Butomissa*, *Cyathophora*, *Rhizirideum*, *Allium*, *Reticulatobulbosa* and *Polyprason* [1].

The palynological study of genus *Allium* is sophisticated but driven characteristics from some researches are able to classify taxonomical level on this genus. Majority of these studies were investigated the following characters: exine ornamentation to specify position of the tribe *Allieae* [6, 7], being single sulcus and having extensive sulcus as a predominant morphological pollen character in the genus *Allium* is determined [8]. [9] examined the

Comment [f1]: Why not cite the family AMARYLLIDACEAE in introduction? And never in text?

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

37 species belong to sections *Codonoprasum* and *Allium* and specified the homogeneity of
38 sulcus and pollen grain ornamentation in the genus *Allium*. Moreover, they indicated that
39 some morphological characters of the pollen grain such as sulcus and presence or lack of
40 operculum that have taxonomic importance in the section level. [2] examined the pollen
41 grain morphology in some species of *Allium* (six sections including *Molium*, *Scorodon*,
42 *Brevispatha*, *Codonoprasum*, *Allium* and *Melanocrommyum*) and characterized the shape
43 of pollen grains was prolate and subprolate. Also, in all species extensive sulcus and the
44 smallest and largest pollen grains are belonged to *A. guttatum* (section *Allium*) and *A.*
45 *roseum* (section *Molium*), respectively. Pollen grains from 30 *Allium* taxa belonging to 15
46 sections were recognized in Iran [10]. In this research, the pollens were heteropolar,
47 peroblate to suboblate shape, rugulate to microrugulate, perforate to striate in subgenus
48 *Melanocrommyum* and striate exine ornamentation [10].

49 According to result of [11], the morphological character of sulcus in investigated species in
50 Iran was observed in genus *Allium*, and section *Allium* that this character wasn't observed
51 about in the other sections. [12] observed an extensive sulcus from the beginning to the end
52 of the pollen grains in all taxa in sections of *Rhizirideum*, *Codonoprasum* and *Allium*. In
53 ultrastructure wall of the pollen grains, the exine semitectate and simplicolumellate were
54 observed in all investigated sections. In addition, the exine ornamentation in these sections
55 and related species were heterogeneous and weren't synchronize in classification of these
56 sections and three exine ornamentation including striate-perforate, striate-rugulate-
57 perforate and rugulate-perforate were observed in these species. They also stated that
58 availability of operculum as an apomorphic character and narrow endexine layer as a
59 taxonomic character in the genus *Allium* determined in the sections of *Rhizirideum*,
60 *Codonoprasum* and *Allium* [12]. [15] separated *A. ursinum* in subspecies level based on

Comment [f2]: How many species are there in each section? Show the relevance of studying only few species of them.

Comment [f3]: Authorities of species names!

Comment [f4]: Change. It is not linking with next sentence.

Comment [f5]: Rewrite! It is difficult to understand.

61 the exine ornamentation and stated that this character is an appropriate taxonomic
62 character (granulate-rugulate, rugulate-striate ornamentation in the subspecies *Ucrainicum*
63 and perforate-rugulate ornamentation in the subspecies *Ursinum*). [14] examined three
64 types of pollen grains in seven subgenera and thirteen sections of *Allium* in Pakistan that
65 weren't in agreement with the mentioned classifications and determined the types of *A.*
66 *fedtschenkoanum* (reticulate ornamentation), *A. grifithianum* (rugulate-foveolate
67 ornamentation) and *A. roylei* (subpsilate ornamentation). [15] demonstrated characteristics
68 of the pollen grains including qualitative and quantitative characters of some European
69 species of genus *Allium* including three subgenera (*Allium*, *Amerallium* and *Rhizirideum*)
70 and five sections that explicitly being synchronize with the sections classification.

71
72 The main aim of our research was to evaluate the pollen ~~grain micromorphological~~
73 ~~characters~~ and its ~~comparison with implications on~~ the taxonomy and phylogeny of ~~some~~
74 ~~species of the~~ genus *Allium* ~~including seven species belong to seven different sections. The~~
75 ~~pollen grain characters were reported in some species for the first time. In addition,~~
76 ~~compared with the previous researches, more pollen grain micromorphological characters~~
77 ~~were evaluated.~~

78 2. MATERIAL AND METHODS

79
80 The pollen grains were provided from the collected samples of Main Botanical Garden of
81 Russian Academy of Sciences (MHA), Moscow, Russia. ~~The specimens~~ Samples were
82 collected since May and June 2003 and identified by E. Kalikov (Table 1). Also, taxonomic
83 relationships among the studied species in this research showed in Table 1 [1]. For light
84 microscopy analysis (LM) ~~observations~~, the pollen grains were acetolyzed based on
85

Comment [f6]: How is close *Allium* pollen of other genera in family? Look for other authors who did LM and SEM in their studies for comparisons!
See:
<http://dx.doi.org/10.1590/S0102-33062007000400021>

Comment [f7]: You can reduce the text, and show main pollen character of the groups.

Comment [f8]: Who is?

86 Erdtman's method [16]. Then, mounted ~~preparations with~~ glycerine jelly ~~glass slides~~.
87 Thirty pollen grains were measured by Leitz Light Microscopy (HM-LUX3) from each
88 studied species and images were taken by Dino camera (AM-423).

89 For ~~scanning electron microscopy analysis (SEM) observations~~, non-acetolyzed pollen
90 grains were put on metal ~~legs stubs~~ and transferred to the EMITECH sputter coater for
91 coating by gold-paladium (K450X). Finally, the obtained micrographs were taken by VEGA-
92 TESCAN Scanning Electron Microscopy. Terminology for LM and SEM observations were
93 explored according to the following references [17, 18, 19].

94 For construction the dendrogram, the obtained characters by SEM observations from the
95 pollen grains were coded (Table 4). Finally, the NTSYS Software [20] was used along
96 based on Single Linkage method and UN1 similarity coefficient (binary coefficient) (Table
97 5). For TEM preparation, the pollen grains were fixed by 2% osmium tetroxide and stained
98 by uranylacetate solution [21]. Then, dehydrated in ethanol series and were put in acetone.
99 Finally, embedded in Epon mixture (Epon 812, Epon Harter DDSA, Epon Harter MNA)
100 according to the standard method of [22]. Ultrathin sections of the pollen grains were
101 obtained by an ultramicrotome (LKB 8800), then stained with lead citrate (LKB 8800,
102 Ultratome III) [21]. The micrographs were made by using a JEOL-JEM-100B Transmission
103 Electron Microscopy.

104 3. RESULTS AND DISCUSSION

105 ~~Based on the LM observations,~~ the pollen grains in the studied species were observed
106 ~~oblate~~ in shape and medium in size (Table 2 and Fig. 1). In SEM micrographs, the pollen
107 grain characters were different in these species (Table 3, Figs. 2 and 3). The exine
108 ornamentation on surface and the exine ornamentation on ~~sulcus edge~~ were different and
109
110

Comment [f9]: Very confuse an long.
Begin with general charactes (size and shape), then describe apertures (size and membrane, or edge here), and exine (thickness and ornamentation). Organize point to point, because it is easier for reader get your information.
Try your text in short paragraphs, yours are too long!

Formatted: Highlight

Comment [f10]: Provide a plate with the main edge ornamentation. It is difficult to see on you pollen plates!

Formatted: Highlight

111 these characters for each species are expressed as follows: striate exine ornamentation on
112 surface and sulcus edge for *A. altissimum*, perforate-striate exine ornamentation on surface
113 and perforate on sulcus edge for *A. fetisowii*, striate-perforate exine ornamentation on
114 surface and striate on sulcus edge for *A. backhousianum*, exine ornamentation on surface
115 and sulcus edge of macrostriate for *A. karataviense*, striate-psilate exine ornamentation on
116 surface and psilate-striate exine ornamentation on sulcus edge for *A. obliquum*, perforate-
117 striate exine ornamentation on surface and psilate exine ornamentation on sulcus edge for
118 *A. rosenbachianum*, perforate-microstriate exine ornamentation on surface and sulcus
119 edge for *A. schoenoprasum* (Table 3). The pollen grains of *A. fetisowii* and *A.*
120 *schoenoprasum* species were acute in the end and in the other species were obtuse (Table
121 3). The size of lumina and muri in the studied species also was different. The scope of
122 changes in lumina's size in surface of the pollen grain almost was observed similar in *A.*
123 *altissimum*, *A. backhousianum* and *A. karataviense* (0.06- 0.13 μm) (Table 3). The number
124 of exine surface lumina, was the least (4 lumina at $2 \mu\text{m}^2$) for *A. altissimum* and the most
125 (30 and 31 lumina at $2 \mu\text{m}^2$) for *A. fetisowii* and *A. backhousianum* species, respectively
126 (Table 3).

127
128 The results of *A. schoenoprasum* TEM micrographs belongs to section *Schoenoprasum*
129 and subgenus *Cepa* revealed that the ectexine is semitectate, infratectum is
130 simplicolumellate, foot-layer is discontinuous and endexine layer is very thin and
131 discontinuous (Fig. 4).

132
133 This research indicated that there were less differences about the LM observations (shape
134 and size); but, the dendrogram of pollen micromorphological characters from SEM

Formatted: Highlight

Comment [f11]: PER or BY or /

Comment [f12]: What does it mean?

135 observations provided useful valuable taxonomical characters on the subgenera and
136 sections classifications (Table 4 & 5, Fig. 5).

137
138 Various shapes of pollen grains were observed in ~~this genus~~*Allium spp.* on previous
139 investigations and the ~~dominant-main~~ shape of the pollen grains in this genus is oblate
140 shape. Medium pollen type was observed in the most species ~~studied here of this genus~~
141 ~~during this research~~ and previous research [2, 9, 10, 11, 13, 14, 15].

142
143 ~~Exine ornamentation was varied in genus Allium [2, 9, 10, 11, 13, 14, 15]. Our results show~~
144 ~~that varied exine ornamentation with the other characters on SEM observation as useful~~
145 ~~taxonomic characters in section level.~~

146
147 Our research on the genus *Allium* confirms previous palynological research on pollen wall
148 structure on *Codonoprasum*, *Allium* and *Rhizirideum* sections [9, 12].

149
150 The perforate-striate exine surface ornamentation for two species including *A. fetisowii* and
151 *A. rosenbachianum* belong to the subgenus *Melanocrommyum* was alike; but, these two
152 species were different in the exine ornamentation on sulcus edge, the number of exine
153 surface lumina, the state of pollen grain apex and the size of lumina and muri.

154 The variations of exine ornamentation on the surface and sulcus edge for the other species
155 were in agreement with the performed sections classification. Moreover, the number of
156 exine surface lumina for each species was also different and the least and the most
157 number of lumina was counted in the subgenus *Melanocrommyum*. Our results also
158 indicated that *A. obliquum* belong to section *Oreiprason* and subgenus *Polyprason* had no

Formatted: Font: Italic

Comment [f13]: How?

159 lumina in the exine surface. In the studied species in Iran, the exine surface ornamentation
160 without lumina was observed in the subgenus *Reticulatobulbosa* in section *Campanulata*
161 and also in the subgenus *Polyprason* in section *Falcatifolia* [11]. Therefore, the exine
162 surface ornamentation without lumina can be defined as a useful micromorphological
163 character in the subgenus *Polyprason* and its related sections. In flora of Pakistan *A.*
164 *rosenbachianum* on section *Megaloprasum*, *A. schoenoprasum* belongs to section
165 *Schoenoprasum* and *A. roylei* in section *Oreiprasum* were placed on *A. roylei* types based
166 on the pollen grain morphology [14].

167
168 The dendrogram of the [palynological-pollen](#) characters analysis was in agreement with the
169 taxonomical ranking and previous phylogram by [1] on these taxa in subgenus and section
170 levels. The phylogram of the studied species were distinguished based on [1]-[research](#). [4]
171 [These authors](#) had done the extensive phylogenetic analysis by molecular data (ITS
172 nuclear marker and *rps16* chloroplast marker) on *Allium* genus. The final dendrogram of
173 our palynological investigation was in agreement with the obtained phylogram by [1]
174 [research](#). According to the pollen [grain](#)-dendrogram obtained from analysis by SEM
175 observations in subgenus *Melanocrommyum*, *A. fetisowii* introduced as a sister group with
176 the other species in this subgenus. In clade *Melanocrommyum*, *A. fetisowii* in section
177 *Longibidentata* determined as the sister group of the remaining species of the subgenus
178 *Melanocrommyum* based on the phylogenetic information of molecular markers [1]. *A.*
179 *schoenoprasum* belongs to subgenus *Cepa* and section *Schoenoprasum* was different from
180 the other species in SEM observation, although in the state of pollen grain apex was similar
181 to *A. fetisowii*. The results also indicated that *A. altissimum*, *A. backhousianum* and *A.*
182 *karataviense* species with similar size in lumina were placed in the subgenus

183 *Melanocrommyum*. The palynological dendrogram in this study put the species *A.*
184 *altissimum* and *A. backhousianum* from subgenus *Melanocrommyum* and in sections
185 *Procerallium* and *Acmeperala* together. Moreover, according to the molecular studies,
186 these two sections were put together [1].

187 **4. CONCLUSION**

188 The dendrogram of palynological analysis data in the related species was in parallel with
189 the divisions of sections and subgenera taxonomical rank and the phylogram of
190 phylogenetic studies. Also, our research on the genus *Allium* confirms some previous
191 palynological research.
192
193

194 **COMPETING INTERESTS**

195 The authors of this manuscript declare that they have no competing interests.
196
197
198
199
200
201

202 **REFERENCES**


- 203 1. Li QQ, Zhou SD, He XJ, Yu Y, Zhang YC, Wei XQ. Phylogeny and biogeography of *Allium*
204 (*Amaryllidaceae*: *Allieae*) based on nuclear ribosomal internal transcribed spacer and chloroplast
205 *rps16* sequences, focusing on the inclusion of species endemic to China. *Ann Bot.* 1990; 106:709-33.
- 206 2. Özhatay N, Kocyigit M. Pollen morphology of *Allium* species (*Liliaceae*) in European Turkey and
207 around Istanbul. *Phytol.* 2009;15(2):199-208.
- 208 3. Fritsch RM, Abbasi M. Taxonomic Review of *Allium* subg. *Melanocrommyum* in Iran, Germany,
209 Leibniz-Institut für Pflanzengenetik und Kulturpflanzenforschung Gatersleben, 2013.
210

- 211 4. Klaas M. Applications and impact of molecular markers on evolutionary and diversity studies in the
212 genus *Allium*. *Plant Breed.* 1998;117:297-308.
- 213 5. Fritsch RM, Friesen N. Evolution, domestication and taxonomy. In: Rabinowitch, H. D. & Currah L.
214 (eds.): *Allium crop science: recent advances*, Wallingford, New York, 5–30, 2002.
- 215 6. Kosenjo VN, Kudryashova GL. Palynomorphology of the family Alliaceae, *Bot. Zhurnal* (st.
216 Petersburg), 1995; 80(6):5-17.
- 217 7. Radulescu D. Recherches morpho-palynologiques sur la famille Liliaceae, *Acta Bot. Horti. Bucur.*
218 1973; 133-248.
- 219 8. Harley MM, Zavada MS. Pollen of the monocotyledons: Selecting characters for cladistic analysis.
220 *In: Monocots: Systematics and Evolution*, K.L. Melbourne, Wilson and D.A Morrison (Eds).
221 CSIRO. 2000; 194-213.
- 222 9. Güler U, Pehlivan S. Pollen morphology of some species belonging to *Codonoprasum* and *Allium*
223 sections of *Allium* (Liliaceae-Alliaceae) genus, *Biologia.* 2006; 61(4):449-55.
- 224 10. Neshati F, Fritsch RM, Zarre S. Pollen morphology of some *Allium* L. species (Alliaceae) from Iran.
225 *Plant Diversity and Evolution.* 2009; 127 (4): 433–451.
- 226 11. Namin HH, Mehrvarz SS, Zarre S, Fritsch RM. Pollen morphology of selected species of *Allium*
227 (Alliaceae) distributed in Iran, *Nordic J. Bot.* 2009; 27(1):54-60.
- 228 12. Özler H, Pehlivan S. Pollen morphology of *Allium* L. (Liliaceae) taxa in Turkey, *Bangladesh J. Bot.*
229 2010; 39(1):37-46.
- 230 13. Bareemizadeh F, Maassoumi SM, Ghasempour H, Karimi N. Pollen morphology of the genus *Allium*
231 in comparison with genus *Calochortus* of order Liliales, *Int. J. Biosci.* 2014; 4(12):237-43.
- 232 14. Perveen A, Qaiser M. Pollen flora of Pakistan–lxviii. Alliaceae, *Pak. J. Bot.* 2015; 47(1): 263-68.

- 233 15. Wronska-Pilarek D, Halbrither H, Krazyminska A, Rzyminska L, Bednorz J, Bocianowski J. Pollen
234 morphology of selected European species of the genus *Allium* L. (Alliaceae), Acta Sci. Pol. Hortorum
235 Cultus. 2016; 15(4):65-84.
- 236 16. Erdtman G. The acetolysis technique: a revised description, Svensk Botanisk Tidskrift, 1960;
237 54:561-64.
- 238 17. Tokarev P. Morphology and ultrastructure of pollen grain. Moscow, KMK Sci, Press Ltd., 2003.
- 239 18. Punt W, Hoen PP, Blackmore S, Nilsson S, Le Thomas A. Glossary of pollen and spore terminology,
240 Rev. Palaeobot. Palynol. 2007; 143:1-81.
- 241 19. Hesse M. Pollen Terminology (An illustrated handbook). Vienna, Springer, 2009.
- 242 20. Rohlf FJ. NTSYS-PC: Numerical Taxonomy and Multivariate Analysis System. Exeter Software,
243 Setauket, NY, USA, 1993.
- 244 21. Reynolds ES. The use of lead citrate at high pH as an electron opaque stain in electron microscopy.
245 Journal of cell biology. 1963; 17: 208–212.
- 246 22. Weakley BS. A Beginner's Handbook in biological Electron Microscopy. Churchill Livingstone,
247 Edinburg, 252 pp, 1972.

Comment [f14]: HESSE, M.; ZETTER, R.; HALBRITTER, H.; WEBER, M.; BUCHNER, R.; FROCHRADIVO, A.; ULRICH, S. 2009. **Pollen Terminology: an illustrated handbook**. Wien: Springer.

UNDER PEER REVIEW



255

256

Table 1. Taxonomic relationships and collecting data from the studied species of *Allium* (Amaryllidaceae). [1].

Taxon	Section	Subgenus	Collector and date of collecting
<i>Allium altissimum</i> Regel.	<i>Procerallium</i>	<i>Melanocrommyum</i>	E. Kalikov, 30.5.2003, gathering from MHA
<i>Allium fetisowii</i> Regel.	<i>Longibidentata</i>	<i>Melanocrommyum</i>	E. Kalikov, 30.5.2003, gathering from MHA
<i>Allium backhousianum</i> Regel.	<i>Acmeperata</i>	<i>Melanocrommyum</i>	E. Kalikov, 10.6.2003, gathering from MHA
<i>Allium karataviense</i> Regel.	<i>Miniprasum</i>	<i>Melanocrommyum</i>	E. Kalikov, 30.5.2003, gathering from MHA
<i>Allium obliquum</i> L.	<i>Oreiprasum</i>	<i>Polyprason</i>	E. Kalikov, 30.5.2003, gathering from MHA
<i>Allium rosenbachianum</i> Regel.	<i>Megaloprasum</i>	<i>Melanocrommyum</i>	E. Kalikov, 30.5.2003, gathering from MHA
<i>Allium schoenoprasum</i> L.	<i>Schoenoprasum</i>	<i>Cepa</i>	E. Kalikov, 30.5.2003, gathering from MHA

Formatted: Font: Italic

Formatted: Font: 10.5 pt

Formatted: Font: (Default) Helvetica, 10.5 pt, Bold, Portuguese (Brazil)

Formatted: Font: 10.5 pt

Comment [f15]: Cite collector's numbers, or herbarium ones. A number is needed!

257

258

259

260

261

262

263

264

265

266

267
268
269
270
271
272
273
274
275
276
277
278
279

Table 2. The obtained pollen data from the Light Microscopy studies Pollen measurements of the studied species of Allium (Amaryllidaceae).

Taxon	Pollen size	Polar axis (µm)	Equatorial axis (µm)	P.E	Pollen	Pollen
	(µm)	Min-Mean±SD-Max	Min-Mean±SD-Max		shape	typesize
<i>A. altissimum</i>	35.18±6.52	17.00-18.28±1.24-21.00	27.50-35.18±6.52-46.00	0.5	Oblate	Medium
<i>A. fetisowii</i>	30.74±5.13	15.00-16.16±1.51-20.00	22.50-30.74±5.13-37.50	0.5	Oblate	Medium
<i>A. backhousianum</i>	29.97±2.34	12.00-16.22±1.79-17.50	27.50-29.97±2.34-35.00	0.5	Oblate	Medium
<i>A. karataviense</i>	31.39±4.92	12.00-16.84±2.03-20.00	22.50-31.39±4.92-41.00	0.5	Oblate	Medium
<i>A. obliquum</i>	31.56±6.09	12.50-16.16±1.60-20.00	25.00-31.56±6.09-41.00	0.5	Oblate	Medium
<i>A. rosenbachianum</i>	33.70±1.92	14.00-19.60±3.63-25.00	30.00-33.70±1.92-37.50	0.6	Oblate	Medium
<i>A. schoenoprasum</i> var. <i>sibiricum</i>	30.82±5.08	15.00-16.46±2.24-20.00	25.00-30.82±5.08-39.00	0.5	Oblate	Medium

Formatted: Font: Italic

Comment [f17]: Smaller or bigger?

Comment [f16]: Pollen size is the greater value of a diameter (equat or polar one). In case of monolcate pollen, there are 2 equatorial diameters (a smaller and a bigger) and one pollen diameter. So, I don't know what POLLEN SIZE means here!

Comment [f18]: No applied in this case! See comments about pollen size.

Comment [f19]: The same

Formatted: Left

Formatted: Left

Formatted: Left

Formatted: Left

Formatted: Left

Formatted: Left

Formatted: Left

280
281
282
283
284
285
286
287
288
289
290

291 **Table 3. The obtained pollen data from the Scanning Electron Microscopy from Pollen data of the studied species of**
292 **Allium (Amaryllidaceae).**

Taxon	Exorn	Sd	S	L	M	P
<i>A. altissimum</i>	Striate	Striate	Obtuse	0.06-0.13	0.13-3.46	4
<i>A. fetisowii</i>	Perforate-striate	Perforate	Acute	0.06-0.20	0.13-0.86	30
<i>A. backhousianum</i>	Striate-perforate	Striate	Obtuse	0.06-0.13	0.13-0.40	31
<i>A. karataviense</i>	Macrostriate	Macrostriate	Obtuse	0.06-0.13	0.06-0.40	12
<i>A. obliquum</i>	Striate-psilate	Psilate-striate	Obtuse	-	-	-
<i>A. rosenbachianum</i>	Perforate-striate	Psilate	Obtuse	0.06-0.26	0.06-1.20	27

Comment [f20]: But if it is MACROSTRATE, it hasn't lumina!

A. schoenoprasum var. *sibiricum* Perforate-microstriate Perforate-microstriate Acute 0.06-0.40 0.06-1.60 22

293

Exorn: Exine ornamentation, Sd: Sulcus edge ornamentation, S: The state of pollen grain apex, L: Lumina size, M: Margin size, P: The number of exine surface lumina. 295

Comment [f21]: Do you mean sulcus membrane ornamentation? Or margin?

Comment [f22]: Width?

Comment [f23]: Number, how? Do you mean density? Lumina by area? I can't understand!

296

297

298

299

300

301

302

303

304

305

306

307

308

Table 4. The pollen traits from Scanning Electron Microscopy and coding of these characters by the analysis with

NTSYS software.


Comment [f24]: The same of table 3! You must change.

Taxon	Exorn (code)	Sd (code)	S (code)	L (code)	M (code)	P (code)
-------	--------------	-----------	----------	----------	----------	----------

<i>A. altissimum</i>	Striate (1)	Striate (1)	Obtuse (1)	0.06 (1)-0.13 (1)	0.13 (1)-3.46 (5)	4 (1)
<i>A. fetisowii</i>	Perforate-striate (2)	Perforate (2)	Acute (2)	0.06 (1)-0.20 (2)	0.13 (1)-0.86 (2)	30 (5)
<i>A. backhousianum</i>	Striate-perforate (3)	Striate (1)	Obtuse (1)	0.06 (1)-0.13 (1)	0.13 (1)-0.40 (1)	31 (6)
<i>A. karataviense</i>	Macrostriate (4)	Macrostriate (3)	Obtuse (1)	0.06 (1)-0.13 (1)	0.06 (2)-0.40 (1)	12 (2)
<i>A. obliquum</i>	Striate-psilate (5)	Psilate-striate (4)	Obtuse (1)	(0)	0	0
<i>A. rosenbachianum</i>	Perforate-striate (2)	Psilate (5)	Obtuse (1)	0.06 (1)-0.26 (3)	0.06 (2)-1.20 (3)	27 (4)
<i>A. schoenoprasum</i> var. <i>sibiricum</i>	Perforate-microstriate (6)	Perforate-microstriate (6)	Acute (2)	0.06 (1)-0.40 (4)	0.06 (2)-1.60 (4)	22 (3)

311
312
313
314
315
316
317
318
319
320
321
322
323

UNDER PEER REVIEW



324
325
326
327
328
329
330

Table 5. The similarity coefficients of the studied species of Allium (Amaryllidaceae) compared with the others by using the

NTSYS software.

332

Taxon	a	b	c	d	e	f	g
a	1.00						
b	0.40	1.000					
c	0.76	0.40	1.00				
d	0.54	0.222	0.66	1.00			
e	0.22	0.00	0.22	0.22	1.00		
f	0.40	0.400	0.40	0.54	0.22	1.00	
g	0.22	0.40	0.22	0.40	0.00	0.40	1.00

333


334
335


a. *A. altissimum*, b. *A. fetisowii*, c. *A. backhousianum*, d. *A. karataviense*, e. *A. obliquum*, f. *A. rosenbachianum*, g. *A. schoenoprasum* var. *sibiricum*.

336
337
338
339
340
341
342
343
344
345
346
347
348

UNDER PEER REVIEW

349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365





367 | Fig. 1. [Light microscopy photographs](#)Pollen grains of the studied species of *Allium* (Amaryllidaceae), seen on
368 | [LM](#). a. *A. altissimum*, b. *A. fetisowii*, c. *A. backhousianum*, d. *A. karataviense*, e. *A. obliquum*, f. *A.*
369 | *rosenbachianum*, g. *A. schoenoprasum* var. *sibiricum*. (Light Microscopy, 1000X, scale: 10 μ m)

370

371

372

373

374

375

376

377

378

379

380

381

382


383

384

385

386

387



388

389 Fig. 2. The Scanning Electron Microscopy micrographs of Pollen grains of the studied species of Allium
390 (Amaryllidaceae), on SEM: (for each species, the equatorial view of pollen grain, exine ornamentation, sulcus
391 edge exine ornamentation and the state of pollen grain apex have been determined, respectively). *A. altissimum*
392 (a–d), *A. fetisowii* (e–h), *A. backhousianum* (i–l) and *A. karataviense* (m–p). (Scale bar: 20 μm for a, e, i & m., scale
393 bar: 2 μm for b, c, d, f, g, h, j, k, l, n, o & p.)

394


395

396

397

398

399



400

401 Fig. 3. ~~The Scanning Electron Microscopy micrographs~~ Pollen grains of the studied species of Allium
402 (Amaryllidaceae), on SEM: (for each species, the equatorial view of pollen grain, exine ornamentation, sulcus
403 edge exine ornamentation and the state of pollen grain apex have been determined, respectively). *A. obliquum*
404 (a–d), *A. rosenbachianum* (e–h) and *A. schoenoprasum* var. *sibiricum* (i–l). (Scale bar: 20 μ m for a, e & i., scale
405 bar: 2 μ m for b, c, d, f, g, h, j, k & l)

406


407

408

409

410

411
412
413



414

Fig. 4. The Transmission Electron Microscopy micrographs Ultra-thin section on pollen grain of *A. schoenoprasum* var. *sibiricum* (Amaryllidaceae): a. Cross section of pollen grain (Scale bar: 5 µm), b: Cross section of pollen wall structure (Scale bar: 1 µm).

415

416

417

Formatted: Font: Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

418

419

420

421

422

423

424

425


426

427


428

429

UNDER PEER REVIEW



130



131

Fig. 5. Dendrogram obtained from [implication the analysis of pollen data morphology on](#) and taxonomic relationships of the studied species [of *Allium* \(Amaryllidaceae\)](#).

434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461

UNDER PEER REVIEW

