

Original Research Article

Mapping a climate change vulnerability index: An assessment in agricultural, geological and demographic sectors across the districts of Karnataka

Abstract

Climate change is a continuous phenomenon ~~and~~ over hundreds of years; the atmosphere has changed considerably around the world. Karnataka, has the second largest drought prone area in the country next only to Rajasthan. Assessment of vulnerability index **could** play a major role in designing appropriate mitigation and adaptation policies to overcome the impacts of climate change. The vulnerability assessment is an exhaustive procedure **influenced** by a large number of indicators. This study **attempted** to capture a picture of composite vulnerability index of different districts of Karnataka by considering agronomic, climatic and demographic indicators. The secondary data on climatic, agronomic and demographic factors were collected from various sources for the year 2017-18. The findings of the study ~~as~~ shown that the average vulnerability index for 30 districts is 0.577 and 16 districts placed above the average composite vulnerability index level. Bidar (0.655) is the most vulnerable district followed by Kolar (0.658) and Yadgir (0.638) districts. Shivamogga (0.440), Davanagere (0.486) and Udupi (0.486) was the districts exhibited the least vulnerability to changing climate. The results suggested that agricultural and climatic indicators are the major factors which influenced vulnerability. So special attention should be given to agricultural and climatic sectors to minimize the impacts of climatic change in the most vulnerable districts.

Key words: Vulnerability index, Climate change, Per capita income, Sensitivity, Exposure and Adaptability

Introduction: Agricultural economy in Karnataka is largely influenced by agro-climatic factors, water and other resource **contributed by** farmers, technology, infrastructures, tradition and social capital and also by the market forces of demand and supply. Karnataka has the second largest drought prone area in the country next only to Rajasthan and water availability is one of the

Comment [H1]: determined

30 major concerns in the state. On average, Karnataka's annual rainfall was 1,151 mm, of which 80
31 per cent was received during the southwest monsoon, 12 per cent in the post monsoon period, 7
32 per cent during summer and 1 percent in **rabi**. Groundwater potential of the area depends on
33 rainfall and efforts to recharge. Changing in climatic conditions directly affects the hydrological
34 cycle and gradually the groundwater table. Obviously the economic impact of climate change
35 will severely affect the food security as well as livelihood security including health security of
36 farmers (Chandrakanth, M. G., 2015).

Comment [H2]: explain a little bit

37 Climate change is a continuous phenomenon ~~and~~ over hundreds of years, since the
38 atmosphere has changed considerably around the world. However, the pace and pattern of
39 changes in climatic factors in recent decades have turned into a matter of concern. **Especially**,
40 since it is very hard to comprehend the effect of change in climatic factors at the small scale level
41 **even, say,** at block or district levels (Raju *et al.*, 2017). The Intergovernmental Panel on Climate
42 Change (IPCC), in its second evaluation report (Anonymous, 1996), characterized vulnerability
43 as the degree to which environmental change may harm or damage a system. It inferred that
44 vulnerability not only depends on a system of sensitivity, but also in addition, on its capacity to
45 adjust to new climatic conditions, the level of economic development and institutions.

Comment [H3]: for example??

46 **2. Methodology:**

47 The key target of this assessment is to analyze the climate vulnerability of different
48 sectors across the districts of Karnataka. ~~Keeping in view of~~ the appraisal of the information
49 relating to different indicators pertaining from agriculture year 2013-14 to 2017-18, and were
50 collected from various sources such as Karnataka State Natural Disaster Monitoring Centre
51 (KSNDMC), Directorate of Economics and Statistics (DES) and Central Groundwater Board
52 (CGB).

53 The vulnerability assesment is an exhaustive procedure influenced by a large number of
54 indicators. However only the most significant and appropriate indicators were chosen for
55 calculation of vulnerability index based on exposure, sensitivity and adaptability to varied
56 climate. Parameters used in this study include

57 **Climatic components:** Variance of annual rainfall (mm²), Variance of South-West monsoon
58 (mm²), Variance of maximum temperature, Variance of minimum temperature and Variance of
59 average temperature (°C).

60 **Agricultural Components:** Geographical area (GA) (ha), Forest area (% of GA), Area under
61 food crops (% of Gross Cropped Area(GCA)), Net sown area (% of GA), Livestock population
62 (No. per ha of GCA), Irrigated area (% of GCA), Cropping intensity (%), Productivity of major
63 crops (Paddy, Ragi, Jowar, Sugarcane, Maize, Groundnut, Sunflower, Cotton, Arecanut,
64 Coconut, Redgram, Cowpea, Chilli), Depth of Groundwater (meter below ground level), Per
65 capita income (Rs per person).

66 **Demographic components:** Density of male population (Persons per sq. ha of GA), Density of
67 female population (Persons per sq. ha of GA), Literacy rate of male (%) and Literacy rate of
68 female (%).

69 Composite Vulnerability Index (CVI) is assessed for each district by using Iyenger and
70 Sudarshan (1982) technique for unequal weight. The assessed CVI is a total of three sub-sectors
71 specifically Climatic Vulnerability, Agriculture Vulnerability and Demographic Vulnerability.
72 Development of vulnerability index and Composite Vulnerability Index comprised of several
73 steps.

74 **Step 1:** The information compiled pertaining to three components was transformed into suitable
75 estimation units and arranged in a rectangular matrix with rows representing districts and
76 columns representing indicators.

77 **Step 2:** Since each of the sub-component was measured using different units and scale, they
78 were needed to be normalized first. The procedure developed by Anand and Sen (1994) for
79 construction of the Human Development Index (HDI) was used to normalize indicators. In any
80 case, before doing normalization, it was imperative to distinguish the functional relationship
81 between the indicators and vulnerability. Two kinds of practical relationships, vulnerability
82 increases with the increase (decrease) in the value of indicators are conceivable.

83 For direct relationship:
$$Y_{ij} = \frac{X_{ij} - \text{Min}(X_{ij})}{\text{Max}(X_{ij}) - \text{Min}(X_{ij})}$$

84

85 For indirect relationship:
$$Y_{ij} = \frac{\text{Max}(X_{ij}) - X_{ij}}{\text{Max}(X_{ij}) - \text{Min}(X_{ij})}$$

86

Comment [H4]: what does this mean? rainfall?
since the unit was mm

87 Where,

88

89 Y_{ij} was the normalized value

90 X_{ij} was the actual value of the indicator

91 $Min(X_{ij})$ and $Max(X_{ij})$ were the minimum and maximum actual values

92 **Step 3:** The degree of vulnerability (\bar{y}_i) is assumed to be the linear sum of X_{ij} as

$$\bar{y}_i = \sum_{j=i}^k w_j x_{ij}$$

93 Where w_j 's were weights and were determined by

$$w_j = \frac{c}{\sqrt{\text{var}(x_{ij})}}$$

94 Where c was the normalizing constant

$$c = \left[\sum_{j=1}^k \frac{1}{\sqrt{\text{var}(x_{ij})}} \right]^{-1}$$

95

96 The vulnerability index lies in the range of 0 and 1. A value of 1 indicates greatest vulnerability
97 and 0 shows absence of vulnerability.

98

99 **3. Results and Discussion:**

100 The Sector wise vulnerability indices and composite index were constructed for all the 30
101 districts of Karnataka. The districts were ranked based on extent of vulnerability index.

102 **3.1 Component wise vulnerability index**

103 **3.1.1 Climatic Vulnerability index**

104 To construct district level vulnerability index five climatic variables were used and the
105 results were presented in the Table 1. The results shown that the Kalaburagi district had the
106 highest climate vulnerability index of 0.747 followed by Kolar (0.720), Bidar (0.720), Raichur
107 (0.712) and Yadgir (0.711) districts. The districts of Kodagu and Udupi had only 0.278 and
108 0.215 vulnerability index respectively, the least in Karnataka state. We can observe highest
109 vulnerability index values in northern districts of Karnataka which was due to large variations in
110 rainfall and temperature during the year. These were the key determinant indicators which
111 explained high climatic fluctuations among districts.

112

113 For instance, prevalence of a high degree of anticipated change in mean precipitation and
114 high inconsistency in minimum and maximum temperatures drove Kalaburagi district to the top
115 of the chart.

116 **3.1.2 Agriculture Vulnerability index**

117 Based on functional relationship of the indicators, Vulnerability index for agricultural parameters
118 were calculated for each district and is presented in Table 2.

119 Kodagu district secured first place with a total vulnerability index value of 0.787
120 followed by Bidar (0.761), Kolar (0.741) and Chitradurga (0.732) districts. Whereas Davanagere
121 has been rated as the least vulnerable district (0.524). Lower productivity, declined forest area,
122 high groundwater table level, lower cropping intensity and low per capita income are the major
123 factors which influence the high level of sensitivity leading to higher vulnerability index.

124 In general Kodagu, Bidar, Kolar and Chitradurga districts are most sensitive districts and
125 highly vulnerable to climate change. On the contrary, Davanagere, Shivamogga, Bellary and
126 Bengaluru Urban districts are less sensitive and less vulnerable to changing climate.

127 **3.1.3 Demographic vulnerability index**

128 The districts having high population density coupled with a lower rate of literacy were
129 identified as vulnerable districts with respect to demographic features.

130 Bengaluru Urban (0.579) district occupied the first place whereas Dakshina Kannada
131 (0.039) district was placed in the last position with respect to demographic vulnerability (Table
132 3). Yadgir (0.449), Raichur (0.353), Chamarajnagara (0.335) and Kalaburagi (0.294) were the
133 districts having higher degree of vulnerability index next to Bengaluru Urban district. The
134 coastal districts of Dakshina Kannada, Udupi (0.051) and Uttara Kannada (0.055) had lower
135 vulnerability index and higher adaptive capacity to changing climate because of high literacy rate
136 and lower population density.

137 **3.2 Composite vulnerability index**

138 Agricultural indicators, climatic indicators and demographic indicators were used to
139 construct composite vulnerability index. Table 4 shows district wise composite vulnerability
140 index which was calculated using all the three sub-components (Agricultural, Climatic and
141 Demographic). Average composite vulnerability index for 30 districts is 0.584 and 17 districts

142 placed above the average composite vulnerability index level. Districts having high composite
143 vulnerability index would be more vulnerable to climate change. Bidar (0.577) district had the
144 highest composite vulnerability index followed by Kolar (0.658) and Yadgir (0.638). These
145 districts were the most vulnerable districts and the results were in line with the report submitted
146 by Anonymous (2011) which used composite vulnerability index. They reported that Kalaburagi
147 and Dakshina Kannada districts were the most and the least vulnerable districts, respectively.
148 Higher composite index was observed mainly due to higher sensitivity of agricultural sector and
149 larger exposure to climate change. Composite vulnerability index was lower for Shivamogga
150 (0.440), Davanagere (0.486) and Udupi (0.486) districts because these districts shown less
151 vulnerability in terms of agriculture and climatic indicators. In addition also demographic
152 variables such as population density and literacy rate contributed to lowering of composite
153 vulnerability index . At district level, contribution of each sub-component to composite index
154 was not uniform. In general agricultural indicators contributed foremost, followed by climatic
155 and demographic indicators. A study conducted by Hiremath and Shiyani (2013) reported that
156 agriculture and occupation sector were the major sectors which have contributed most to
157 composite vulnerability index in Saurashtra.

158
159

160 **4. Conclusion:**

161 Karnataka was the second most drought prone state after Rajasthan. District wise
162 vulnerability mapping was carried out to calculate the vulnerability index of each district. Sector
163 wise indicators were selected based on exposure, sensitivity and adaptive capacity to climate
164 change. All the indicators were considered to calculate composite vulnerability index. Findings
165 of the analysis showed that Bidar was the most vulnerable district and Shivamogga was the least
166 vulnerable. Major component which contributed to composite index was the Agricultural
167 vulnerability. The results of agricultural vulnerability index analysis has highlighted the
168 indicators such as productivity of the major crops, cropping intensity and per capita income were
169 the major drivers in determining the vulnerability of districts. Therefore, it is suggested that
170 Bidar, Kolar, Yadgir, Koppal and Chtradurga districts should be considered as the priority to
171 minimize the degree of vulnerability. There was a need to take up adaptive practices such as
172 varieous selection according to prevailing weather, contingent cropping, soil and water

173 conservation measures, in-situ moisture conservation, rainwater harvesting and augmenting
174 recharging of groundwater for supplementary irrigation. In addition, better education and
175 infrastructure development in rural areas would also play a catalytic role in enhancing adaptive
176 capacity of these districts.

177 **5. References:**

178 ANAND, S. AND SEN, A., 1994, Human development index : Methodology and measurement.
179 United Nations Development Programme, New York : Human Development Report
180 Office.

181 ANONYMOUS, 1996, Climate change second assessment report (1995). Intergovernmental
182 panel on climate change. Geneva, Switzerland.

183 ANONYMOUS, 2011, Karnataka Climate Change Action Plan, Bangalore Climate Change
184 Initiative–Karnataka, Bengaluru (India) Final Report, Government of Karnataka.

185 CHANDRAKANTH, M. G., 2015, *Water resource economics: Towards sustainable use of water*
186 *for irrigation in India*, Springer, Switzerland.

187 HIREMATH, D. B. AND SHIYANI, R. L., 2013, Analysis of vulnerability indices in various
188 agro-climatic zones of Gujarat. *Indian Journal of Agricultural Economics*, **68**(1):122-
189 137.

190 IYENGAR, N. S. AND SUDHARSHAN, P., 1982, A method of classifying regions from
191 multivariate data, *Economic and Political Weekly, Special Article* : 2048-1052.

192 RAJU, K. V., DESHPANDE, R. S. AND SATYASIBA BEDAMATTA, 2017, Vulnerability to
193 Climate Change: A Sub-regional Analysis of Socio-economic and Agriculture Sectors in
194 Karnataka, India, *Journal of Development Policy and Practice*, **2**(1):24–55.

195

196

197

198 **Tables**

199 **Table 1: Index of climate vulnerability across the various districts of Karnataka**

Sl. No	Districts	Annual rainfall	S-W monsoon	Max Temp	Min Temp	Avg Temp	Index total
1	KALABURAGI	0.177	0.174	0.102	0.169	0.125	0.747
2	KOLAR	0.170	0.183	0.189	0.107	0.073	0.720
3	BIDAR	0.173	0.169	0.120	0.189	0.069	0.720
4	RAICHUR	0.183	0.183	0.082	0.127	0.138	0.712
5	YADGIR	0.181	0.177	0.088	0.130	0.135	0.711
6	VIJAYAPURA	0.185	0.182	0.087	0.147	0.100	0.701
7	RAMANAGARA	0.163	0.178	0.165	0.085	0.103	0.693
8	BALLARI	0.186	0.187	0.100	0.054	0.154	0.681
9	KOPPALA	0.183	0.184	0.063	0.096	0.141	0.667
10	BAGALKOTE	0.188	0.185	0.078	0.109	0.095	0.656
11	DHARWAD	0.181	0.179	0.118	0.094	0.070	0.643
12	DAVANAGERE	0.178	0.179	0.066	0.046	0.125	0.593
13	GADAG	0.189	0.186	0.061	0.075	0.082	0.593
14	CHITRADURGA	0.183	0.185	0.060	0.044	0.119	0.591
15	CHIKKABALLAPURA	0.176	0.183	0.055	0.077	0.094	0.585
16	BELAGAVI	0.174	0.166	0.058	0.103	0.083	0.584
17	TUMAKURU	0.176	0.181	0.044	0.055	0.105	0.561
18	HAVERI	0.179	0.174	0.072	0.051	0.082	0.559
19	MANDYA	0.172	0.185	0.038	0.026	0.116	0.537
20	MYSURU	0.173	0.182	0.023	0.036	0.101	0.514
21	CHAMARAJANAGARA	0.172	0.189	0.000	0.028	0.086	0.475
22	UTTARA KANNADA	0.078	0.074	0.139	0.077	0.106	0.474
23	BENGALURU RURAL	0.164	0.175	0.023	0.052	0.056	0.470
24	BENGALURU URBAN	0.157	0.168	0.015	0.038	0.055	0.431
25	HASSAN	0.152	0.149	0.024	0.043	0.052	0.421
26	CHIKKAMAGALURU	0.128	0.126	0.076	0.039	0.042	0.412
27	SHIVAMOGGA	0.103	0.082	0.053	0.058	0.076	0.372
28	DAKSHINA KANNADA	0.031	0.030	0.045	0.000	0.189	0.294
29	KODAGU	0.080	0.082	0.094	0.022	0.000	0.278
30	UDUPI	0.000	0.000	0.024	0.022	0.170	0.215

200

201

202

203

204

205 **Table 2: Agricultural vulnerability index across the districts of Karnataka**

District	Geographical area(Ha)	Forest area(% to GA)	Total food crops(% to GCA)	Net sown area(% to GA)	Livestock pon (No. per Ha of GCA)
Kodagu	0.008	0.032	0.055	0.027	0.065
Bidar	0.013	0.051	0.026	0.016	0.063
Kolar	0.007	0.051	0.012	0.027	0.057
Chitradurga	0.026	0.048	0.031	0.024	0.059
Koppal	0.014	0.051	0.018	0.010	0.062
Hassan	0.019	0.048	0.022	0.021	0.061
Gadag	0.010	0.049	0.020	0.001	0.064
Dakshin Kannada	0.011	0.036	0.013	0.034	0.063
Dharwad	0.008	0.049	0.023	0.003	0.064
Haveri	0.011	0.048	0.021	0.006	0.062
Chikballapura	0.008	0.046	0.016	0.024	0.058
Bengaluru Rural	0.000	0.051	0.026	0.022	0.061
Kalaburagi	0.036	0.052	0.006	0.009	0.064
Raichur	0.026	0.053	0.014	0.018	0.061
Tumkuru	0.035	0.051	0.037	0.024	0.059
Mysuru	0.017	0.047	0.021	0.016	0.062
Chamarajanagara	0.014	0.021	0.017	0.034	0.061
Yadgir	0.013	0.050	0.028	0.014	0.061
Ramanagara	0.005	0.041	0.019	0.024	0.060
Chikkamagaluru	0.021	0.035	0.032	0.027	0.063
Vijayapura	0.034	0.054	0.003	0.000	0.064
Bagalkot	0.018	0.046	0.005	0.010	0.060
Uttar Kannada	0.033	0.000	0.004	0.044	0.059
Udupi	0.006	0.035	0.015	0.035	0.061
Mandya	0.011	0.051	0.010	0.026	0.058
Belagavi	0.046	0.045	0.014	0.016	0.065
Bellari	0.026	0.046	0.019	0.020	0.057
Bengaluru Urban	0.000	0.053	0.017	0.041	0.000
Shivamogga	0.026	0.032	0.000	0.035	0.060
Davanagere	0.015	0.044	0.005	0.013	0.062

206
207
208
209
210
211
212
213

214 Contd....

Irrigated area(% to GCA)	Cropping intensity (%)	Productivity	Per capita income	Depth of groundwater (mbgl)	Index total
0.049	0.045	0.443	0.046	0.018	0.787
0.041	0.034	0.438	0.051	0.030	0.761
0.038	0.056	0.442	0.045	0.006	0.741
0.032	0.037	0.410	0.048	0.017	0.732
0.027	0.040	0.426	0.050	0.012	0.708
0.032	0.037	0.378	0.042	0.038	0.697
0.035	0.029	0.402	0.046	0.038	0.694
0.012	0.041	0.449	0.014	0.011	0.684
0.041	0.000	0.429	0.041	0.024	0.683
0.028	0.039	0.394	0.048	0.020	0.677
0.029	0.048	0.381	0.046	0.012	0.669
0.034	0.057	0.358	0.037	0.020	0.667
0.040	0.041	0.332	0.051	0.033	0.663
0.023	0.039	0.379	0.049	0.000	0.663
0.025	0.044	0.345	0.040	0.003	0.662
0.026	0.024	0.371	0.045	0.030	0.661
0.019	0.037	0.392	0.044	0.017	0.657
0.023	0.034	0.379	0.051	0.003	0.655
0.034	0.053	0.352	0.039	0.028	0.654
0.036	0.035	0.353	0.029	0.021	0.651
0.025	0.053	0.348	0.050	0.018	0.649
0.014	0.036	0.378	0.041	0.037	0.646
0.024	0.050	0.364	0.043	0.021	0.644
0.027	0.045	0.386	0.024	0.007	0.641
0.005	0.033	0.356	0.039	0.039	0.628
0.011	0.027	0.325	0.049	0.025	0.623
0.014	0.032	0.297	0.040	0.013	0.564
0.029	0.049	0.361	0.000	0.003	0.552
0.000	0.047	0.266	0.036	0.025	0.527
0.012	0.049	0.267	0.048	0.010	0.524

215
216
217
218
219
220
221
222

Table 3: Demographic vulnerability index across the districts of Karnataka

District	Density of male population	Density of female population	Literacy rate of male (%)	Literacy rate of female (%)	Index total
Bengaluru Urban	0.281	0.281	0.015	0.001	0.579
Yadgir	0.006	0.006	0.225	0.212	0.449
Raichur	0.006	0.006	0.165	0.176	0.353
Chamarajanagara	0.003	0.003	0.184	0.145	0.335
Kalaburagi	0.007	0.007	0.137	0.144	0.294
Bellari	0.010	0.010	0.120	0.129	0.270
Vijayapua	0.005	0.005	0.116	0.136	0.262
Ramanagara	0.011	0.012	0.119	0.112	0.255
Koppal	0.007	0.008	0.106	0.132	0.253
Bagalkot	0.010	0.010	0.101	0.128	0.249
Mandya	0.015	0.016	0.108	0.107	0.246
Chikkaballapura	0.011	0.011	0.112	0.112	0.245
Bidar	0.012	0.012	0.102	0.112	0.238
Mysuru	0.022	0.023	0.107	0.085	0.237
Belagavi	0.014	0.015	0.080	0.097	0.206
Kolar	0.016	0.017	0.083	0.086	0.202
Chitradurga	0.004	0.004	0.086	0.091	0.184
Davanagere	0.013	0.013	0.078	0.075	0.179
Tumakuru	0.008	0.008	0.075	0.083	0.174
Gadag	0.006	0.006	0.062	0.093	0.167
Bengaluru Rural	0.020	0.020	0.061	0.067	0.167
Hassan	0.008	0.009	0.069	0.077	0.163
Haveri	0.013	0.013	0.067	0.068	0.160
Dharwad	0.019	0.020	0.049	0.053	0.142
Chikkamagaluru	0.001	0.002	0.056	0.054	0.114
Shivamogga	0.005	0.005	0.052	0.046	0.107
Kodagu	0.000	0.000	0.043	0.030	0.073
Uttara Kannada	0.000	0.000	0.026	0.028	0.055
Udupi	0.011	0.014	0.013	0.013	0.051
Dakshina Kannada	0.018	0.021	0.000	0.000	0.039

228 **Table 4: Composite index of vulnerability**

Sl. No	Districts	Composite index	Sl. No	Districts	Composite index
1	BIDAR	0.677	16	CHAMARAJANAGAR	0.579
2	KOLAR	0.658	17	MYSURU	0.574
3	YADGIR	0.638	18	TUMKUR	0.573
4	KOPPAL	0.636	19	HASSAN	0.571
5	RAICHUR	0.628	20	BENGALURU RURAL	0.558
6	CHITRADURGA	0.628	21	MANDYA	0.557
7	KALABURAGI	0.625	22	BELAGAVI	0.555
8	RAMANAGARA	0.604	23	BALLARI	0.543
9	VIJAYAPURA	0.602	24	BENGALURU URBAN	0.538
10	GADAG	0.599	25	CHIKKKAMAGALURU	0.531
11	DHARWAD	0.596	26	UTTARA KANNADA	0.530
12	KODAGU	0.594	27	DAKSHINA KANNADA	0.528
13	CHIKBALLAPUR	0.593	28	UDUPI	0.486
14	BAGALKOT	0.590	29	DAVANGERE	0.486
15	HAVERI	0.580	30	SHIVAMOGGA	0.440
Average=0.577					

229

230