Original Research Article

1 2 3

Future Impact of Climate Change on the Yield of Cocoa in Ondo State, Nigeria

This study was carried out to determine the trend of cocoa yield and climatic variables and

4 5

7

Abstract

also to assessment of the impacts of climate change on the future yield of cocoa in Ondo 8 State, Nigeria. Annual trend statistics for cocoa yield and climatic variables were analyzed 9 for the state using Mann-Kendall test for trend and Sen's slope estimates. Downscaled 10 data from six Global Circulation Models (GCMs) were used to examine the impact of 11 12 climate change on the future yield of cocoa in the study area. The results of trends analysis in Ondo State showed that yield decreased monotonically at the rate of 492.18 13 tonnes/yr (P<0.05). An increased significant trend was established in annual rainfall trend. 14 While The Maximum temperature, minimum temperature, and mean temperature all 15 increased at the rate of 0.02 °C/yr (P<0.001). The ensemble of all the GCMs projected a 16 mid-term future decrease of about 9,334 tonnes/yr by 2050 and a long-term future 17 decrease of 13,504 tonnes/yr of cocoa by 2100. The economic implication of these is that, 18 if the projected change in the yield of cocoa as predicted by the ensemble of all the GCMs 19 20 should hold for the future, it means that Ondo state may experience a loss of about \$22,470,018.22 and \$32,308,584.32 by the year 2050 and 2100 respectively according to 21 the present price of the commodity in the world market. This research should be extended 22 23 to other cocoa producing areas in Nigeria.

Comment [M1]: What is the rate of increase of annual rainfall

Keyword: Climate change; cocoa yield; climate variable; Mann-Kendall trend; future yield

1. Introduction

One of the most important cash crops contributing to the gross domestic product (GDP) of Nigeria is cocoa (*Theobroma cacao*) (Oyekale *et al*, 2009). It has contributed greatly to the economic development and social wellbeing of the people in the cocoaproducing areas and also boosted the financial status of the farmers after oil. The beans derived from cocoa is used in the production of chocolate products, biscuits, cocoa bread, cream, soap, livestock feeds, cocoa powder and amongst others (Hamsat *et al.*, 2003; Olubamiwa *et al.*, 2000).

In the time past, Nigeria used to be the second largest producer of cocoa in the world and Ondo state was also the largest producing state but the production dwindled due to discovery of oil, change in weather and management practices, and Ceurrently, Nigeria is the fourth producer of cocoa in the World and due to some limiting factors, the production of cocoa declined drastically because of change in weather and climate change, management practices, oil exploration, etc. (ICCO, 2008). Ayanlaja (2000) reported that cocoa production declined from 310,000 tonnes/yr despite increase insecticide application, land area, and introduction of a high yielding variety of cocoa in the country. Weather and climate change over the years has greatly affected cocoa production which is a major cash crop in Nigeria where Ondo state the worst hit of the menace.

Comment [M2]: where is the eference

The vVariation in the two climatic variables: Rrainfall and temperature were discovered to have much influence on the sprouting, production, and growth of cocoa trees (Anim-Kwapong and Frimpong, 2005). However, most of the developing countries are already experiencing low yield of the crop, as a result, extreme weather and climate change (Odjugo, 2010). Extreme weather is a situation best described as extreme in terms of historical distribution, severe or unfavorable weather (ICCO, 2003). Climate change was reported to have played a vital role in the alteration, development of cocoa pests and pathogens thereby shifting their interactions (Oyekale et al., 2009). This, in turn, leads to lower yield, which brings about low yield, which brings about reduced income and livelihood for the farmers. Cocoa production is highly sensitive to change in rainfall, the intensity of sunshine, temperature, water supply, soil condition due to evapotranspiration effects cocoa production (Anim-Kwapong and Frimpong, 2005). Climate change has been reported to be one of the most serious environmental threats affecting humans and their crops in the world today (Enete and Amusa, 2010). It also has a great effect on agricultural production.

Comment [M3]: REference not listed

Unfortunately, the recent trends pattern of rainfall had—either been excess increasing thereby—leading to the infestation of black pod disease which also leads to losses in cocoa yield. Insufficient rainfall also leads to seed mortality, drought and bush burning. This study was therefore carried out gives us the opportunity to examine the trends and impact of climate change on cocoa yield in Ondo state.

Therefore, the aim of this study is to evaluate the trends in historical cocoa yield, climatical variables—and determine—the impacts of climate change on the future yield of cocoa in

Ondo state, Nigeria.

Formatted: Indent: First line: 0"

2. Methodology

70

71

81

83

84

85

86

87

88

89

2.1 Study Area

72 Ondo state is located within the rainforest agro-climatic zone of Southwest Nigeria. It lies between latitudes 5°45' and 7° 52'N and longitudes 4° 20' and 6° 5' E. The major 73 74 occupation of the people is agriculture, which provides income and employment for about 75 70% of the total population. The major arable crops cultivated include: yam, rice, cassava, 76 tomatoes, maize, etc and some tree crops cultivated include: cocoa, coffee, oil palm and 77 timber (OSMARD, 2004). Ondo state consists of 18 local government areas producing 78 about 45% to 65% of the total cocoa production figures in Nigeria. OSMARD (2004) 79 reported that 9 local government areas (LGAs) are producing about 95% of the total cocoa 80 production in the state which include: Akure-North, Akure -South, Ondo- East, Ese Odo,

Odigbo, Ile Oluji / Okeigbo, Ondo -West, Owo, and Ilaje.

82 2.2 Data Source

The climatic data—used for this study—was (rainfall, Maximum temperature, Minimum temperature and Mean temperature) were extracted between from 1976 andto 2014. These data were retrieved from the Climate Research Unit (CRU) dataset (www.cru.uea.ac.uk). The cocoa yield data from 1976 to 2014 was obtained from Ondo State Ministry of Agriculture and Natural Resources (between 1976 and 2014 and also from Food and Agricultural Organization statistics (FAOSTAT, www.faostat.org).

2.3 Data Analysis

To evaluate the trend in cocoa yield and meteorological parameters in the study area, MAKESENS (Mann-Kendall test for trend and Sen's slope estimates), An Excel template which was developed for detecting and estimating trends in the time series was used. The Mann-Kendall test statistic S is given by Salmi *et al.* (2002) as:

94
$$S = \sum_{k=1}^{n-1} \sum_{j=k+1}^{n} \operatorname{sgn}(x_{j} - x_{k})$$
 (1)

Wherein is the length of the time series $x_i...x_n$, and sgn(.) is a sign function, x_j and x_k are valued in years j and k, respectively. The expected value of S equals zero for series without trend and the variance is computed as:

99
$$\sigma^{2}(S) = \frac{1}{18} \left[n(n-1)(2n+5) - \sum_{p=1}^{q} t_{p}(t_{p}-1)(2t_{p}+5) \right]$$

Where q is the number of tied groups and t_p is the number of data values in p^{th} group. The test statistic Z is then given as:

$$Z = \begin{cases} \frac{S-1}{\sqrt{\sigma^2(S)}} & \text{if } S > 0\\ 0 & \text{if } S = 0\\ \frac{S+1}{\sqrt{\sigma^2(S)}} & \text{if } S < 0 \end{cases}$$
(3)

No assumptions as to the underlying distribution of the data are very significant as a non-parametric test, The Z statistic was then used to test the null hypothesis, H_o that the data are randomly ordered in time, against the alternative hypothesis, H_I , where there is an increasing or decreasing monotonic trend. A positive (negative) value of Z indicates an upward (downward) monotone trend. H_o will be rejected at a particular level of significance if the absolute value of Z is greater than $Z_{I-\alpha/2}$, where $Z_{I-\alpha/2}$ is obtained from the standard normal cumulative distribution tables. Hobbins et al. (2001) noted that the Mann-Kendall test is non-dimensional and does not quantify the scale or the magnitude of the trend but the direction of the trend. To estimate the true slope of an existing trend the Sen's non-parametric method will be used (Salmi *et al.*, 2002).

3. Results and Discussion

3.1 Descriptive trends in annual cocoa yield and climatic variables

The summary statistics of the Mann-Kendall monotonic trend statistics and nonparametric sen_"s slope estimate test for annual trends in cocoa yield and climatic variables in Ondo State is presented in Table 1. The results of the analysis of trends showed that cocoa yield decreased monotonically at the rate of 492.18 tonnes/yr (P<0.05), which agrees with Oguntunde *et al.*(2014), that noticed attributed a decreasing trend in cocoa yield in the study area_Ondo state which was attributed to variations in weather elements. No significant correlation was established in annual rainfall. While there was a positive significant trend in Mmaximum- temperature, Mminimum- temperature and mean temperature all at the rate of 0.02 °C/yr (P<0.001). Similarly, the statistical trend of annual yields of cocoa and the climatic variable is presented in Figures 1 to 5. Cocoa Yield

Comment [M4]: Reference not listed

Comment [M5]: Reference not listed

showed a declining temporal trend while Rainfall, <u>Mmaximum</u> temperature, <u>Mminimum</u> temperature and mean temperature showed a positive statistical trend. This may be due <u>to</u> variations in the amount of rainfall in the study area which is in line with the findings of Thompson, (2013) who carried out a study on <u>the</u> climate change and <u>the</u> cocoa production in Ekiti and Ondo States of Nigeria: A <u>using a cointegration</u> analysis. He reported that the availability of rainfall will have much effect on cocoa yield over time.

There was also an increasing positive statistical trend for climatic variables during the time under study. This result also confirms the empirical study carried out by Oyekale et al. (2009) who reported that the reduction in the cocoa yield was as a result of excessive rainfall which was deduced in the time under study. These also reduce the quality of cocoa as a result of the climatic failure. In terms of correlation, rainfall was not significantly correlated with the yield of cocoa; Ttemperature maximum, Ttemperature minimum, and Ttemperature mean were the only variables that are significantly correlated with cocoa yield in the study area.

3.2 Development of climate-yield regression Models

The correlation between cocoa and climatic variables has given us an understanding of the time characteristics of each variable. Therefore, to establish a relationship between cocoa yield and climatic variables, the variables which were identified to have a significant relationship with the yield using multiple linear regression analysis were regressed with cocoa yield. The summary of stepwise regression between cocoa yield and climatic variables using multiple linear regression for the four states are presented in Table 2.

In the study area, the model I <u>of</u> the correlation coefficient (R) = 0.52 showing that the regressor (TY_{12}) in the model I accounted for 52% in the variability of cocoa yield. Model II also have R = 0.68 which implies that the regressors (TY_{12} and R6) are responsible for 68% variability in cocoa yield in Ondo State. Looking at Model III, where R = 0.74, this implies that the regressors (TY_{12} , R₆, and TX_5) are responsible for 74% of the variability in cocoa yield. Examining Model IV, with R = 0.78, this simply means that the regressors in model IV (TY_{12} , R₆, TX_5 , and TZ_3) accounted for 78% in the variability in cocoa yield. Model IV from the stepwise regression analysis was also selected in order to predict the future yield of cocoa by 78% assurance based on RCP 4.5 emission scenario.

The Global Circulation Models (GCMs) for the projection of the future climate data by the IPCC based on the RCP 4.5 emission scenario were used for this study. This includes CCCMA, ICHEC, MIROC, NCC, NPI, and NOAA. The present-day cocoa yield estimation for the study area present daytime series (1976 - 2005) Midterm (2021-2050) and long term (2071-2100) for the six GCMs based on RCP 4.5 are presented in Figures 6 to Figure 8.

3.3 Impact of climate change on the future yield of cocoa in Ondo state

Figure 9 shows the impact of climate change on the yield of cocoa in Ondo state. Looking at The CCCMA model, shows a mid-term decrease of 7,413 tonnes/yr of cocoa was projected for for the study area by the year 2050, and for At the long term, a decrease of 10,383 tonnes/yr of cocoa was projected for the study area by 2100. A decrease of 7,992 tonnes/yr cocoa was projected by ICHEC model for the study area for the mid-term by the year 2050. For At the long-term projection by ICHEC, a decrease of 11,852 tonnes/yr was projected by the year 2100 for in the study area. Considering MIROC, also in Figure 9, 15,960 tonnes/yr decrease yield of cocoa was projected by the year 2050 while a decrease of 28,146 tonnes/yr yield of cocoa was projected for in the study area by the year 2100.

For NCC and at the same emission scenario, there will <u>be</u> a mid-term future decrease of 8,926 tonnes/yr of cocoa by the year 2050 <u>forin</u> the study area. <u>For At</u> the long-term, a decrease of 8,162 tonnes/yr of cocoa was projected by the year 2100. With reference to MPI, there was a decrease of 6,335 tonnes/yr of cocoa for the mid-term future projection for the study area and decrease of 6,395 tonnes/yr in the long-term future was projected for the study area by the year 2100.

______For the NOAA model, there was a decrease in both mid-term and long-term future with 9,379 tonnes/yr and 16084 tonnes/yr of cocoa by the year 2050 and 2100 respectively forin the study area. The ensemble of all the GCMs projected a mid-term

future reduction of 9,334 tonnes/yr by 2050 and a long-term future decrease of 13,504 tonnes/yr of cocoa by 2100.

The study of the impact of climate change on the future yield of cocoa both by the midterm (2050) and long term (2100) in the study area cannot be overemphasized being the highest cocoa producing state. From the study, climate change will have a negative impact experienced in the study area. The variation in the future projected yield of cocoa may be due to variability in rainfall distribution across the study area by the year 2050 and 2100. This agrees with Oluyole (2010); Edet *et al.* (2018); Amos and Thompson (2015) that variability in rainfall has much influence on the cocoa yield. Thompson (2013) established that the yield of cocoa is mostly affected by rainfall variability in the long run, that is, the yield of cocoa is highly susceptible to drought and excess rainfall.

Anim-Kwapong and Frimpong (2005) reported that cocoa is highly sensitive to rainfall and water application. Also, yearly variations in the yield of cocoa were affected by more by rainfall than any other factors in Nigeria (Ajewole and Iyanda, 2010).

3.4 The economic implication of the impact of climate on cocoa yield

The projected change in the future yield of cocoa using RCP 4.5 future climate scenario in the future for all the GCM, showed that a loss of 9,334 tonnes/yr of cocoa was projected by the year 2050 and a loss of 13,504 tonnes/yr of cocoa was also projected by the year 2100 Now, if the projected change in the yield of cocoa as predicted by the ensemble of all the GCMs should hold for the future, it means that Ondo state may

Comment [M6]: Reference not listed

Comment [M7]: Reference not listed

Comment [M8]: Reference not listed

Formatted: Indent: First line: 0"

Comment [M9]: REference not listed

208 experience a loss of about \$22,470,018.22 and \$32,308,584.32 by the year 2050 and 2100 209 respectively. 210 211 4. Summary and Conclusion From the trend analysis, the yield decreased monotonically at a rate of 492.18 tonnes/yr 212 (P<0.05). The increasing trend was established in annual rainfall trend. And there was a 213 214 positive significant trend in maximum temperature, minimum temperature and mean temperature all at the rate of 0.02 °C/yr (P<0.001). The impact of climate change on the 215 yield of cocoa in the study areas, there was a projected yield decrease of 9,334 and 9,379 216 tonnes/yr by the year 2050 and 2100 respectively 217 218 219 References Anim-Kwapong, G. J. and Frimpong, E. B. (2005). The vulnerability of Agriculture to 220 Climate - Change impact of climate on Cocoa production. Cocoa Research 221 Institute, New TafoAkim, Ghana. 222 223 Ayanlaja, S. A. (2000). The development of an alley cropping system: a viable agroforestry technology for sustained cocoa production in Nigeria. Cocoa 224 growers' Bulletin', No. 52. Pp. 9-18 225 Enete, A. A. and T. A. Amusa, (2010). Challenges of Agricultural Adaptation to Climate 226 Change in Nigeria: a Synthesis from the Literature », Field Actions Science 227

Comment [M10]: The sentence is floating. Complete the sentence

| 228 | Reports [Online], Vol. 4 2010, Online since 20 December 2010, connection on 30 | | |
|---|--|--|--|
| 229 | September 2016.URL: http://factsreports.revues.org/678 | | |
| 230 | Hamzat, R. A. Umana, T. E. and Babayemi, O. J. (2003). Potentials of cocoa Husk pulp as | | |
| 231 | a sole Feed ingredient for African giant land Snails (Archachatina marginata). P. | | |
| 232 | 151. In the book of abstracts of the 14 th International Cocoa Conference | | |
| 233 | Organized by Cocoa producer's Alliance (COPAL) with the Government o | | |
| 234 | Ghana, Accra. 13 th – 18 th October. | | |
| 235 | International Cocoa Organization (ICCO) (2005). World Cocoa production, 2005. | | |
| 236 | Retrieved from http://www.icco.org. | | |
| 237 | ICCO (International Cocoa Organization) (2008). Annual Report 2008/2009. London: | | |
| | | | |
| 238 | ICCO. Accessed June, 011.www.icco.org/pdf/Annual | | |
| 238 239 | ICCO. Accessed June, 011.www.icco.org/pdf/Annual _report/AnnualReport2008/2009. | | |
| | | | |
| 239 | _report/AnnualReport2008/2009. | | |
| 239 240 | _report/AnnualReport2008/2009. Odjugo, P. A. O. (2010). General Overview of Climate Change Impacts in Nigeria. | | |
| 239240241 | _report/AnnualReport2008/2009. Odjugo, P. A. O. (2010). General Overview of Climate Change Impacts in Nigeria. Journal Hum Ecol, 29(1): 47-55. | | |
| 239240241242 | _report/AnnualReport2008/2009. Odjugo, P. A. O. (2010). General Overview of Climate Change Impacts in Nigeria. Journal Hum Ecol, 29(1): 47-55. Olubamiwa, O., odewumi, W. O. Longe, O. G. and Hamzat, R. a. (2000). Practical | | |
| 239240241242243 | _report/AnnualReport2008/2009. Odjugo, P. A. O. (2010). General Overview of Climate Change Impacts in Nigeria. Journal Hum Ecol, 29(1): 47-55. Olubamiwa, O., odewumi, W. O. Longe, O. G. and Hamzat, R. a. (2000). Practical inclusion of cocoa bean shell in poultry feed A Preliminary Report. Pp. 981-986, | | |
| 239240241242243244 | _report/AnnualReport2008/2009. Odjugo, P. A. O. (2010). General Overview of Climate Change Impacts in Nigeria. Journal Hum Ecol, 29(1): 47-55. Olubamiwa, O., odewumi, W. O. Longe, O. G. and Hamzat, R. a. (2000). Practical inclusion of cocoa bean shell in poultry feed A Preliminary Report. Pp. 981-986, in Proceedings of the 13 th International Cocoa Research Conference. Organized by | | |

Comment [M11]: the reference not cited in the text

| 248 | Oyekale, A. S., Bolaji, M. B., and low a, O. W. (2009). The effects of climate change on |
|-----|--|
| 249 | cocoa production and vulnerability assessment in Nigeria. Agricultural Journal, |
| 250 | 4(2), 77-85. |
| 251 | Salmi, T., Maatta, A., Anttila, P., Ruoho-AirolaAmnell, T.,(2002). Detecting trends of |
| 252 | annual values of atmospheric pollutants by the Mann-Kendall test and Sen's slope |
| 253 | estimates. Publications on air quality, No 31. Helsinki, Finland |
| 254 | Thompson, (2013). The Effect Climate change on Cocoa Production in Ekiti and Ondo |
| 255 | States of Nigeria: A co-integration analysis. Journal of Sustainable Technology, |
| 256 | 4(2) pp 66-77 |
| 257 | |

Table 1: Trends results of Annual Yield and climatic variables for Ondo State

260

| Time series | Test Z | Significance | Slope |
|------------------|--------|--------------|---------|
| Yield | -2.03 | * | -492.18 |
| Rainfall | 0.53 | | 1.88 |
| Max. Temperature | 4.15 | *** | 0.02 |
| Min. Temperature | 4.09 | *** | 0.02 |
| Mean Temperature | 4.61 | *** | 0.02 |

*** Significant at 0.001, **significant at 0.01, * significant at 0.05, * significant at 0.1




Figure 1: Annual trends in yield of cocoa between 1976 and 2014 in Ondo State




Figure 2: Annual trends in rainfall between 1976 and 2014 in Ondo State




Figure 3: Annual Trends in Maximum. Temperature between 1976 and 2014 in Ondo State




Figure 4: Annual trends in Minimur temperature between 1976 and 2014 in Ondo

272 State

271




Figure 5: Annual trends in mean temperature between 1976 and 2014 in Ondo State

Table 2: Summary of the stepwise regression between cocoa yield and the climatic variables

| Models | States | Regression Model | R |
|--------|--------|--|------|
| i. | Ondo | $Y_c = \beta_0 + \beta_1 T Y_{12} + e_i$ | 0.52 |
| ii. | Ondo | $Y_c = \beta_0 + \beta_1 T Y_{12} + \beta_2 R_6 + e_i$ | 0.68 |
| iii. | Ondo | $Y_c = \beta_0 + \beta_1 T Y_{12} + \beta_2 R_6 + \beta_3 T X_5 + e_i$ | 0.74 |
| iv. | Ondo | $Y_c = \beta_0 + \beta_1 T Y_{12} + \beta_2 R_6 + \beta_3 T X_5 + \beta_3 T Z_3 + e_i$ | 0.78 |
| 17. | Ondo | $r_c = \rho_0 + \rho_1 + \rho_2 + \rho_3 $ | 0.70 |

Where R = Correlation coefficient, Yc = Yield, e_i = error term, $\beta_0 - \beta_3$ are constant. R_1 - R_{12} (Rainfall of January – December), R13 (Annual Rainfall); TY_1 - TY_{12} (Max Temperature of January – December), TY_{13} (Annual max temperature); TX_1 - TX_{12} (Min. Temperature of January – December), TX_{13} (Annual max temperature); TZ_1 - TZ_{12} (Mean Temperature of January – December), TZ_{13} (Annual mean temperature).




Figure 6: Present Day (1976 – 2005) cocoa yield in Ondo State




Figure 7: Projected Midterm (2021 – 2050) for different GCMs output based on RCP 4.5 in
Ondo State




Figure 8: Projected Long term future (2071-2100) yield for Six GCMs output based on

RCP 4.5




Figure 9: Change in yield of cocoa for the mid-term (MT; 2021-2050 in green) and long-term (LT; 2071 -2100) in blue under RCP 4.5 emission scenario

Table 3: Descriptions of acronyms used for the study

| 300 | Acronyms | Meaning of the Acronyms |
|-----|-----------------|--|
| 301 | GCM | General Climate model / Global circulation model |
| 302 | RCP | Representative Concentration Pathways |
| 303 | IPCC | Intergovernmental Panel on Climate Change |
| 304 | IITA | International Institute of Tropical Agriculture |
| 305 | CRIN | Cocoa Research Institute |
| 306 | FAO | Food and Agricultural Organization |
| 307 | R | Correlation Coefficient |
| 308 | GHG | Green House Gases |
| 309 | CO ₂ | Carbon dioxide |
| 310 | NPC | National Population Commission |
| 311 | CRU | Climate Research Unit |
| 312 | Tmax | Maximum Temperature |
| 313 | Tmin | Min Temperature |
| 314 | Tmean | Mean Temperature |
| 315 | | |