

Original Research Article

Evaluation of carbon stock across different forest physiognomy in a tropical rainforest ecosystem at Obafemi Awolowo University Ile-Ife, Nigeria

Abstract

This study investigated carbon stock in above-ground biomass across different physiognomies in Obafemi Awolowo University tropical rainforest ecosystem. This was with a view of increasing the understanding of carbon cycle in tropical rainforest in Nigeria.

Two 20 m x 20 m plots were marked out in the secondary forest, *Tectona grandis* and Riparian vegetations. Total enumeration was carried out for the living tree, the Diameter at Breast Height (DBH) of trees ≥ 10 cm were measured at 1.3 m above the ground and height was also determined using a ranging pole and Haga altimeter.

Aboveground carbon stocks in standing trees ranged from 218.24 to 318.92 C t ha⁻¹ with the highest value in *Tectona grandis* plantation. Trees with DBH size class 11-20 cm contributed more to Carbon stock in secondary forest and *Tectona grandis* plantation, while size class ≥ 60 cm contributed more in the riparian vegetation. *Tectona grandis* plantation proved to be better in mitigating carbon in our environment and this result will enhance better estimates of local and regional carbon stock which is crucial to addressing the problems of climate change.

Keywords: Allometric, Atmosphere, Climate, Human, Plantation, Sequestration

Introduction

Tropical rainforest and plantation ecosystems sequester carbon in terrestrial ecosystems and therefore serve as an important natural brake on climate change (Gibbs *et al.* 2007). These ecosystems are unique environmental resources that provide numerous global benefits and play crucial role with respect to global carbon pools and fluxes as they store about half of the world's biomass (Brown and Lugo, 1992). It has been previously reported that they represent important pools of biological, ecological and economic resources (Sheikh *et al.* 2012), which greatly influence the lives of other organisms as well as human societies (Komiya *et al.* 2008). The tropical forest and plantation ecosystems are long-lived dynamic systems that are involved in climate regulation (Egbe and Tabot, 2011); as well as prominent sites for the study of climate change in terms of total net carbon emission and global storage capacity (Terakunpisut *et al.* 2007).

The main carbon pools in tropical forest and plantation ecosystems are the living biomass of trees, understorey vegetations, mass of litters, woody debris and soil organic matter (Ludang and Jaya, 2007). The carbon stored in the aboveground living biomass of trees is typically the largest and the most directly impacted upon by human disturbances (Gibbs *et al.* 2007). Stable tropical forest and plantation ecosystems with less disturbances are important as carbon sinks and are currently sequestering carbon dioxide (CO₂) from the atmosphere which are critical to future climate stabilization (Stephens *et al.* 2007) and this can be strengthened by increasing the density of vegetations cover in currently vegetated areas or increasing the areas covered by vegetations (Karjalainen *et al.* 2002).

Forest and plantation ecosystems management practices can play a significant role in climate change mitigation by sequestering carbon through photosynthesis (Strassburg *et al.* 2009). Knowledge of the aboveground living biomass density is useful in determining the amount of carbon stored through photosynthesis in the forest stands. Forest also releases carbon to the atmosphere through plant respiration and organic material decomposition, although the loss of carbon into the atmosphere is usually less than the gain (Fonseca *et al.* 2011).

The issue of aboveground biomass and carbon stock has received tremendous attention across the world; however, little information is available in Nigeria. This study is focusing on carbon sequestration specifically in terms of aboveground biomass and carbon stock. The estimates of carbon stock are important for scientific and management issues such as forest productivity and nutrient cycling.

In addition, aboveground biomass is a key variable in the annual and long term changes in the global terrestrial carbon cycle and other earth system interactions. Hence, a study on evaluation of carbon stock in the aboveground biomass of tropical rainforest and plantation ecosystem was conducted in Obafemi Awolowo University estate, Ile-Ife, with the aim of providing information on carbon stock across different forest vegetations that is critical to better understanding of the issues of global climate change. The specific objective of this study was to estimate carbon stock in aboveground biomass across different vegetations (secondary forest, *Tectona grandis* plantation and Riparian vegetation) based on allometric models.

Materials And Methods

Study area

The study was conducted at the Obafemi Awolowo University, Ile-Ife, Osun state, Nigeria. Ile-Ife is located on Latitude N 07° 31' and Longitude E 04° 30' and the elevation of Ife ranges from 215 m to 457 m above sea level (Hall, 1969). The study sites lies between Latitude N 07° 032' and Longitude E 04° 031' while the elevation ranges from 243 m to 274 m above the sea level. The climate of the area is a tropical type with two prominent seasons, the rainy and the dry season. The dry season is short, usually lasting 4 months from November to March and the longer rainy season prevails during the remaining months. The annual rainfall average 1400 mm yr⁻¹ (Oke and Isichei, 1997) and it showed two peaks, one in July and the other in September, the mean annual temperature range from 27° C to 34° C (Oke and Isichei, 1997).

The soil of the area is derived from material of old basement complex which is made up of granitic metamorphosed sedimentary rock (Hall, 1969). Five major soil types have been recognized in this area: inselberg soils, Hill creep soils, and sedimentary non-skeletal soils, drift soils, alluvial deposits (Hall, 1969). The soil has been classified as lixisols and utisols (FAO/UNESCO, 1974). The original vegetation of Ile-Ife is lowland rainforest as climax vegetation (Keay, 1959). White (1983) described the vegetation as the Guinea-Congolian drier forest type. Most of the original lowland rain forests have been massively destroyed leaving remnant of secondary forest scattered around. Tree crops plantations like *Theobroma cacao*, *Cola nitida*, *Tectona grandis*, and *Elaeis guineensis* are now common around the area.

Sampling procedure

Two samples plots, each of 20 × 20 m were marked out within the secondary forest, *Tectona grandis* plantation and riparian vegetation in the Obafemi Awolowo University community. The secondary forest is 29 years old having been last disturbed by ground fire that engulfed the forest in 1983. It is located within the Biological Garden and lies within latitude 07° 32' 23.11"N and longitude 04° 31' 23.09"E. Some of the dominant species present in the secondary forest in the area includes: *Celtis zenkeri*, *Funtumia elastica*, *Newbouldia laevis* and *Trichilia prieuriana*. The plantation is 38 years old going by the time of its establishment in the year 1967, it was last harvested in 1975. It is a monoculture of *Tectona grandis* trees lying within latitude 07° 32' 26.08"N and longitude 04° 31' 25.19"E and the Riparian vegetation whose age cannot be less than 40 years old, though the actual age cannot be ascertained due to unavailable statistics, is located on latitude 07° 32' 30.06"N and longitude 04° 31' 31.11"E. Some of the dominant species encountered in the riparian vegetation includes: *Celtis mildbraedii*, *Funtumia elastica*, *Pycnanthus angolensis* and *Sterculia tragacantha*.

Estimation of aboveground biomass and carbon stock

Aboveground biomass and carbon stock were estimated in each plot across the different physiognomy. The girth size of all the trees (GBH-1.3 m) greater than or equal to 10 cm in height were enumerated, measured with a tape rule and identified to species level and converted to DBH using the equation

$$DBH = GBH/\pi \quad 1$$

Where: DBH = Diameter at Breast Height, GBH = Girth at Breast Height. $\pi = 22/7$

All identified trees were marked to avoid double enumeration. Tree heights in the secondary forest were measured using a 4m range pole and estimated by the ruler method as stated by Egbe and Tabot (2011). This method was preferred to the altimeter-based measurement because of the closed canopy in the secondary forest. Tree heights in the *Tectona grandis* plantation and the Riparian vegetation were measured using Haga altimeter. The heights of trees and the GBH of all the trees were measured and grouped into different size classes in all the sample plots. Aboveground biomass was calculated using site-specific generated allometric equations developed from measurements such as DBH and tree total height as predictors for the various studied sites.

The site-specific generated equations were developed by plotting DBH as the independent variable against total height, the dependent variable using scattered plot line. The biomass regression equations used for the estimation of the tree species biomass in the secondary forest, *Tectona grandis* plantation and Riparian vegetation were developed from the data obtained from these vegetations using the DBH and the height of the tree species as predictors. A total number of 65 trees in the secondary forest with a DBH ranging from 3 to 37 cm, 87 trees with a DBH ranging from 3 to 34 cm in the plantation and Riparian vegetation having 49 trees with a DBH between 3 and 79 cm were used for the development of individual allometric equations used in the estimation of aboveground biomass in each of these vegetations. The carbon stock was estimated by multiplying the aboveground biomass by a factor of 0.5 (carbon fraction) (IPCC 2003).

Data analysis

The data were first tested for normality and homogeneity in order to satisfy assumptions of Analysis of Variance (ANOVA). One Way Analysis of Variance was employed to test for significant difference between carbon stock in aboveground biomass, soil across the different vegetations. Descriptive statistics was also employed in presenting some of the results. Means of the main effects were compared using Least Significant Difference (LSD) test, using SPSS 17.0 software package.

Results

Aboveground biomass across the different physiognomies

Relationships between biomass of trees in kg, DBH in cm and height in m of the tree species employed in the estimation of the biomass of the vegetations studied are shown in figure 1 to 3. The R^2 -values of the allometric equations explain the relationship between the outcome (biomass) and the values of the DBH and height used for predicting the biomass. It is a measure of how well the allometric equation appropriates the real data points. The R^2 -value indicates a positive, nonlinear relationship between the biomass; DBH and height in all the vegetations (Figure 1-3).




Figure 1: Allometric relationship in the secondary forest; Allometric relationship between aboveground biomass (kg), diameter (cm) and height (m) for tree species in the secondary forest.




Figure 2: Allometric relationship in the *Tectona grandis* plantation; Allometric relationship between aboveground biomass (kg), diameter (cm) and height (m) for tree species in the *Tectona grandis* plantation.




Figure 3: Allometric relationship in the riparian vegetation; Allometric relationship between aboveground biomass (kg), diameter (cm) and height (m) for tree species in the Riparian vegetation.

The high R^2 -values (97% in the secondary forest, 88% in the plantation and 93% in the Riparian vegetation) indicate that DBH and tree height are good predictors of forest aboveground biomass and that the allometric equations are reliable for the estimation of forest biomass.

Aboveground biomass accumulation was found to be higher in the *Tectona grandis* plantation followed by secondary forest and the least value was recorded in the Riparian vegetation (Table 1). The mean aboveground biomass ranged from 7.49 ± 0.90 in the plantation; 8.27 ± 1.07 in the secondary forest to 8.90 ± 3.02 in the Riparian vegetation (Table 1). Across, the three studied sites, there was no significant ($F_{2,198} (0.05) = 0.202$; $P = 0.817$) difference in the mean aboveground biomass (Table 1).

Table 1: Aboveground biomass ($t\ ha^{-1}$) across the various study sites

Name	Maximum	Minimum	Mean \pm std error	Total
Secondary forest	43.21	0.87	$8.27 \pm 1.07^{2.46}$	537.73
<i>Tectona grandis</i> plantation	43.28	0.83	$7.49 \pm 0.90^{2.46}$	637.83
Riparian vegetation	97.52	0.16	$8.90 \pm 3.02^{2.46}$	436.47

*Value in superscript is the LSD value used in comparing the mean difference and mean difference is not significantly different across the column at $P = 0.05$.

The distribution of the aboveground biomass across the different tree size classes across the study sites are presented in Table 2. The 11-20 cm size class contributed more to tree aboveground biomass in secondary forest and *Tectona grandis* plantation, while in the Riparian vegetation; the above 60 cm size class contributed the most (Table 2). The 31-40 cm size class contributed the least to the aboveground biomass in the secondary forest; the 41-50 cm size class is contributing the least in the *Tectona grandis* plantation and the size class 11-20 cm in the Riparian vegetation respectively (Table 2).

Table 2: Size class distribution of tree aboveground biomass ($t\ ha^{-1}$) recorded across the study sites

Size class (cm)	Secondary forest	<i>Tectona grandis</i> plantation	Riparian vegetation
0-10	100.32 (18.7)	122.15 (19.2)	17.51 (4.0)
11-20	276.33 (51.4)	194.88 (30.6)	11.11 (2.5)
21-30	82.96 (15.4)	175.68 (27.5)	50.62 (11.6)
31-40	78.12 (14.5)	115.92 (18.2)	31.75 (7.3)
41-50	Nil	29.20 (4.5)	Nil
51-60	Nil	Nil	43.10(9.9)
Above 60	Nil	Nil	282.38(64.7)

*The percentage contributions of each of the size classes to the tree aboveground biomass are in parenthesis.

The distribution of tree basal area across the study plots are presented in table 3. In the secondary forest, the trees within the 0-10 cm size class had the least basal area and the most was recorded in the 11-20 cm size class (Table 3). Whereas in the *Tectona grandis* plantation, trees within the 41-50 cm size class had the lowest basal area while the highest was recorded in the 11-20 cm size class. In the Riparian vegetation, the above 60 cm size class had the highest basal area and the 11-20 cm size class had the lowest basal area (Table 3).

Table 3: Size class distribution of tree basal area ($\text{m}^2 \text{ha}^{-1}$) recorded across the study sites

Size class (cm)	Secondary forest	<i>Tectona grandis</i> plantation	Riparian vegetation
0-10	4.18	6.71	3.21
11-20	18.16	18.38	2.25
21-30	7.33	15.80	10.34
31-40	7.73	13.71	6.68
41-50	Nil	5.44	Nil
51-60	Nil	Nil	9.32
Above 60	Nil	Nil	58.37

Aboveground carbon stock across the different physiognomies

The estimated amount of carbon accumulated in the trees in the various study sites are presented in Table 4. The estimated carbon stock in the Aboveground carbon stock did not vary significantly ($P < 0.05$) across the various vegetations studied (Table 4).

Table 4. Aboveground carbon stock (t C ha^{-1}) across the various vegetations studied

Name	Maximum	Minimum	Mean \pm std error	Total
Secondary forest	21.61	0.44	$4.14 \pm 0.54^{1,23}$	268.86
<i>Tectona grandis</i> Plantation	16.01	0.42	$3.66 \pm 0.42^{1,23}$	318.92
Riparian vegetation	48.76	0.08	$4.45 \pm 1.51^{1,23}$	218.24

*Value in superscript is the LSD value used in comparing the mean difference and mean difference is not significantly different across the column at $P=0.05$.

Discussion

Aboveground biomass across the different physiognomies

Aboveground biomass was estimated at the different forest types in order to indicate the proportion of biomass. The variation in aboveground biomass from site to site in the study areas might be due to different tree growth stages and tree density. The basal area, especially of the biomass of bigger trees has been reported to be the largest component of above ground forest's biomass (Ogawa *et al.* 1965). The higher aboveground biomass recorded in *Tectona grandis* plantation compared with the estimate in the secondary forest (about 15.6 %) and higher value (about 31.6 %) in secondary forest compared to the Riparian vegetation, can be attributed to higher tree density in the *Tectona grandis* plantation (2175 trees ha^{-1}) followed by (1625 trees ha^{-1}) in the secondary forest and least in Riparian vegetation (1225 trees ha^{-1}). This observation is consistent with the findings of Egbe and Tabot (2011) in their study in Southwestern Cameroun, where it was reported that pure stands of high density trees are bound to have higher carbon stock resulting from higher aboveground biomass than those in mixed stands of tropical forest. The higher tree density recorded in the *Tectona grandis* plantation might be as a

result of high number of tree stands planted or found in the area. The reduction in tree densities in the secondary forest and the Riparian vegetation might be as a result of disturbance (fire) that ravaged the secondary forest some 29 years ago and the human disturbances observed in the Riparian vegetation respectively.

The aboveground biomass estimated for *Tectona grandis* plantation (637.83 Mg ha⁻¹) in this study was higher compared to other studies from plantations across the world. For instance, Duguma *et al.* (2001) reported aboveground biomass of 304 Mg ha⁻¹ for cocoa plantation in South Cameroun, Egbe and Tabot (2011), reported aboveground biomass of 600.72 Mg ha⁻¹ for a *Ricinodendron heudelotii* and of 494.84 Mg ha⁻¹ for *Cola lepidota* plantations in Southwestern Cameroun. Redondo (2007) reported 24.8 to 158.2 Mg ha⁻¹ of aboveground biomass in Costa-Rica. Odiwe *et al.* (2012) also reported aboveground biomass in the *Tectona grandis* plantation to be 38.33 Mg ha⁻¹ in Nigeria. Chittachumnonk *et al.* (2002) who studied carbon sequestration of *Tectona grandis* plantations in Thailand, reported 78.15 Mg ha⁻¹ for aboveground biomass. The general differences in aboveground biomass has been reported to be related to factors such as climatic conditions, solar radiation, disturbances, age of forest, species composition and soil characteristics which varies across different regions (Liao *et al.* 2010). It has also been pointed out that biomass accumulation varies greatly among forest types and ages of forest and that carbon sequestration potential relies on tree size class (Terakunpisut *et al.* 2007).

The highest stem density in size class 0-10 cm and the lowest contribution to biomass accumulation in the secondary forest in the study sites might have resulted to the lowest stem volume and basal area. The implication of this observation is that this vegetation is recovering from disturbance and its developmental stages might be slow. The size class 11-20 cm, 31-40 cm and 41-50 cm in the riparian vegetation, secondary forest and *Tectona grandis* plantation accumulated the least tree biomass respectively. Their low contributions to aboveground biomass accumulation in this study sites was related to low basal area and low stem density which had resulted from the previous fire disturbances in the secondary forest and human disturbance noticed in the riparian vegetation. The low aboveground biomass in the 41-50 cm size class in the *Tectona grandis* plantation might be as a result of the harvest of trees that was done some years ago (1975).

Comparison of the size class distribution and aboveground biomass showed some evidence of biomass reduction in larger size classes from 31-40 cm to above 60 cm especially in the secondary forest and this might be attributed to the ground fire that ravaged this place sometimes ago (Muoghalu and Odiwe 2001). Ground fire is a threat to tropical forests damaging forest stands especially at the young stage of development preventing these forest stands from developing into larger stands which can accumulate more of the aboveground biomass.

The contribution of large trees (DBH ≥ 60 cm) to aboveground biomass in the Riparian vegetation recorded in this study was consistent with the findings of Terakunpisut *et al.* (2007) in Thailand where most aboveground biomass accumulation was found in trees of higher size classes' ≥ 80 –100 and ≥ 100 cm. This indicates that trees of higher size classes play an important role in the biomass accumulation of tropical forest.

Aboveground carbon stock across different physiognomies

Results on carbon sequestration in the different physiognomies showed that the highest amount of carbon was stored in the biomass of trees in the *T. grandis* plantation because of the higher tree density encountered in the *T. grandis* plantation compared to the secondary forest and Riparian vegetation. Hence, calculated carbon stock was higher in the *T. grandis* plantation.

However, tree aboveground carbon stock in the secondary forest and the Riparian vegetation in this study was higher than the results of Hertel *et al.* (2009), where 120 Mg C ha⁻¹ was reported for aboveground carbon storage in a non-Dipterocarp forest in Indonesia. A carbon pool of 150 to 200 Mg C ha⁻¹ has been reported in old-growth forests in South America (Saatchi *et al.* 2007). Brown and Lugo (1982), also reported total carbon sequestration for tropical forest in three countries; Malaysia, Cameroon and Sri Lanka, to be 76.50 Mg C ha⁻¹ in disturbed tropical rain forest (Sri Lanka) and 223 Mg C ha⁻¹ in relatively undisturbed mature tropical rain forest (Cameroun and Malaysia). The highest value was recorded in Malaysia (112.5-223 Mg C ha⁻¹), followed by Cameroun (119-170.5 Mg C ha⁻¹), and the least in Sri Lanka (76.5-110.5 Mg C ha⁻¹). Likewise, aboveground carbon stock in this study in the secondary forest and riparian vegetation were also found to be higher than the result (188 Mg C ha⁻¹) reported by FAO (2010) in Cote d'Ivoire in Tai National park and the results of Sishir and Stephan (2012), where aboveground carbon stock recorded in a naturally forested landscape was 146 Mg C ha⁻¹ in Gabon. The variation in aboveground carbon stocks generally have been pointed out to depend on a number of

factors such as species composition, climate, nutrient conditions, topography, forest age, disturbance and land history management (Vieira et al. 2004, de Castilho et al. 2006, Hertel et al. 2009), and allometric model equation used. All of these factors will influence the development of large-scale policy prescriptions aimed at C-sequestration and that carbon sequestration depended not only on rates of productivity, but also on the size of the trees (Huston and Marland 2003).

The higher carbon sequestration value recorded in the *Tectona grandis* plantation in this study can be attributed to higher tree density in the plantation. The value of aboveground carbon stock (318.92 Mg C ha⁻¹) in the plantation was found to be higher than the carbon stock reported by other workers in other places. For instance, Duguma et al. (2001) reported aboveground biomass carbon stock of 152 Mg C ha⁻¹ for a cocoa agroforestry in South Cameroun; Egbe et al. (2012) reported carbon stock in oil palm to range from 66 to 88 Mg C ha⁻¹ and in rubber to range from 248 to 264 Mg C ha⁻¹ in Cameroun. van Vuuren et al. (1978) has also reported carbon storage for a 25 years old *Pinus patula* plantation and *Eucalyptus grandis* plantation to be 62.6 and 269.9 Mg C ha⁻¹ respectively in South Africa. Chavan and Rasal (2012) reported total standing carbon stock for *Mangifera indica* to be 82.83 Mg C ha⁻¹ in India. Odiwe et al. (2012) reported aboveground carbon stock in *Tectona grandis* plantation to be 38.33 Mg C ha⁻¹ in Nigeria. Chittachumnonk et al. (2002) in their study on carbon sequestration of *T. grandis* plantations in Thailand reported aboveground carbon stocks of 78.15 Mg C ha⁻¹. The difference in tree carbon stock estimates in all these study sites is largely as a result of the form of the regression curve for trees in plantation and the high levels of variability in aboveground carbon estimates. This is as a function of different assumed allometric relationships which affects the size of individual tree canopies, tree-management practices, and crown architecture and this differ considerably by forest type (Nair et al. 2009), species-specific allometry is needed to improve the precision of carbon estimates.

Conclusion

The lower size class 11-20 cm had the highest contribution both in the secondary forest and *Tectona grandis* plantation, unlike the Riparian vegetation where the above 60 cm size class had the highest contribution. This indicated that the secondary forest and *T. grandis* plantation are younger or be relatively disturbed and are just recovering from the disturbance.

References

- Brown S and Lugo AE. Aboveground biomass estimates for tropical moist forests of the Brazilian Amazon. *Interciencia*. 1992;17:8-18.
- Chavan B and Rasal G. Total sequestered carbon stock of *Mangifera indica*. *Journal of Environment and Earth science*. 2012;2:37-48.
- Chittachumnonk PC, Sutthisrisinn S, Samran C, Viriyabuncha and Peawsad K. Improving estimation of annual biomass increment and aboveground biomass of Teak plantation using site - specific allometric regressions in Thailand. 2002.
- de Castilho CV, Magnusson WE, de Araujo RNO, Luizao RCC, Lima AP, Higuchi N. Variation in aboveground tree live biomass in a central Amazonian forest: effects of soil and topography. *Forest Ecology and Management*. 2006;234:85-96.
- Duguma B, Gockowski J and Bakala J. Small holder Cacao (*Theobroma cacao* L.) cultivation in agroforestry systems of West and Central Africa: challenges and opportunities. *Agroforestry System*. 2001;51:177-188.
- Egbe EA and Tabot PT. Carbon sequestration in eight woody non timber forest species —and their economic potentials in Southwestern Cameroon. *Applied Ecology Environmental Research*. 2011;9:369-385.
- Food and Agriculture Organisation. Global Forest Resources Assessment (FRA) 2010 Main Report. *FAO Forestry Paper* 163, Food and Agriculture Organization (FAO) of United Nations, Rome. 2010.

320
321 Fonseca W, Rey Benayas JM and Alice FE. Carbon accumulation in the biomass and soil of different
322 aged secondary forests in the humid tropics of Costa Rica. *Forest Ecology and Management*.
323 2011;262:1400-1408.
324
325 Gibbs Holly K, Sandra Brown, John O Niles and Jonathan A Foley. Monitoring and estimating tropical
326 forest carbon stocks: making REDD a reality. *Environmental Research Letters* (2) 2007.
327
328 Hall JB. 1969. The Vegetation of Ile-Ife. University of the Ife Herbarium Bulletin 1.
329
330 Hertel D, Moser G, Culmsee H, Erasmi S, Horna V, Schuldts B and Leuschner C. Below and above-ground
331 biomass and net primary production in a paleotropical natural forest (Sulawesi, Indonesia) as compared
332 to neotropical forests. *Forest Ecology and Management*. 2009;258: 1904-1912.
333
334 Huston MA and Marland G. Carbon management and biodiversity. *Journal of Environmental*
335 *Management*. (2003).
336
337 Karjalainen T, Pussinen A, Liski J, Nabuurs G, Erhard M, Eggers T et al. An approach towards an
338 estimate of the impact of forest management and climate change on the European Forest Sector
339 carbon budget: Germany as a case study. *Forest Ecology and Management*. 2002;162: 87-103.
340
341 Keay RWJ. 1959. An Outline of the Nigerian Vegetation (3rd edn). Government Printer: Lagos, Nigeria.
342
343 Komiya A, Ong JE and Pongpan S. Allometry, biomass, and productivity of mangrove forests: a
344 review. *Aquatic Botany*. 2008;89:128-137.
345
346 Liao C, Luo Y, Fang C and Li B. Ecosystem carbon stock influenced by plantation practice: Implications
347 for planting forests as a measure of climate change mitigation. *Plos ONE* 5. 2010.
348
349 Ludang Y and Jaya HP. Biomass and carbon content in tropical forest of Central Kalimantan. *Journal of*
350 *Applied Sciences in Environmental Sanitation*. 2007;2:7-12.
351
352 Muoghalu JI and Odiwe AI. Ecosystem dynamics in a Nigerian Secondary Lowland Rainforest 14 years
353 after a ground fire: Tree species population dynamics. *Nigeria Journal of Botany*. 2001;14:7-24.
354
355 Nair PKR, Kumar BM and Nair VD. Agroforestry as a strategy for carbon sequestration. *Journal of Plant*
356 *Nutrition and Soil Science*. 2009;172:10-23.
357
358 Ogawa H, Yoda K, Ogino K and Kira T. Comparative ecological studies on three main
359 type of forest vegetation in Thailand II. Plant Biomass. *Nature and Life in Southeast Asia*. 1965;4: 49-80.
360
361 Oke SO and Isichei AO. Floristic and structure of the fallow vegetation in Ile-Ife area of Southwestern
362 Nigeria. *Nigeria Journal of Botany*. 1997 vol. 10, 30-50.
363
364 Redondo A. Growth, carbon sequestration, and management of native tree plantations in humid regions
365 of Costa Rica. *New Forests*. 2007;34:253-268.
366
367 Saatchi SS, Houghton RA, Dos Santos Alvala RC, Soares JV and Yu Y. Distribution of aboveground live
368 biomass in the Amazon Basin. *Global Change Biology*. 2007;138:16-37.
369
370 Sheikh MA, Kumar S and Kumar M. Above and below ground organic carbon stocks in a sub-tropical
371 *Pinus roxburghii* Sargent forest of the Garhwal Himalayas. *Research Article For Students China*.
372 2012;14(3):205-209.
373

Formatted: Indent: Left: 0", Hanging: 0.5"

374 | Stephens BB, Gurney KR, Tans PP, Sweeney C, Peters W, Bruhwiler L et al. Weak northern and
375 | strong tropical land carbon uptake from vertical profiles of atmospheric CO₂. Science. 2007; 22(316):
376 | 1732-1735.

377 |
378 | Strassburg B, Turner RK, Fisher B, Schaeffer R and Lovett A. -Reducing emissions
379 | from deforestation—The combined incentives mechanism and empirical simulation. Global
380 | Environmental Change. 2009;19: 265-278

381 |
382 | Terakunpisut J, Gajasen N and Ruankawe N. Carbon sequestration potential in aboveground biomass of
383 | Thong Pha Phum National Forest, Thailand. Applied Ecology And Environmental Research.
384 | 2007;5(2):93-102.

385 |
386 | Van Vuuren NJJ, Banks CH and Stohr HP. Shrinkage and Density of Timbers used in the Republic of
387 | South Africa. Bullentin 57. 1978. Department of Forestry, Pretoria.

388 |
389 | Vieira S, de Camargo PB, Selhorst D, da Silva R, Hutya L, Chambers JQ et al. Forest structure and
390 | carbon dynamics in Amazonian tropical rain forests. Oecologia. 2004;140: 468-479.

391 |
392 | White F. -The Vegetation of Africa Descriptive Memoir to Vegetation Map Africa.UNESCO: Paris. 1983.
393 |
394 |
395 |