

Original Research Article

Trends in epidemiology, susceptibility pattern and serotypes of Salmonellae at a tertiary care hospital: An eight year study

Comment [C1]: Corrected

Comment [C2]: ???

Abstract:

Background- Enteric fever is a global disease. In India Enteric fever is endemic with *Salmonella enterica* serovar Typhi being the predominant etiological agent. Due to changing antimicrobial resistance patterns, knowledge of local epidemiology, antimicrobial resistance pattern helps in initiation of appropriate empiric therapy.

Comment [C3]: ???

Comment [C4]: ????

Comment [C5]: ????

Methodology A prospective study on Salmonellae isolated from blood and stool specimens over 8 year period was conducted. Antimicrobial susceptibility was done as per CLSI guidelines. Serotyping was done by using commercial antisera and later confirmed at Central Research Institute, Kasauli.

Comment [C6]: ????

Results Out of 52 salmonellae, 43 (82.6%) were from blood and 8 (15.3%) from stool and 1 (1.9%) from pus specimen. We observed a change in spectrum and susceptibility pattern of salmonellae the 8 year study period. In 2011, 2013, 2016 and 2018, *Salmonella* Typhi (serotype-9,12,vi:d:-) was the predominant etiological agent accounting for 81.8% , 66.6%, 51% and 80% of the total cases of salmonella respectively. *Salmonella* Paratyphi B (4,12:b:1,2) was predominant in 2012 (100% of cases). *Salmonella* Serotype Typhimurium (4,12;i:1,2) was predominant in 2014 (50%) while *Salmonella* Typhi and *S. Paratyphi* B contributed equally to infections in 2015 (40% each). From 2011, Non typhoidal salmonellae (NTS) steadily increased. 19 (36.6%). The most effective antimicrobials against typhoidal salmonellae were chloramphenicol, ceftriaxone and co-trimoxazole with all most 100% sensitivity from 2011 to 2018. Ciprofloxacin maintained good sensitivity in 2013, 2014 and 2015, 2016 and 2018 but ampicillin was ineffective in our set-up .

Comment [C7]: ???

Conclusions- Due to changing trends in spectrum and sensitivity of salmonellae, continuous monitoring is essential.

Introduction-

Typhoid fever remains an important global public health problem accounting for 12-33 million cases worldwide.^[1] Around 80% of these cases occur in Asia alone.^[2] Many published Hospital based studies and outbreak studies suggests that Typhoid fever is a major public health concern in India with *Salmonella enterica* serovar Typhi (*Salmonella* Typhi) being main etiological agent.^[3] Antimicrobial treatment is the mainstay of treatment of Typhoid and Paratyphoid fever.^[4] Emergence of Antimicrobial resistance can pose a

38 challenge for effective management of typhoid fever,^[5] especially the emergence and spread
39 of multidrug resistant strains.

40 In India, drug resistant Salmonellae have been reported since 1960, first outbreak of
41 multidrug resistant *Salmonella* Typhi occurring in Calicut. Since then multi drug resistant
42 *Salmonella* Typhi have appeared throughout the world, especially in South America, the
43 Indian sub continent, Africa and South East Asia.^[6] Later an outbreak due to chloramphenicol
44 resistant *Salmonella* Typhi was reported from Chandigarh.^[7]

45 Subsequently, resistance to commonly used antibiotics such as chloramphenicol, ampicillin
46 and cotrimoxazole has been reported from different parts of India.^[5,7]

47 The present study was undertaken to know the trends in serotypes and antibiograms of
48 Salmonellae isolates in a hospital setting over a 8 year period.

49 Methodology – A prospective study was conducted in the department of microbiology of
50 Sassoon General Hospital Pune over a period of 8 years. Various clinical specimens like
51 blood, stool, urine, and pus were processed for culture by routine methods and Salmonellae
52 isolates were included in the study. Identification of salmonella was done by standard
53 microbiological methods.[8] All the salmonella isolates were tested by commercially
54 available(DENKA SEIKEN) Salmonella polyvalent antisera and group specific antisera (O9,
55 O4 and O2). The isolates were preserved and also sent to CRI, Kasauli for serotyping. The
56 antimicrobial susceptibility testing was done by Kirby Bauer disk diffusion method as per
57 CLSI guidelines.[9] According to CLSI, antimicrobial sensitivity for non-typhoidal
58 salmonellae is not recommended. So sensitivity was not analysed for non-typhoidal
59 salmonellae The data was entered in WHONET.

60 Results-

61 A total of 52 salmonellae were isolated over a period of 8 years. Out of the 52 isolates, 43
62 (82.6%) were obtained from blood cultures and 8 (15.3%) were obtained from stool
63 specimens and 1 (1.9%) from pus. Demographic data revealed that males (56.4%) were more
64 affected by Salmonella than females (43.6%).

65 Out of 52 salmonellae, 31 (59.6%) were *Salmonella enterica serovar* Typhi followed by
66 *Salmonella* Paratyphi B. 9 (17.3%), *Salmonella* Typhimurium, 4 (7.6%), *Salmonella*
67 Paratyphi A 3 (5.7%) . *Salmonella* Jaffna and *Salmonella* Enteridis each were 3.8% of the
68 isolates,. There was one isolate *Salmonella* Welteverden. Year wise distribution of the
69 isolates revealed changing trends in etiology of typhoid fever. In 2011, 2013, 2016 and 2018,
70 *Salmonella* Typhi (serotype-9,12,vi:d:-) was predominant etiological agent accounting to
71 81.8%, 66.6%, 51% and 80% of the total cases of salmonella respectively. *Salmonella*
72 Paratyphi B was predominant in 2012 (100% of cases). *Salmonella* Ser. typhimurium was
73 predominant in 2014 (50%) while *Salmonella* Typhi and *Salmonella*.Paratyphi B contributed
74 equally in 2013, 2015 (40% each). In 2011 *Salmonella* Typhi was predominant isolate but
75 after that non typhoidal salmonellae are steadily increasing.(Table 1)


76

77

78 Table 1-Yearwise distribution of Salmonellae from clinical samples (n=52)

Year	Isolates	No. of isolates	Serotypes
2011 (n=22)	<i>Salmonella</i> Typhi	18	9,12,vi:d:-
	<i>Salmonella</i> Paratyphi B	2	4,12:b:1,2
	<i>Salmonella</i> Typhimurium	2	4,12;i:1,2
2012 (n=2)	<i>Salmonella</i> Paratyphi B	2	4,12:b:1,2
2013 (n=6)	<i>Salmonella</i> Typhi	2	9,12,vi:d:-
	<i>Salmonella</i> Paratyphi B	2	4,12:b:1,2
	<i>Salmonella</i> Enteridis	2	9,12:g,m:-
2014 (n=4)	<i>Salmonella</i> Paratyphi A	1	2,12;a:-
	<i>Salmonella</i> Typhi	1	9,12,vi:d:-
	<i>Salmonella</i> Typhimurium	2	4,12;i:1,2
2015 (n=5)	<i>Salmonella</i> Typhi	2	9,12,vi:d:-
	<i>Salmonella</i> Paratyphi B	2	4,12:b:1,2
	<i>Salmonella</i> Welteverden	1	3,10:r:z6
2016 (n=6)	<i>Salmonella</i> Typhi	3	9,12,vi:d:-
	<i>Salmonella</i> Paratyphi A	1	2,12;a:-
	<i>Salmonella</i> Jaffna	2	9,12:d:Z39
2017 (n=2)	<i>Salmonella</i> Typhi	1	9,12,vi:d:-
	<i>Salmonella</i> Paratyphi A	1	2,12;a:-
2018 (n=5)	<i>Salmonella</i> Typhi	4	9,12,vi:d:-
	<i>Salmonella</i> Paratyphi B	1	4,12:b:1,2

79 Fig 1- Antibiogram of Typhoidal Salmonella showing % susceptibility. (n=35)



80

81

82 The most effective antimicrobial agent against typhoidal salmonellae was chloramphenicol
 83 with 100% sensitivity. Next drug found to be effective was Co-trimoxazole with again 100%
 84 sensitivity over all years except in 2014 (50%). Fluroquinolones like ciprofloxacin has
 85 maintained good sensitivity in 2013, 2014 and 2015, 2016 and 2018. Surprisingly in our

86 study we found very low resistance to 3rd generation cephalosporins in 2011 and 2014.
87 However, ampicillin was ineffective in our set-up with almost 0% sensitivity in 2012, 2014
88 and 2017.

89 Discussion-

90 Antimicrobial resistance is a major hindrance in successful treatment of typhoid fever.
91 Environmental conditions like poor sanitation, bad personal hygiene, poor quality water
92 aggravates the problem. In the present study, we analysed trends in etiology of typhoid fever
93 and the susceptibility pattern of Salmonellae isolated from a tertiary care centre in western
94 Maharashtra. Demographic data in our study revealed that males were affected more than
95 females by salmonella which is in accordance to the finding by Saba et al^[10] who also found
96 that 90 (58%) isolates were obtained from male and 64 (42%) from female patients.
97

98 *Salmonella* Typhi was predominant pathogen isolated over the study period. It was also the
99 commonest Samonella in 2011, 2013, 2016 and 2018 accounting for 81.8% ,66.6%, 51% and
100 80% of the cases respectively. Similar finding has been noted by V. Laxmi et al from
101 Hyderabad, India in 2006.^[11]

102 In the present study, from year 2012 onwards non-typhoidal salmonellae (NTS) emerged.
103 And formed 32.6% of the isolates from 2012 to 2018. *Salmonella* Typhimurium, *Salmonella*
104 Welteverden and *Salmonella* Enteridis, *Salmonella* Jaffna were the NTS detected. Similar
105 findings have been reported by Suman Kanungo et al^[12] [2008, kolkatta, India] in their
106 review article. They have shown an increasing incidence of invasive salmonellosis due to
107 Non-typhoidal salmonellae. In a study in Thailand, 135 cases of NTS bacteraemia have been
108 reported.^[13] But in contrast to this study only 2 cases of NTS from 1500 blood cultures has
109 been reported by an Indian study.^[2]

110 Drug resistance is a major challenge when treating typhoid fever. Chloramphenicol has
111 remained the treatment of choice for typhoid fever for around six decades now.
112 Chloramphenicol therapy is reduces mortality due to typhoid fever from 20% to 1% and
113 duration of fever from 14-28 days to 3-5 days.^[14] However, chloramphenicol has its own side
114 effects like bone marrow toxicity, high carriage rates and emergence of drug resistance. In
115 1980s there was emergence of plasmid mediated chloramphenicol resistance in many
116 countries including India.^[15] In this scenario next options were ampicillin and Co-
117 trimoxazole^[16] This was followed by emergence of multidrug resistant (MDR) strains
118 (combined resistance to chloramphenicol, ampicillin and co-trimoxazole) initially reported
119 from India [karnataka,1999]^[17], Pakistan and Middle East and then from all over the
120 world.^[18]

121 In the present study good sensitivity to chloramphenicol and co-trimoxazole was observed
122 over the 8 years. Similar findings have been mentioned by other Indian authors like shorey et
123 al[Mumbai,1993]^[19] and Nath et al from Varanasi,2003.^[20] In the current study there was
124 very low sensitivity to ampicillin almost 0% in 2012, 2014 and 2017. Increasing use of
125 ampicillin seems to have decreased its efficacy in treatment of typhoid fever

126 Parenteral administration of 3rd generation cephalosporins specially ceftriaxone is often the
127 treatment of choice for typhoid fever due to its short duration of therapy as compared to long
128 duration of chloramphenicol. In the present study, in 2011 and 2014, we observed diminished
129 sensitivity to ceftriaxone (66%) but it regained sensitivity in 2015 to 2018 (100%). This
130 finding is similar to findings by saba et al,^[10] who observed increased sensitivity to
131 salmonella from 92% to 100%. Other Indian studies by Nath et al^[20] and Gautam et al^[6] also
132 mentioned increasing sensitivity to 3rd generation cephalosporins in their studies.

133 Ciprofloxacin was used as a good alternative to chloramphenicol when it was initially
134 introduced in 90s but because of overuse and misuse of the drug it also showed resistance.

135 Gautam et al reported diminished sensitivity to ciprofloxacin from 89% to 81% from 1997 to
136 2001.^[6] In the present study, it was observed that there was diminished sensitivity to
137 ciprofloxacin in 2011 (66.6%) and 2012 (50%) but it improved to 100% in 2013, 2015, 2016
138 and 2018. This could be due to the fact that resistance to this drug made bit ineffective and
139 most clinicians stopped using this drug for the treatment of typhoid fever. Hemlatha et al from
140 Hyderabad also observed 95% sensitivity to ciprofloxacin in the year 1999.^[21] In the present
141 study, we observed a lot of variation in serotypes of salmonellae causing typhoid and also in
142 susceptibility pattern of salmonella species. So continuous monitoring of Isolates causing
143 enteric fever and their susceptibility to antimicrobials is recommended.

144
145 Conclusion- The study highlights the changing trends in etiology and susceptibility pattern of
146 salmonellae causing typhoid fever. So, continuous monitoring of microorganisms causing
147 enteric fever is important for optimum treatment of typhoid fever.

148
149 References-

- 150 1) Miller SI, Pegues DA. Salmonella species, including *Salmonella Sertype Typhi*,
151 chapter 210. In: Mandell GL, Bennett JE and Dolin R, *Principles & Practice of*
152 *infectious diseases*, 5th edn. (Churchill Livingstone, London); 2000. p. 2344–63.
- 153 2) Ochiai RL, Acosta CJ, Danovaro-Holliday MC , Baiqing D, Bhattacharya SK, Agtini
154 MD et al . A study of typhoid fever in five Asian countries: disease burden and
155 implications for controls. Bull World Health Organ 2008; 86: 260-8.
- 156 3) AS Damle, RP Karyakarte, AB Deshmukh, MP Bansal. Salmonella phage types
157 prevalent at Aurangabad. Indian journal of medical microbiology 1988; 6, 151-154.
158
- 159 4) Preeti Behl, Varsha Gupta, Atul Sachdev, Vishal Guglani & Jagdish Chander.
160 Patterns in antimicrobial susceptibility of Salmonellae isolated at a tertiary care
161 hospital in northern India. Indian J Med Res 145, January 2017, pp 124-128.
- 162 5) Jacob John, Carola J. C. Van Aart, Nicholas C. Grassly. The Burden of Typhoid and
163 Paratyphoid in India: Systematic Review and Meta-analysis. PLOS Neglected
164 Tropical Diseases | DOI:10.1371/journal.pntd.0004616 April 15, 2016.
- 165 6) Gautam V, Gupta NK, Choudhary U, Arora DR. Sensitivity Pattern of Salmonella
166 serotypes in Northern India. *Braz J Infect Dis* 2002;6:281–7.
- 167 7) Kapil A, Ayyagiri A, Garg RK, Agarwal KC. *Salmonella Typhi* with transferable
168 Chloramphenicol Resistance Isolated in Chandigarh during 1983–87. *Indian J Pathol*
169 *Microbiol* 1994;38:179–83.
- 170 8) Forbes BA, Sahm DF, Weissfeld AS. In: Bailey and Scott's Diagnostic Microbiology.
171 12th ed. Missouri: Mosby Elsevier; 2007. p. 779.
- 172 9) CLSI performance standards for antimicrobial susceptibility testing. Twelfth
173 international supplement. NCCLS document M100-S12 [ISBN1-56238-454-6].
174 NCCLS, 940 WEST VALLEY Road, suite 1400, Wayne Pennsylvania 19087-1898
175 USA, 2002.
- 176 10) Saba Qaiser, Seema Irfan, Erum Khan, Tanwir Ahsan, Afia Zafar. In Vitro
177 Susceptibility of typhoidal Salmonellae against newer antimicrobial agents: A search
178 for alternate treatment options. J Pak Med Assoc, May 2011; Vol. 61(5):462-465.
- 179 11) V Lakshmi, R Ashok, J Susmita, VV Shailaja . Changing trends in the antibiograms
180 of *Salmonella* isolates at a tertiary care hospital in hyderabad. *Indian Journal of*
181 *Medical Microbiology*, (2006) 24 (1):45-8.
- 182 12) Suman Kanungo, Shanta Dutta, and Dipika Sur1. Epidemiology of typhoid and
183 paratyphoid fever in India. *J Infect Developing Countries* 2008; 2(6): 454-460.

- 184 13) Kiratisin P (2008) Bacteraemia due to non-typhoidal *Salmonella* in Thailand: clinical
185 and microbiological analysis. *Trans R Soc Trop Med Hyg* 102(4):384-8. Epub 2008
186 Mar 5.
- 187 14) Mandal S, Mandal MD, Kumar NP. Reduced minimum inhibitory concentration of
188 chloramphenicol for *Salmonella enterica* serovar typhi. *Indian J Med Sci* 2004;**58**:16-
189 23.
- 190 15) Srivastava L, Aggarwal P. Multidrug resistant *Salmonella Sertype Typhi* in Delhi.
191 *Indian J Med Microbiol* 1994;**12**:102-5.
- 192 16) Parry CM. The treatment of multi-drug resistant & Nalidixic acid-resistant typhoid
193 fever in Vietnam. *Trans Roy Soc Trop Med Hyg* 2004;**98**:413-22.
- 194 17) Ciraj AM, Seetha KS, Gopalakrishna BK, Shivananda PG .Drug resistance
195 pattern and Phage types of *Salmonella typhi* isolates in Manipal, South
196 Karnataka. *Ind J Med Sci* 1999 ;53(11): 486-89
- 197 18) Bhutta ZA. Current concepts in the diagnosis and treatment of typhoid fever. *BMJ*
198 2006; 333: 78-82.
- 199 19) Sheorey HS, Kaundinya DV, Hulyalkar VS, Deshpande AK. Multidrug Resistant
200 *Salmonella Sertype Typhi* in Bombay. *Indian J Pathol Microbiol* 1993;**36**:8-12.
- 201 20) Nath G, Tikoo A, Manocha H, Tripathi AK, Gulati AK. Drug resistance in *Salmonella*
202 *Sertype Typhi* in North India with special reference to Ciprofloxacin. *J Antimicrobial*
203 *Chemother* 2003;**46**:145-53.
- 204 21) Hemalatha R, Vijayalakshmi P, Gyaneshwari, Rao MV, Ramani A. Multidrug
205 resistant salmonella typhi in Hyderabad. *Indian J Med Microbiol* 1999;**17**:39-41.

206
207