

# Review Article

## State of the art: Soil physical attributes

### ABSTRACT

Proper soil management interferes with the result of the installed crop. The knowledge of the positive and/or negative influence on the production systems is essentially essentialimportant to improve the physical, chemical and biological quality of the soil, for that, there are some attributes that act as indicators of soil quality. The practices carried out improperly will result in problems in soil structure, as compaction, lack of availability of water and air in the soil and for plants, soil loss among others. Some properties that as soil porosity, soil aggregation, soil compaction, and soil water infiltration are used to measure soil quality. GivenIn view of this, the use of these attributes as indicators is signified extremely important for excellent good productivity, since management practices used can directly influence the development of plants.

*Keywords: Soil quality, soil properties, soil management.*

### 1. INTRODUCTION

The quality of the soil is variable to its formation, textural composition and type of management adopted, which determines their behavior in the face of anthropogenic activities. The conversion of forest into agricultural areas or pasture areas has been causing severerious problems due to the adoption of inadequate management. There are indicators that determine soil quality and verify the effectiveness of the practices adopted [1].

The use of unsuitable practices in the soil can result in severerious problems to its structure, aggregate stability, the degree of compaction, resulting in insufficient infiltration of water into it, which hinders the availability of the resource to crops, in addition to significantly increase erosive processes. Several attributes must be measured to to evaluate how management influences soil characteristics as well as their relationship to the plant [2].

The understanding of the physical behavior of a soil is of utmost importance, since it guides the proper activities that must be performed in the system, so that in this manner, it reaches an adequate crops development. This diagnosis involves the arrangement of particles and pores, soil bulk density, aggregation structure, mechanical penetration resistance, soil water infiltration, water availability to plants [3].

### 2. SOIL POROSITY

Due to the structure or arrangement between the soil particles, in addition to the fraction or volume of solids, there is also a volume of voids (pores), in which factors such as retention, movement and availability of water, aeration, availability of nutrients, resistance to root penetration, aggregate stability and compaction, to a lesser or greater degree.

Formatted: No underline, Underline color: Auto

Formatted: No underline, Underline color: Auto

Formatted: No underline, Underline color: Auto

Formatted: No underline, Underline color: Auto

Formatted: No underline, Underline color: Auto

Formatted: No underline, Underline color: Auto

Formatted: No underline, Underline color: Auto

54 According to Teixeira et al. [4], porosity is a physical property defined by the  
55 relationship between the pore volume and the total volume of a particular certain  
56 material, and according to Embrapa[5], porosity is constituted by the porous space,  
57 after the arrangement of the components of the substantialolid part of the soil and  
58 which, under natural conditions, is occupied by water and air, being divided into  
59 primary and secondary.

60 Primary porosity is developed with the sediment or rock, being characterized in  
61 the sedimentary rocks by the spaces between clasts or grains (intergranular porosity)  
62 or stratification planes. Worth noting that in sedimentary materials, the size and shape  
63 of the particles, their degree of selection and the presence of cementation influence the  
64 porosity. The secondary porosity develops after the formation of igneous, metamorphic  
65 or sedimentary rocks, by fracture or failure during their deformation (fracture porosity)  
66 [4].

67 Sands retain an inadequate, poor amount of water because their large porous  
68 space allows free water drainage from the soils. Clays absorb relatively large amounts  
69 of water, and their smaller porous spaces hold it against the forces of gravity.

70 In short, porosity consists of the physical quantity given by the volume of the  
71 porous space, constructed by the arrangement of the components of its  
72 substantiallyolid part and which, under natural conditions, is occupied by water and air  
73 [6].

74 Regarding to the distribution and size of the pores is oriented by three types of  
75 classification, consisting of macropores (pores with larger diameter, which directly  
76 influences the infiltration capacity, soil drainage and its aeration capacity); mesopores  
77 (pores with intermediate diameter, responsible for the conduction of water during the  
78 redistribution process, that it, after infiltration, when the macropores are emptied);  
79 micropores (pores with the smallest diameter responsible for the retention and storage  
80 capacity of water and solutes in the soil [7].

81 According to Lorenzo[6], the macropores (Ma) are results of the arrangement of  
82 the aggregates, the action of the mesofauna and roots and the expansion and  
83 contraction of the soil mass. They are related to the gas exchange of oxygen and  
84 carbon dioxide and to the flow of water by gravity: infiltration, drainage, and transport of  
85 solutes; and micropores (Mi) are in-aggregated and are related to water retention due  
86 to molecular adhesion that entraps gases, vapors or solids in the surface of solid  
87 bodies. Kiehl [8] classifies as macro and micropores, pores with a larger and smaller  
88 diameter, respectively, than 0.06 mm. Several authors include mesopores in this  
89 classification as an intermediate class, such as Luxmoore [9], which suggested a  
90 classification in which the micropores have a diameter smaller than 0.01 mm; the  
91 mesopores have a diameter between 0.01 and 1.0 mm; and the macropores, diameter  
92 greater than 1.0 mm.

93 Soil porosity interferes with aeration, conduction, and retention of water,  
94 resistance to penetration and root branching in the soil and, consequently, in the use of  
95 available water and nutrients [10].

96 Ideal soil must present a volume and size of pores suitable for the entry,  
97 movement and retention of water and air to meet crop needs [11]. The distribution of  
98 pores in the soil matrix plays a fundamental role in the relationships between the solid,  
99 liquid and gaseous phases, determining the spatial and temporal evolution of the  
100 processes that involve the movement of water in the soil [12]. According to Ribeiro et  
101 al. [12], soil porosity is determined by the way the solid particles are arranged,  
102 emphasizing that if they are arranged in close contact, predominance of solids occurs  
103 in the sample and the porosity is low; and if, on the contrary, the particles are arranged  
104 in aggregates, there is a predominance of voids in the soil sample and the porosity is  
105 high.

106  
107

### 3. SOIL AND PARTICULATE DENSITY

108 The diversity of the mineral and organic components presents in the composition  
109 of the soils, as well as the proportion between them, determine the density of the  
110 material. This physical attribute besides being determinant of the composition is also  
111 related to soil texture and aggregation, water infiltration rate and erosion,  
112 macroporosity and root development, soil consistency (dry, wet and wet), the degree of  
113 compaction, which interferes with root development and management techniques and  
114 agricultural productivity. The density is oriented by determining the soil density (ratio of  
115 the sample mass to the volume occupied by solids, considering the pore space) and  
116 the density of particles (ratio of the sample mass to the volume occupied by the  
117 particles, disregarding the porous space).  
118

### 119 3.1 Soil density

120 Soil density is defined by the ratio of the mass of dry solids to the soil volume,  
121 being affected by crops that alter the structure, consequently the arrangement and  
122 volume of the pores. These changes influence soil physical properties, such as  
123 aeration porosity, soil water retention, plant water availability and resistance to root  
124 penetration [3].

125 A soil sample of the surface horizon, rich in organic matter (substrate), when  
126 compared to a portion of any of the horizons in depth, it is perceived that the superficial  
127 sampling is lighter. The significant increase in soil density in depth can be explained by  
128 the pressures exerted by the upper layers, causing compaction and reduction of pore  
129 volume [8].

130 Association of the concepts of density and porosity, between the masses and the  
131 volume of the soil constituents, are developed by porosity, which determines the  
132 existent space between the aggregates, occupied by air or water, being calculated from  
133 density, the pore space occupied varies in the inverse ratio of soil density [13].

134 This physical attribute is expressed in grams per cubic centimeters and the  
135 amplitudes of variation for each type of soil is within the following limits: clayey soils  
136 ( $0.90$  to  $1.25 \text{ g cm}^{-3}$ ); sandy soils ( $1.25$  to  $1.60 \text{ g cm}^{-3}$ ); humic soils ( $0.75$  to  $1.00 \text{ g cm}^{-3}$ );  
137 turfous soils ( $0.20$  to  $0.50 \text{ g cm}^{-3}$ ).

138 The determination methods are based on obtaining the mass and volume of the  
139 soil sample. The mass is readeasily determined by weighing the dry soil in an oven,  
140 and the determination of the volume is varied from the use of some methods, which are  
141 described below:  
142

#### 143 3.1.1 Volumetric ring method

144 There are several types of samplers, the most usual it's a stainless-steel  
145 cylinder with sharp edges, nailed directly into the soil. This method presents  
146 particularcertain difficulties in the removal of the ring from the soil, may occurring loss  
147 of sample, since there is no soil surplus at the top and bottom of the cylinder under  
148 comparable structure conditions, the higher the clay content of a soil, the lower its  
149 density, always considering the composition of the soil analyzed [3].

150 This method has been used since 1914, suitable for well-structured soils.  
151 However, when the soil has thick roots or is a compact horizon, it is unfeasible to use  
152 and is not recommended in these situations [8].  
153  
154

#### 155 3.1.2 Method of the waterproofed clod

156 Based on the Archimedes' Law, which defines the buoyancy of a body is equal to  
157 the weight of the volume of liquid displaced when it is immersed into it. This method is  
158 not recommended for mobilized soils, since in this condition the aggregates will be of  
159 equal density to that of before the preparation. The volume of the clods is determined  
160 by the volume of water displaced by them immersed in water [8].

Formatted: No underline, Underline color: Auto

161 | This method presents a certain disadvantage, due to the possibility of  
162 | segregating the soil sample during the collection process, thus generating a disregard  
163 | for the existence of macropores in the clods.

164

### 165 | **3.2 Density of particles**

166 | This soil physical attribute aims to measure the average density of the mineral  
167 | and organic particles of the soil, reflecting its average composition. This density is  
168 | related to the volume effectively occupied by solid matter, without considering the  
169 | porosity. The mineralogy and soil composition are characteristics that naturally  
170 | influence the density of individual soil particles [14].

171 | Some incorrectly practices performed may increase soil density, such as  
172 | excessive tilting or use of poor's conservation practices, which may cause structural  
173 | alteration, a decrease of macroporosity and total porosity, among other damages  
174 | [15,1].

175 | The problem of having a compacted soil and consequently the increases of its  
176 | density and resistance, is the difficulty that the root system will have to penetrate and  
177 | exploit this soil, thus reducing the pore diameter of the soil, reducing permeability and  
178 | flow of water, as well the air capacity, which may affect the development of plants and  
179 | the anatomical structures of its roots [16].

180 | The mean values for each soil type depend on its predominant mineral  
181 | constituents, with a mean variation between the limits of 2.3 to 2.9 g cm<sup>-3</sup>. The  
182 | vasgreat majority of the soils are composed of quartz, feldspar and colloidal aluminum  
183 | silicates, whose particle density is around 2.65 g cm<sup>-3</sup>.

184 | The methods for determining the density of soil particle are based on obtaining  
185 | the sample mass value and then the volume of present solids, the mass is obtained by  
186 | simple weighing, and the volume can be obtained by the volumetric flask method, more  
187 | accurate among existing methods. The differential of this method is the practicality  
188 | offered, in which it is summarized in a single weighing, pipetting and buret reading, of  
189 | the displaced volume [8].

190

## 191 | **4. SOIL AGGREGATION**

192 | Aggregate is characterized as a grouping of strongly adhered particles, the size of  
193 | the aggregate determines its susceptibility to movement by the wind, water, and porous  
194 | space, interfering in the percolation of the water and the volume occupied by the air of  
195 | the soil, being conditioned from the environment to the growth of the root system of  
196 | plants. Organic matter is an essentialimportant cementing agent of soil particles,  
197 | vegetation and its residues protecting the aggregates from the surface, against  
198 | disaggregation due to the impact of rainfall and sudden variations of humidity [3].

199 | The soil structure is adequate to allow propergood flow of water, inner's aeration,  
200 | resistance to erosion and traffic of machinery, development of living organisms and  
201 | proper development of plant roots [17].

202 | The soil structure is represented by the aggregation, that is, the result of the  
203 | interaction between the size, shape, and arrangement of the solid particles and porous  
204 | spaces of the soil, being highly variable and associated with physical, chemical and  
205 | biological factors [18]. These properties, with the genetic potential of the plants,  
206 | determine the productivity of the crops [19].

207 | The dynamics of soil aggregation is influenced by the soil management system.  
208 | This management comprises a set of practices that, when rationally used, promote  
209 | better crop productivity, but when improperly used, cause physical, chemical and  
210 | biological degradation of the soil and, also, a reduction of productivity [20, 21].

211 | In the last years, soil quality studies have evolved due to the need to evaluate the  
212 | behavior of different soil attributes [22].

213 | Soil aggregation is one of the attributes used as indicators of soil quality, defined  
214 | as the ability to sustain agricultural productivity, maintain the quality of the environment,

215 | and ensure human, animal, and plant health [23]. Aand is related to essentialimportant  
216 | processes, such as erosion resistance and infiltration capacity [24].

217 | Soil erosion is one of the significantmajor environmental problems, because in  
218 | addition to soil and nutrient losses, it is associated with flooding, sedimentation, and  
219 | pollution of water bodies, and this process is affected by different factors such as soil  
220 | cover and management practices. However, soils with good aggregation are more  
221 | resistant to erosion [25, 17].

222 | Infiltration is also an important indicator of structuring and aggregation, influencing  
223 | the improvement of soil support capacity [26]. Besides that, their knowledge is  
224 | indispensable for the elaboration of an irrigation project, aimed at providing a greater  
225 | yield to the crops, and the better the aggregation, the highgreater the water infiltration  
226 | capacity [27].

227 | Another important aspect is the protection of soil organic matter, and its increase is  
228 | partially determined by the link between the recycling of macroaggregates, formation of  
229 | microaggregates and stabilization of carbon within the microaggregates. In order to  
230 | have a propergood formation and stabilization of these aggregates requires an  
231 | interaction of several factors such as, for example, soil fauna, roots, inorganic agents  
232 | and environmental variables [20].

233 | The organic compounds participate in the bonds between individual soil particles,  
234 | acting as cementing agents of the structural units by their diverse surface  
235 | characteristics, thus, there is a correlation between the organic matter and the stability  
236 | of the aggregates, since the organic compounds are the main cementing agents of the  
237 | soil particles and, at the same time, the state of greater aggregation promotes greater  
238 | physical protection of the organic matter of the soil thus allowing its accumulation [24,  
239 | 28].

240 | Cultural practices are primordial when optimum productivity is expected; besides  
241 | that, an inadequately performed activity can cause degradation of soil and natural  
242 | resources [29].

243 | Conventional preparation breaks the aggregates in the prepared layer and  
244 | accelerates the decomposition of the organic matter, reflecting negatively in the  
245 | resistance of the soil aggregates. Bertol et al. [30], evaluating the physical properties of  
246 | the soil under conventional tillage and direct sowing in rotation and succession of  
247 | crops, compared to the native ones, verified that the physical properties are altered  
248 | with the management, in which conventional cultivation resulted in a lower organic  
249 | carbon content, implying a greater soil degradation when compared to direct sowing.

250 | Studies made by Loss et al. [31] also observed when analyzing total organic  
251 | carbon and soil aggregation in an agroecological and conventional no-tillage system of  
252 | onion, that the use of single or intercropping cover crops in the planting system was  
253 | efficient to recover and increase the weighted average diameter (WMD), geometric  
254 | mean diameter (DMG) indices in relation to the conventional tillage system, in which  
255 | forage turnip increased the aggregation of the soil in the layer of 10-20 in relation to the  
256 | other treatments.

257 | The intensity of the structural stability of the aggregates varies according to the  
258 | type of soil and the cultural practices applied at the place of cultivation. When there is  
259 | soil rotation, the percentage of aggregates in the larger diameter classes reduces,  
260 | consequently, there is an increase in the class of smaller diameter, resulting in an  
261 | advers negative effect on the stability of the aggregates [18].

262 | In places arising from civil works, the common denominator of degraded areas is  
263 | the removal of the superficial horizon containing organic matter, causing serious  
264 | physical, chemical and biological problems to soil [26].

265 | An alternative to maintaining or recovering soil quality, is the usage of conservation  
266 | practices, as the no-tillage system, which, due to the absence of soil rotation and  
267 | maintenance of the straw on the surface, contributes to the improvement of soil  
268 | aggregation. Ssoil and consequently for the increase of carbon stocks in the soil, being

269 | more effective when associated [with](#) the use of cover crops, either by rotation or  
270 | succession of crops [32].

271 | According to Loss et al., (2014), analyzing the aggregation, light organic matter  
272 | and mineralizable carbon in soil aggregates, found that the conventional tillage system  
273 | reduced the aggregation index (WMD and DMG) and the organic matter content and  
274 | total organic carbon in relation to the forest area and using the direct tillage system and  
275 | pasture it was possible to recover these original values.

276

277

## 5. MECHANICAL RESISTANCE TO ROOT PENETRATION

278

279 | Soil compaction is an old problem and has been intensified with the expansion  
280 | of the agricultural frontier and the usage of basically two annual crops, mainly by use of  
281 | [more massive](#) heavier machinery and agricultural implements for the management of  
282 | soils and exploited crops [33,34]. Soil compaction refers to the compression of the  
283 | unsaturated soil during which there is an increase of its density because of the  
284 | reduction of its volume, resulting from the expulsion of air from the pores, causing a  
285 | denser rearrangement of the soil particles and consequent reduction of porosity [35].

286 | Thus, the increase of soil density becomes a limiting factor for the development  
287 | of the plants and, consequently, harming the achievement of higher yield indices  
288 | [36,37], due to the decrease of the water infiltration capacity [38], the low development  
289 | of the root system [39] due to the mechanical impedance, which results in a lower  
290 | volume of soil explored, a reduction in nutrient availability and losses of nitrogen by  
291 | denitrification [38], causing the increase of CO<sub>2</sub> and phytotoxins [40].

292 | The limitation to root growth, is clearly guided [41], within classes determined by  
293 | the values found in the resistance analysis (Mpa) as without limitation (<1,1); little  
294 | limitation (1.1 - 2.5); some limitations (2.6 - 5); serious limitations (5.5 - 10); roots  
295 | hardly grow (10,1-15); roots do not grow (> 15).

296 | The decomposition of a soil is done through the application of organic matter  
297 | into it in order to reduce its density, and green fertilization can be used [42,43], animal  
298 | manures, compost prepared on the farm, vegetable cakes and various industrial  
299 | wastes [15,44], among others.

300 | Several methods are used to recognize soil compaction, for example: trench  
301 | opening, vegetation cover visualization, soil density, and soil penetration resistance.

302 | The trench opening consists [of](#) the observation of the root system, especially  
303 | [about](#) in relation to subsurface compaction or grid footing. When there is subsurface  
304 | compaction, it is possible to observe a [significan](#) great concentration of roots in the  
305 | superficial layer, by not being able to cross the compacted layer [45].

306 | There is also the determination of soil density, which is the ratio between the  
307 | mass of a dry soil sample and the volume occupied by this sample, but the density  
308 | values may vary from soil to soil and difficult to correlate with plant growth [45].

309 | In order to solve this problem, it can use the relative density, which is the ratio  
310 | of the soil density to the maximum density, reached on the compacted sample in the  
311 | Procter test or [in](#) the uniaxial compression test. Hakansson and Lipiec [46] affirm that  
312 | the relative density isolates the effect of the texture in the density of the soil, being  
313 | possible to compare soils of different textures as the level of compaction.

314 | For determination of the resistance of soil to penetration, can be used  
315 | penetrometers or penetrometers, the penetrometer perform specific evaluations of  
316 | resistance to penetration, [pen](#)etrographs record the resistance throughout the soil  
317 | profile. Both equipments uses the same principle of operation, varying [the](#) only model,  
318 | having various types as the impact's ones, the torque spring and the prints that use  
319 | load cells [45].

320 | Soil compaction has become a global problem as a result of intensive  
321 | cultivation, increased use of heavy machinery, short crop rotations, and inadequate soil  
management practices [47, 48].

322 | The damages include both the compression as the shear of [the](#) structure of the  
323 | pores of the soil, so that simple indexes, as changes in the density of the soil, generally  
324 | provide an indicator of [lousy bad](#) damage to compaction [49, 50].

325 | Soil resistance and aeration are dynamic parameters mainly affected by soil  
326 | structure, texture, and water content. The interactions between water content and soil  
327 | density on soil resistance and aeration make it [challenging difficult](#) to characterize soil  
328 | compaction effects, considering individual soil properties [47].

329 | It is [essential important](#) to cultivate the soil with the correct humidity, so that  
330 | compaction is minimized [51]. As soil density increases and total porosity decreases,  
331 | soil resistance to root penetration increases, preventing root growth and restricting  
332 | water and air circulation throughout the profile resulting in poor aeration of the root  
333 | system [48].

334 | Intensive traffic in agricultural machinery is [standardcommon](#) in most  
335 | agricultural operations, even in no-tillage systems. Plowing, harvesting and spreading  
336 | chemicals or fertilizers are common operations on most farms. Most, when not all these  
337 | operations are carried out by heavy wheeled machines. Soil compaction by wheels is  
338 | characterized by a decrease in soil porosity located in the area below the wheel and  
339 | formation of grooves in the soil surface [51, 52, 53].

340 | The compaction's degree depends on the mechanical strength of the soil, which  
341 | is influenced by intrinsic properties of it, as texture and soil organic matter content;  
342 | structure of the plow layer on the wheel and its state of water; and loading, which  
343 | depends on axle load, tire size and speed, as well [as](#) tire solo interaction [51, 53].

344 | Increasing the pressure on the soil increases the chances of soil compaction.  
345 | Increasing the frequency of machine passes over a soil increases its bulk density and  
346 | cone index, resulting in soil compaction and [lowinadequate](#) soil physical conditions for  
347 | seed emergence. However, most of the total compaction of the soil is caused by the  
348 | first pass, or initial passages of the machine and [ten10](#) passages can affect the soil up  
349 | to 50 cm depth [53, 52].

350 | The depth of compaction varies widely from 10 to 60 cm but is more  
351 | [evidentobvious](#) in the surface soil (about 10 cm). Though, cone index increments  
352 | (penetrometer reading) between 16 and 76% may occur in the first 40 cm of the  
353 | surface layer, and the bulk density may also increase, but increases were limited to a  
354 | depth of 15 cm. However, in a pasture situation, differences between heavy and light  
355 | loads in the lower depth range (surface soil) were not found [51].

356 | Soil type also influences soil compaction. In soil with [a](#) thick texture, the  
357 | dominant stress penetration was in the vertical direction, while in thinner textured soil  
358 | the propagation of stress was multidirectional. However, they suggested that in soil  
359 | with good structure (aggregate soil) the compaction due to the axle load was not as  
360 | deep. The effects of axle load on soil compaction have been researched by many  
361 | workers around the world in the last decade [51].

362 | Animal trampling can cause compaction and degradation of soil structure. The  
363 | compaction caused by the grazing of animals through the action of the hull will  
364 | probably be more widespread in the pickets compared to the compaction caused by  
365 | mechanical implements that are limited under the rails. The trampling of the animals  
366 | [about in relation to](#) soil compaction can affect soil density, hydraulic conductivity,  
367 | macropore volume and resistance to soil penetration. The effects of grazing animals on  
368 | soil physical properties, nitrogen and soil carbon were discussed in detail in the  
369 | literature [53].

370 | Improved land management techniques are vital to ensure that [physical soil soil](#)  
371 | [physical](#) conditions are not compromised and practices that increase organic content,  
372 | reduce crop yield and sustain agricultural land use [51].

373 | Crops of coverage with aggressive and extensive root systems help in the  
374 | formation of soil aggregates, thus facilitating root growth of later crops and increased  
375 | water infiltration. Soil aggregation is generally improved by management systems,  
376 | including crops with a high capacity to form roots and increase soil organic matter. The

377 contribution of SOM to the formation of stable aggregates is attributed to processes  
378 such as the formation of cationic bridges, cementation between particles and stability  
379 promoted by root and microbial exudates around and within aggregates. Therefore, this  
380 could be a mechanism whereby the use of rotating hedge plants with the main crop  
381 would have a lasting effect on the alleviation of soil physical limitations [48].  
382

## 383 6. INFILTRATION OF WATER IN SOIL

384 Infiltration is a process by which water crosses the surface of the soil and  
385 redistributes in its profile. An essentialimportant process for the supply of underground  
386 aquifers, determining the water balance in the root zone of the crops, directly interfering  
387 in the runoff, responsible for erosion and flooding processes. The infiltration of water is  
388 a physical attribute sensitive to changes in soil planning, management, and  
389 conservation.

390 . The distribution of water in the soil profile, submitted to a hydraulic load on the  
391 surface, is distinguished in four respective zones to the increase of depth, according to  
392 Brandão [54].

393 Saturation zone: is located below the surface of the soil, usually a narrow layer, in  
394 which the soil is saturated.

395 Transition zone: layer characterized by a marked decrease of humidity.

396 Transmission zone: the region where the water is transmitted, characterized by  
397 increasing thickness with the continuous increase of application of water load, with a  
398 small variation of humidity aboutin relation to space and time.

399 Moistening zone: narrow layer, with significangreat reduction of humidity with  
400 increasing depth.

401 Moistening front is the visible limit of soil water movement, as a reflection of the  
402 variation of moisture exists in the system (soil), which is also affected by the physical,  
403 chemical and biological conditions of the soil.

404 The infiltration process has relations of dependence with some factors in which  
405 they can be divided into classes being, soil- related factors, surface related and soil  
406 preparation/ / management. These relations of dependence exert a function in the  
407 properties related to the porous space of soil composition, combined with the flowing  
408 fluid, determining the hydraulic conductivity, as well the occurrence of the surface  
409 crushing process caused by the impact of the raindrops on the soil, which allows the  
410 rearrangement of the particles, densification and consolidation of a surface structure,  
411 modifying the thickness of the surface layer [55].

412 Soils with a sandy (thick) texture, have a higher amount of macropores, when  
413 compared to clayey (fine) soils, in which they present higher hydraulic conductivity and  
414 infiltration rate, the contribution of the clay as an inorganic solid having loads is an  
415 excellengreat value for the structuring and aggregation of the soil.

416 The aggregation of soil particles contributes positively to the process of  
417 infiltration of water in the soil, besides promoting spaces to soil organisms. Infiltration is  
418 an important attribute that controls the leaching, flow, and availability of water to crops.  
419 Lack of residue coverage and direct exposure of soil to high- intensity rains result in  
420 poor aggregation, providing crust formation, as well reducing the availability of water to  
421 crops, contributing to poor water quality [56]

422 The type of soil surface cover is a determinant factor for the infiltration process,  
423 being responsible for the increase of the macroporosity of the surface layer, reduces  
424 surface crumbling, promotes a high infiltration potential and considerably reduces water  
425 and soil losses.

426 Water infiltration in the soil can contribute to a better understanding of the  
427 erosive dynamics, since the lower the infiltration rate, the high greater the possibility of  
428 surface runoff, reflecting the degree of soil compaction [57]

429 Studies by Marchini et al. [26] showed that the values of the infiltration rate  
430 ranged from 19.62 for exposed soil and 36.06 cm<sup>-1</sup> for Gonçalo Alves + Bean. The  
431 superiority of the treatment with vegetal cover can be explained, by the factors of soil

Formatted: No underline, Underline color: Auto

Formatted: No underline, Underline color: Auto

Formatted: No underline, Underline color: Auto

432 revolving, due to the preparation for the sowing, or by the effect of the roots of the  
433 green manure.

434 Influence of the factors related to the surface in the infiltration process was  
435 found by Bonini et al. [58], where the crop-livestock-forest system presented lower  
436 rates of water infiltration when compared to the eucalyptus forest and the crop-livestock  
437 system, this behavior can be attributed to the higher compaction of these systems,  
438 verified by the high values of resistance to root penetration.

439 Similar results were also observed by Marchão [39], where the crop-livestock  
440 system presented higher infiltration rates because of three main effects: absence of  
441 preparation during the grazing cycle, [the](#) presence of a dense root system and an  
442 increase in activity microbial and macrofauna of the soil.

443 The water infiltration process must be determined by simple methods with the  
444 potential to adequately represent the soil conditions [54].

445 In hydrological studies, infiltration rate determination equipment is used, with  
446 specific attributions, with the ring infiltrator, rainfall simulator and infiltrometer of mini-  
447 disk.

448

## 449 7. FINAL CONSIDERATIONS

450 Physical attributes reveal soil quality and indicate whether the management is  
451 appropriate. Attributes as soil mechanical resistance and water infiltration in the soil are  
452 fast and with low data acquisition costs. Already the porosity and density of the soil  
453 together with the aggregation, take time for the determination of the same and are  
454 costly. Analyzing soil attributes is extremely important for [excellentgood](#) productivity,  
455 since inappropriately used practices can influence plant development

456

## 457 CONFLICT OF INTEREST

458 The authors have no conflicts of interest to declare.

459

460

## 461 8. REFERENCES

462

463 1. Soares MDR, Campos MCC, Oliveira IA, Fonseca JS, Souza ZM. Physical attributes  
464 of the soil in areas under different systems of uses in the region of Manicoré, AM. Rev  
465 Cienc Agrar.2016; 59: 9-15.

466 DOI: 10.4322 / rca.2020.

467

468 2. Loss A, Junior ES, Schmitz D, Veiga M, Kurtz C, Comin JJ. Soil physical attributes in  
469 onion cultivation under no - tillage and conventional tillage systems. Revista  
470 Colombiana de Ciencias Hortícolas.2017; 11 (1): 105-113.

471 DOI: 10.17584 / rcch.2017v11i1.6144

472

473 3. Klein VA. Soil physics.3th ed. Passo Fundo: UPF .; 2014.

474

475 4. Teixeira W, Fairchild TR, Toledo MCM, Taioli F. Deciphering the Earth - 2nd ed. São  
476 Paulo: Companhia Editora Nacional .; 2009.

477

478 5. EMBRAPA. Irrigated Cotton Crops. 2003.

479 Available:

[http:](#)

480 //systemsproduction.cnptia.embrapa.br/FontesHTML/Algodao/AlgodaoIrrigado

481 (Accessed 10 March 2019)

482

483 6. Lorenzo, M. PEDOLOGIA - Morphology: Soil Porosity. 2010.

484 Available: [https://marianaplorenzo.com/2010/10/17/pedologia-%E2%80%93-](https://marianaplorenzo.com/2010/10/17/pedologia-%E2%80%93-morologia-porosidade-do-socio)  
485 [morologia-porosidade-do-socio](https://marianaplorenzo.com/2010/10/17/pedologia-%E2%80%93-morologia-porosidade-do-socio)

486 (Accessed 14 december 2018)

487  
488 7. Richart A, Filho JT, Brito OR, Llanillo RF, Ferreira R. Soil compaction: causes and  
489 effects. *Semina: Agrarian Sciences*.2005; 26: (3) 321-344  
490  
491 8. Kiehl EJ. *Manual of Edaphology*. São Paulo: Agronomic, 1979.  
492  
493 9. Luxmoore RJ. Micro, meso and macroporosity of soil. *Soil Science Society American*  
494 *Journal*. 1981; 45: 671-672.  
495  
496 10. Tognon AA. Physical-water properties of the Purple Latosol of the Guairá-SP  
497 region under different cropping systems. School of Agriculture of Luiz de Queiroz,  
498 Piracicaba, p. 85, 1991.  
499  
500 11. Hillel D. *Fundamentals of soil physics*. New York: Academic, p. 413, 1980.  
501  
502 12. Ribeiro KD, Menezes SM, MGBF Mosque, Sampaio FMT. Physical properties of  
503 the soil, influenced by the pore distribution, of six classes of soils of the region of  
504 Lavras - MG. *Science and agrotechnology*2007; 31: (4) 1167-1175.  
505 DOI: 10.1590 / S1413-70542007000400033  
506  
507 13. Lepsch IF. *Nineteen lessons of pedology*. São Paulo: Oficina de Textos, 2011. 456  
508 p.  
509  
510 14. Teixeira PC, Donagemma GK, Fontana A, Teixeira WG. *Manual of Methods of Soil*  
511 *Analysis*. 3rd ed. Brasília rev. ampl. : Embrapa. 2017; 574p.  
512  
513 15. Bonini CSB, AlvesMC. Physical quality of a Red Latosol recovering for seventeen  
514 years. *R. Bras. Eng. Agríc. Environmental*. 2012; 16 (4): 329-336.  
515  
516 16. Kormanek M, Głęb T, Banach J, Szewczyk G. Effects of soil bulk density on sessile  
517 oak *Quercus petraea* Liebl. *Eur J Forest Res*. 2015; 134 (6): 969-979.  
518 DOI: 10.1007 / s10342-015-0902-2  
519  
520 17. Salton JC, Tomazi M. Plant root system and soil quality. Press Release, Embrapa  
521 Agropecuária Oeste, 2014.  
522  
523 18. Prevedello J, Vogelmann ES, Kaiser DR, Fontanela E, Reinert DJ, Reichert JM.  
524 Aggregation and organic matter of an argisol under different soil preparation for  
525 Eucalyptus plantation. *Pesq. flower. bras*.2014; 34 (78): 149-158.  
526 DOI: 10.4336 / 2014.pfb.34.78.456  
527  
528 19. Tiecher T. Management and conservation of soil and water in small rural properties  
529 in southern Brazil: alternative management practices aiming at soil and water  
530 conservation. Porto Alegre, UFRGS, p. 186, 2016.  
531  
532 20. Arcângelo L, Costa EM, Pereira MG, Beutler SJ. Aggregation, light organic matter  
533 and mineralizable carbon in soil aggregates. *Revista de la Facultad de Agronomía, La*  
534 *Plata*.2014; 113 (1): 1-8.  
535  
536 21. Beutler AN, Munareto JD, Greco AMF, Pozzebon BC, Galon L, Guimarães S, Burg  
537 G, Schmidt MR, Deak EA, Giacomeli R, Alves GS. Soil management, residual straw  
538 and flooded rice yield. *Semina: Agrarian Sciences*.2014; 35 (3): 1153-1162.  
539 DOI:10.5433/1679-0359.2014v35n3p1153  
540

541 22. Mantovanelli BC, Silva DAP, Campos MCC, Gomes RP, Soares MD, Santos LAC.  
542 Evaluation of soil attributes under different uses in the region of Humaitá, Amazonas.  
543 Rev. Cienc. Agrar.2015; 58 (2): 122-130.  
544 DOI: 10.4322 / rca.1822  
545

546 23. Parron LM, Garcia JR, Oliveira EB, Brown GG, Prado RB. Environmental Services  
547 in Agricultural and Forest Systems of the Atlantic Forest Biome. Embrapa Forests,  
548 2015.  
549

550 24. Borges, C.A .; Ribeiro, B. T .; Wendling, B .; Cabral, D. A. Soil aggregation, organic  
551 carbon and CO2 emission in areas under different uses in the Cerrado, Triângulo  
552 Mineiro region. Rev. Ambient. Water.2015; 10 (3): 660-675.  
553 DOI: 10.4136 / ambient-water.1573  
554

555 25. Almeida WS, Carvalho DS, Panachuki E, Valim WC, Rodrigues SA, Varella CAA.  
556 Water erosion in different cropping systems and levels of soil cover. Pesq. agropec.  
557 bras.2016; 51 (9): 1110-1119.  
558 DOI: 10.1590 / S0100-204X2016000900010  
559

560 26. Marchini DC, Ling TC, Alves MC, Crestanha S, Filho SNS, Arruda OG. Organic  
561 matter, infiltration and tomographic images of Oxisol undergoing recovery under  
562 different types of management. R. Bras. Eng. Agríc. Environmental.2015; 19 (6): 574-  
563 580.  
564 DOI: 10.1590 / 1807-1929 / agriambi.v19n6p574-580.  
565

566 27. Neto GS, Santos KP, Fernandes GAG, Oliveira CF. Determination of the equation  
567 and infiltration velocity curve of red latosol water. 5th Academic Seminar - Science,  
568 Innovation and Technology in the Closed Biome.2011; 5 (1).  
569

570 28. Portugal AF, Juncksh I, Schaefer CERG, Neves JCL. Stability of aggregates in  
571 argisol under different uses, compared to forest. Rev. Ceres.2010; 57 (4): 545-553.  
572

573 29. Loss A, Pereira MG, Giacomo SG, Perin A, LHC Angles. Aggregation, carbon and  
574 nitrogen in soil aggregates under no-tillage with crop-livestock integration. Pesq.  
575 agropec. bras.2011; 46 (10): 1269-1276.  
576

577 30. Bertol I, Albuquerque JA, Leite D, Amaral AJ, Zoldan JWA. Physical properties of  
578 the soil under conventional tillage and direct seeding in rotation and succession of  
579 crops, compared to the native field. R. Bras. Ci. Solo.2004; 28 (1): 155-163.  
580 DOI: 10.1590 / s0100-06832004000100015.  
581

582 31. Loss A, Basso A, Oliveira BS, Koucher LP, Oliveira RA, Kurtz C, Lovato PE, Curmi  
583 P, Brunetto G, Comin JJ. Total organic carbon and soil aggregation in conventional  
584 agroecological and conventional onion system. R. Bras. Ci. Solo.2015; 39 (4): 1212-  
585 1224.  
586 DOI: 10.1590 / 01000683rbcs20140718  
587

588 32. Loss A, Costa EM, Pereira MG, Beutler SJ. Aggregation, light organic matter and  
589 mineralizable carbon in soil aggregates. Journal of The Faculty of Agronomy, La Plata,  
590 2014; 113 (1): 1-8.  
591

592 33. Pedrotti A, Dias JMS. Soil compaction: how to avoid it. Agropecuária  
593 Catarinense.1996; 9 (4): 50-52.  
594

- 595 34. Reichert JM, Suzuki LEAS, Reinert DJ. Soil compaction in agricultural and forest  
596 systems: identification, effects, critical limits and mitigation. *Topics Ci. Solo*.2007; 5 (1):  
597 49-134.  
598
- 599 35. Dias JMS, Pierce RJ. The process of soil compaction and its modeling. *Brazilian*  
600 *Journal of Soil Science*. 1996; 20 (2): 175-182.  
601
- 602 36. Alakukk L, Elomen P. Long-term effects of a single compaction by heavy field traffic  
603 on yield and nitrogen uptake of annual crops. *Soil & Tillage Research*.1994; 36 (3-4):  
604 141-152.  
605 DOI: 10.1016 / 0167-1987 (95) 00503-X  
606
- 607 37. Cavichiolo SBV, Dedecek RA, Gava JL. Modifications in the physical attributes of  
608 soils submitted to two systems of preparation in regrowth of *Eucalyptus equine*. *R.*  
609 *Árvore*.2005; 29 (4): 571-577.  
610
- 611 38. Oliveira VS, Rolim MM, Vasconcelos RF, Costa YD, Pedrosa, E. M. Compaction of  
612 a ultisol submitted to different managements. *Rev. bras. eng. Agríc*.2010; 14 (9): 914-  
613 920.  
614 DOI: 10.1590 / S1415-43662010000900002  
615
- 616 Physical quality of a Red Latosol under systems of integrating livestock farming in the  
617 Cerrado, Brazil. *Pesq. agropec. bras* 2007; 42 (): 873-82.  
618 DOI: 10.1590 / S0100-204X2007000600015  
619
- 620 40. Marschner H. Mineral nutrition of higher plants. 2th ed. San Diego, Academic Press,  
621 p. 889, 1995.  
622
- 623 41. Canarache A. Penetr-a generalized semi-empirical model estimating soil resistance  
624 to penetration. *Soil & Tillage Research*. nineteen ninety; 16 (1-2): 51-70.  
625 DOI: 101016 / 0167-1987 (90) 90021-5  
626
- 627 42. Kitamura AE, Alves MC, Suziki LGAS, Gonzalez AP. Recovery of degraded soil  
628 with the application of green manures and sewage sludge. *Rev. Bras. Ciênc. Solo*.  
629 2008; 32 (1): 405-416.  
630 DOI: 10.1590 / S0100-06832008000100038  
631
- 632 43. EMBRAPA - Brazilian Agricultural Research Corporation. Ministry of Agriculture,  
633 Livestock and Supply. Technical information for common bean cultivation in Central-  
634 Brazilian region: 2012-2014. 1ed.Santo Antônio de Goiás-GO, p. 248, 2012.  
635
- 636 44. Bonini CSB, Alves MC, Montanari R. Recovery of the structure of a degraded red  
637 Latosol using sewage sludge. *Brazilian Journal of Agricultural Sciences / Brazilian*  
638 *Journal of Agricultural Sciences* 2015; 10 (1): 34-42.  
639 DOI: 10.5039 / agraria.v10i1a4513  
640
- 641 45. Sá MAC, Santos JJDG. Soil compaction: consequences for plant growth.  
642 *Documents* 136, Embrapa, p. 24, 2005.  
643
- 644 46. Hakansson I, Lipiec JA. review of the usefulness of relative bulk density values in  
645 studies of soil structure and compaction. *Soil & Tillage Research*. 2000; 53(2):71- 85.  
646 DOI:10.1016/S0167-1987(99)00095-1  
647
- 648 47. Chen G,Weil RR,Hill RL. Effects of compaction and cover crops on soil least  
649 limiting water range and air permeability. *Soil & Tillage Research*.2014; 136(2):61-69.

650 DOI:10.1016/j.still.2013.09.004  
651  
652 48. Calonego JC,Rafhael JPA,RigonJPG,Neto LO,Rosolem CA. Soil compaction  
653 management and soybean yields with cover crops under no-till and occasional  
654 chiseling. *Europ. J. Agronomy*.2017; 85(5):31-37.  
655 DOI:10.1016/j.eja.2017.02.001  
656  
657 49. Chamen WT, Moxey AP, Towers W, Balana B, Hallett PD. Mitigating arable soil  
658 compaction: A review and analysis of available cost and benefit data. *Soil & Tillage  
659 Research*. 2015; 146(2):10-25.  
660 DOI:10.1016/j.still.2014.09.011  
661  
662 50. Alaoui A,Rogger M,Peth S,Blöschl G. Does soil compaction increase floods? A  
663 review. *Journal of hydrology*, 2017;557 (2), 631-642.  
664 DOI:10.1016/j.jhydrol.2017.12.052  
665  
666 51.Hamza MA,Anderson WK. Soil compaction in cropping systems: A review of the  
667 nature, causes and possible solutions. *Soil & Tillage Research*.2005; 82(2):121-145.  
668 DOI:10.1016/j.still.2004.08.009  
669  
670 52. Lima RP, León MJ, Silva AR. Soil compaction of different textured classes in areas  
671 of sugarcane production. *Rev. Ceres*.2015; 60 (1): 16-20.  
672 DOI: 10.1590 / S0034-737X2013000100003  
673  
674 53. Nawaz MF, Bourrie G, Trolard F. Soil compaction impact and modeling. A review.  
675 *Agronomy for sustainable development*. 2013; 33 (2): 291-309.  
676 DOI: 10.1007 / s13593-011-0071-8  
677  
678 54. Brandão VS, Cecílio RA, Pruski FF, Silva DD. *Infiltration of Soil Water*. 3rd ed.  
679 current and ampl. Viçosa: Editora UFV, 2012.  
680  
681 55. Valentin C, Bresson LM. Morphology, genesis and classification of surface crusts in  
682 loam and sandy soils. *Geoderma*.1992; 55 (3-4): 225-245.  
683 DOI: 10.1016 / 0016-7061 (92) 90085-L  
684  
685 56. Franzluebbers AJ. Water infiltration and soil structure related to organic matter and  
686 its stratification with depth. *Soil & Tillage Research*.2002; 66 (2): 197-205.  
687 DOI: 10.1016 / S0167-1987 (02) 00027-2  
688  
689 57. Mathias DT, Lupinacci CM, Moruzzi RB. Distribution of infiltration rates in areas  
690 affected by accelerated erosive processes and covered by anthropogenic materials.  
691 *Sanitary and Environmental Engineering*.2018; 23 (5): 1-9.  
692 DOI: 10.1590 / S1413-41522018167570  
693  
694 58. Bonini CSB, Lupatini GC, Andrighetto C, Mateus GP, Heinrichs R, Aranha AS,  
695 Santana EAR, Meirelles GC. Fodder production and soil chemical and physical  
696 attributes in integrated agricultural production systems. *Pesq. agropec. bras*.2016; 51  
697 (9): 1695-1698.  
698 DOI:10.1590/s0100-204x2016000900070  
699  
700  
701