

1
2
3
4
5
6
7
8
9
10

Original Research Article

Control of *Alternaria alternata* using Melaleuca essential oil (*Melaleuca alternifolia*)

ABSTRACT

Aims: This study aimed to evaluate the fungitoxic potential of Melaleuca essential oil on the mycelial growth of *Alternaria alternata* under *in vitro* condition and the treatment of cowpea beans.

Study desing: The experiments comprised completely randomized designs: Eleven treatments with five replicates on *in vitro* test; and six treatments with five replicates on *in vivo* test.

Place and Duration of Study: The work was carried out at the Center for Agrifood Science and Technology of the Federal University of Campina Grande, Pombal, Brazil, ~~between~~ since February 2018 to February 2019.

Methodology: In the *in vitro* experiment, the essential oil was incorporated into the culture medium and poured into Petri dishes. The treatments consisted of different concentrations of the essential oil (0.0125, 0.025, 0.05, 0.1, 0.2, 0.25, 0.50, 0.75, and 1.0%), a negative control (0.0%), and a positive control (Thiram). Discs of culture medium with fungal mycelia were inoculated in the center of the plates and incubated for seven days at 27±2°C. We calculated the percentage of mycelial growth inhibition (PGI) and the index of mycelial growth speed (IMGS) to verify the difference between treatments. In the *in vivo* experiment, the bean seeds were treated with different concentrations of oil (0.0, 0.2, 0.5, 1.0, and 5.0%), a negative control (0.0%), and positive control (Thiram). Seeds were inoculated with colonies of the fungus for 48 hours, and after that, we performed the seed sanity test.

Results: Under *in vitro* conditions, all concentrations of melaleuca essential oil reduced the mycelial growth of *A. alternata*. The oil reached complete inhibition of fungal growth from 0.2% concentration and above. In the cowpea treatment, the essential oil had no significant control over the percentage of infected seeds.

Conclusion: The melaleuca essential oil had a fungitoxic effect on the *A. alternata* under *in vitro* conditions. However, using the adopted methodology, on the cowpea bean seed treatment, the essential oil had was not reduce the incidence of *A. alternata*.

Keywords: Alternative control, Cowpea bean, Mycelial growth, Phytopathogenic fungi, Tea-tree, Seeds disease unguiculata.

1. INTRODUCTION

Cowpea bean (*Vigna unguiculata* (L.) Walp), popularly known as the string bean or macaçar bean, is a source of protein and staple food for a large part of the population of the North and Northeast of Brazil, thus one of the most important crops in the country [1]. According to CONAB [2], Brazil occupies the third position in world bean production with a cultivation area of approximately one million hectares, with the North and Northeast regions accounting for about 90% of the cultivated area [3].

Comment [R1]: All names in latin should be in italics all over the manuscript.

Comment [R2]: Using just «oil» may confuse reders. 'Oil' intuitively associates with grease. Please always use 'essential oil' ore abbreviation 'EO' all over manuscript.

Comment [R3]: Please use in whole manuscript impersonal form. For instance, in this case I suggest: 'The percentage of mycelial growth inhibition (PGI) was calculated...'

Comment [R4]: Please correct the grammar.

Comment [R5]: Please consider 'Tea-tree essential oil'.

11
12

13
14
15
16
17
18
19
20
21
22
23
24

Cowpea cultivation has a very competitive production cost, a factor that has increased the farmers' interest in the crop. In addition, Brazilian production is of high quality, enabling the product to have good acceptance in all members of its production chain [4]. However, diseases represent a limiting factor to income expansion.

Fungi are the main phytopathogens that cause economic losses in bean crop. When present in the seed, they can cause miscarriages, deformations and discoloration of the bark, which always leads to the reduction of seed germination potential and vigor, and when allocated in the field will result in low or no yielding uneven plant stands [5]. Diseases caused by fungi with the greatest economic impact on bean crop are caused by *Macrophomina phaseolina* [6], *Fusarium* spp. [7], *Rhizoctonia solani* [8], *Curvularia* spp., *Trichoderma* spp. [9], *Alternaria* spp. [10], *Aspergillus* sp. and *Penicillium* sp. [11].

Considering bean pathogen control practices, treatment with synthetic agrochemicals has been a conventionally used one. However, the use of these products has been associated with significant damage to human health and the environment due to their high toxicity [12,13] besides favoring the emergence of resistant strains [14].

In this scenario, it is necessary to use alternative products to chemical pesticides that have similar efficacy but are not harmful to human health and the environment. Among the products studied are essential oils extracted from aromatic plants, which have fungitoxic properties on phytopathogens [15,16,17].

Melaleuca essential oil (*Melaleuca alternifolia*) has been studied for some years and its antimicrobial activity has been well documented. The main components of this oil are: terpineol, cineol, terpenene, cymene, limonene and sabinene [18]. Most compounds have inhibitory activity against fungi and bacteria [19]. Among these, terpineol is the main antifungal constituent [20]. In the control of phytopathogens its use has shown promising results in the control of fungi *Cercospora beticola* [21], *Aspergillus niger*, *M. phaseolina*, *Penicillium* sp. and *Sclerotinia sclerotiorum* [22], demonstrating a strong antimicrobial activity.

The antifungal action of oils is related to their ability to dissolve in lipid media, causing modifications in the cell membrane structure [23]. Due to their low toxicity and rapid degradation in the environment, the use of essential oils to combat phytopathogens may be a good alternative to synthetic pesticides [24]. Thus, this work aimed to evaluate the fungitoxic potential of melaleuca essential oil on the mycelial growth of *Alternaria alternata* under *in vitro* conditions and in the treatment of cowpea seeds.

2. MATERIAL AND METHODS

2.1 Place of experiments

The work was conducted at the Center of Science and Technology Agrifood (CCTA) of the Federal University of Campina Grande (UFCG), Campus of Pombal. The experiments were carried out in the Phytopathology laboratory, between February 2018 to and February 2019.

2.2 Sampling

We used the fungal strain of *Alternaria alternata* 0878 yielded by the collection of phytopathogenic fungi Prof. Maria Menezes of the Federal Rural University of Pernambuco (UFRPE). The fungi were preserved in sterile distilled water by the Castellani method until the assay [25].

The pure essential oil of Melaleuca (*Melaleuca alternifolia*) was purchased at a local store specialized in natural products. The cowpea bean seeds (*Vigna unguiculata* L. Walp) were purchased at a commercial house in the city of Patos, Paraíba.

2.3 Screening of the antifungal activity of Melaleuca essential oil in vitro

Eleven treatments were used, 9 oil concentrations (0.0125, 0.025, 0.1, 0.2, 0.5, 0.25, 0.50, 0.75 and 1.0%), a negative control (without essential oil supplementation=0.0%) and a positive control (supplemented with 1 mL L⁻¹ of the fungicide Thiram, which is the dosage indicated by the manufacture's). Five replicates of each treatment were arranged in completely randomized design (CRD).

The treatments were incorporated into PDA (Potato Dextrose Agar) culture medium just before pouring in sterilized Petri dishes. After solidification, one-centimeter mycelial disks were taken from the margins of 7days old culture and transferred to the center of each plate containing the treatments. The plates were then wrapped in plastic film and incubated in a BOD (Biochemical Oxygen Demand) at a temperature of 27±2°C.

Comment [R6]: I suggest 'terpine-4-ol' here and further.

Comment [R7]: Please combine this two sentences. Also please check the rest of manuscript to avoid a lot of short sentences. Then the text will be easier to read.

Formatted: Highlight

86 The concentrations were chosen from an initial concentration based on the literature [26,27] and then gradually reduced
87 until the addition of oil to the medium was no longer able to prevent the fungal growth. To obtain the final concentrations,
88 the direct dilution procedure in a culture medium [28] was used.
89

90 Colony growth was measured daily until the colony took the entire surface of the culture medium in one of the plates or in
91 a maximum period of 7 days. Mycelial growth evaluation consisted of daily measurements of the diameter of the colonies
92 obtained through the average of two perpendicular measurements, using a digital caliper, resulting in the average daily
93 growth for each repetition of each treatment.
94

95 The percentage of mycelial growth inhibition (PGI; [29]) and mycelial growth rate index (IMGS; [30]) were calculated
96 according to formulas (1) and (2):
97

$$98 \quad PGI = \frac{[(negative\ control\ growth - treatment\ growth)] \times 100}{negative\ control\ growth} \quad (1)$$

$$100 \quad IMGS = \sum \frac{current\ mycelial\ growth - previous\ mycelial\ growth}{number\ of\ days\ of\ incubation} \quad (2)$$

102 The minimum inhibitory concentration was considered the lowest oil concentration capable of totally inhibiting *Alternaria*
103 *alternata* mycelial growth.
104

105 **2.4 Screening of the antifungal activity of Melaleuca essential oil *in vivo* (on cowpea bean seeds)**

106 The experiment consisted of a completely randomized ~~desingdesign~~. The treatments consisted of sterilized distilled water
107 solutions supplemented with 4 oil concentrations (0.2, 0.5, 1.0 and 5.0%), a negative control (without essential oil
108 supplementation=0.0%) and ~~and~~ a positive control (supplemented with 1 ml L⁻¹ of the fungicide Thiram, which is the
109 dosage indicated by the manufacture's). The concentrations used were determined based on the *in vitro* test. To emulsify
110 the oin in the water Tween 80 (1 mL L⁻¹) was used [31].
111

112 The cowpea bean seeds were ~~desinfected~~ ~~disinfected~~ in 2.0% sodium hypochlorite solution for five minutes, washed with
113 sterile distilled water twice and dried at room temperature. Afterwards they were immersed for five minutes in different
114 solutions (treatments). After drying at room temperature, the artificial inoculation was performed.
115

116 The inoculation was done depositing the seeds on 7 days colonies of *A. alternata* ~~with 7 days of age~~. The seeds and the
117 fungal colonies stayed for 48 hours in a BOD 27±2°C, with a 12-hour photoperiod [32].
118

119 After the treatment and inoculation, the samples were submitted to the sanity test, which was performed by the filter paper
120 method with freezing [33]. Six hundred of cowpea bean seeds (100 per treatment) were used, distributed in Petri dishes
121 (Ø=14 cm). In this method, ten seeds were placed at equal distances on each plate on triple layer of filter paper previously
122 moistened in sterile distilled water and incubated initially for 24 hours on BOD at 27±2°C, with a 12-hour photoperiod.
123 After this period, they were subjected to freezing (-20°C) for 24 hours, and then returned to the incubator for another five
124 days.
125

126 After incubation, the seed were examined individually, using a stereoscopic microscope, for the quantification of seeds
127 infected by *Alternaria alternata*. The results were expressed as percentage of infected seeds.
128

129 **2.5 Statistical analysis**

130 The effect of oil concentration on fungal growth was analyzed using regressions in quadratic plateau model for *in vitro*
131 experiment and in linear model for *in vivo* experiment.
132

133 To test the difference between treatments with the essential oil and the treatment containing the fungicide (positive
134 control), Mann-Whitney (Tukey nonparametric) multiple comparisons were applied. Non-parametric tests were used
135 because of the lack of variance in the results of some treatments. Differences with a probability values below 5% were
136 considered significant. The regressions were performed in the program R CoreTeam 3.5.1[34].
137

138 **3. RESULTS AND DISCUSSION**

139 **3.1 *In vitro* antifungal assay**

Comment [R8]: Are there available any pictures of Petri dishes with *in vitro* assay results? It will be a very enriching addition to this manuscript.

3.1.1 Effects of Melaleuca essential oil on *Alternaria alternata*

All tested concentrations of melaleuca essential oil reduced the mycelial growth and growth speed of *Alternaria alternata*. The inhibition percentages increased significantly with the concentrations ($P < .001$) reaching the maximum value (PGI=100%) the 0.2% concentration of the oil (Fig 1A), which is the minimum inhibitory concentration (MIC). On the other hand, applying the regression equation in a quadratic plateau model, the estimated minimum concentration (MCest) was 0.33%.

The mycelial growth rate is a variable inversely proportional to the inhibition percentage. For this reason, it presented opposite behavior, with significant reduction with the tested oil concentrations ($P < .001$). The mycelial growth rate was more effectively reduced from the 0.2% concentration, in which growth paralyzed (IMGS=0.00 cm day⁻¹) (Fig 1B), differing from the negative control, which presented the highest growth speed (0.63 cm day⁻¹).




Fig 1. Inhibition percentage and mycelial growth speed of melaleuca essential oil against *Alternaria alternata*.

***Level of significance below 0.1% ($P < .001$)

According to the literature, terpinenol (terpinen-4-ol) is the major constituent of the melaleuca essential oil, which is associated with your high fungitoxic potential [35]. One of the antifungal mechanisms of action of melaleuca essential oil is the change in the permeability and fluidity of the cell membranes of the microorganisms. As these organisms are permeable to oil, the main effects found are inhibition of cell respiration and alteration in membrane structure and integrity, as well as leakage of essential intracellular materials. These events cause growth inhibition or even cell death [36,37].

Using tea tree oil at concentrations close to or greater than ours, other authors obtained similar inhibition results. For example, Martins et al. [22] obtained total inhibition of *Macrophomina phaseolina* and *Sclerotinia sclerotiorum* at concentration 0.2%. While in the control of *Alternaria radicina* and *A. dauci*, Ricioni and Orzali [38] reached the maximum inhibition from the concentration 0.5%.


Using the essential oil of the other plant species on control of *A. alternata*, other authors obtained similar results as superior or inferior to ours. For example, the total inhibition was achieved by Chutia et al. [39], Guimarães et al. [40] and Barboza [41] using mandarin orange (*Citrus reticulata*), lemongrass (*Cymbopogon citratus*) and alecrim-da-chapada (*Lippia gracilis*) essential oil at concentrations of 0.2 mL/100mL (0.2%), 14.49 µg mL⁻¹ (0.0014%) e 750 µL L⁻¹ (0.075%),

Comment [R9]: Please be consistent. In line 32 you had used *Macrophomina phaseolina* full name. In line 48 shorter version – *M. Phaseolina*. I suggest to use full name of microorganism when first time mentioned in the text, and later shorter version. Since *Sclerotinia sclerotiorum* was used in line 48, now it should be '*S. Sclerotiorum*'. Etc.

Comment [R10]: All numbers should be with full stop (.) not with coma (,).

179 respectively. On the other hand, using peppermint essential oil (*Mentha piperita*), França et al. [42] obtained a maximum
 180 inhibition of 41.6% at a concentration of 0.8%. Thus, both the fungitoxic potential of essential oils on *A. alternata*, as well
 181 as their minimum inhibitory concentrations will vary depending on the plant species from which the essential oil was
 182 extracted [43]. In addition, increasing inhibitory power as a function of increased concentration can either potentiate the
 183 effect or generate product waste.

184
 185 To understand the potential of melaleuca essential oil as a fungicide on *A. alternata*, we compared its fungitoxic effect with
 186 that obtained by the fungicide Thiram (commercial synthetic fungicide). We observed strong inhibition effect of the oil
 187 concerning the fungicide from the concentration of 0.2% (Fig 2). This result suggests that under *in vitro* conditions the oil
 188 could replace the use of this agrochemical.



190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212 **Fig 2. Effect of different treatments (melaleuca essential oil and the control treatments) on the mycelial growth**
 213 **inhibition of *Alternaria alternata*.**


214 Superscript concentrations with the same letter were not significantly different from each other by the Mann-Whitney test ($P > .05$)

215
 216 Due to the chemical complexity, the antifungal control promoted by essential oils is associated with their different
 217 constituents [44] through different mechanisms of action that act simultaneously on different targets [15]. These peculiar
 218 characteristics guarantee the advantage over synthetic fungicide, since they reduce the possibility of resistance by
 219 phytopathogens [45].

221 3.2 *In vivo* antifungal assay

222 3.2.1 Effects of Melaleuca essential oil on cowpea beans seed infected with *Alternaria alternata*

223
 224
 225 Contrary to that observed on the *in vitro* test, in cowpea bean seed treatment, the melaleuca essential oil was ineffective
 226 in combating the incidence of *A. alternata*. Increasing concentrations did not reduce the number of seeds infected by the
 227 phytopathogen ($P < .001$; Fig 3).



228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000


240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263

Fig 3. Effect of concentrations of melaleuca essential oil in the incidence of infected cowpea bean seeds by *Alternaria alternata*.

The incidence of infected seeds at all oil concentrations was higher than the negative control (treatment without the addition of oil). One of the hypotheses raised by the authors is that the essential oil did not adhere to the seed surface due to the high volatilization of its constituents. Thus, during the incubation period some constituents may have evaporated and reduced to their antimicrobial capacity.

Khalili et al. [46] emphasize that the formation of oils by volatile compounds and their subsequent degradation may be influenced by ambient temperature. And according to Simões and Spitezer [47] and Rozwalka et al. [48], the volatilization of oil constituents as well as their instability in the presence of light, heat and humidity, modify the atmosphere inside the Petri dishes, leading to the loss of the effectiveness of an oil that, under other conditions, inhibited fungal growth.

The effect of melaleuca essential oil on the incidence of infected seeds was lower than Thiram fungicide at all concentrations tested (Fig 4). Between 0.2 and 1%, the oil did not differ significantly from the negative control ($P > .05$), while at the 5% concentration, the oil promoted an increased incidence of phytopathogen in relation to the control. On the other hand, the treatment containing the Thiram fungicide prevented the development of phytopathogen in the seeds.



Comment [R12]: There is a lack of discussion regarding the increased incidence of seeds, when treated with 5% EO concentration. This result is really surprising and controversial. It is needed to bring forward some ideas that will explain that. Maybe such high EO concentration has affected the surface of the seeds, what made easier pathogen to colonize it?

264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281

Fig 4. Percentage of infected cowpea bean seeds by *Alternaria alternata* after the treatment with different concentrations of melaleuca essential oil and the control treatments.

In the present study, although the melaleuca essential oil did not provide satisfactory results in the *in vivo* experiment, against other phytopathogens the use of essential oils in the treatment of bean seeds was successful. In the treatment of carioquinha bean seeds. Using carioquinha bean seeds treated with lemongrass (*Cymbopogon flexuosus* and *C. citratus*) and melaleuca (*Melaleuca* sp.) essential oils, Morais et al. [49] obtained a significant reduction in the incidence of seeds infected by *Aspergillus* sp. and *Penicillium* sp. Whereas, Wanderley et al. [50] proved the efficacy of citronella

(*Cymbopogon* sp.) fennel (*Pimpinella anisum*) and alfavaca (*Ocimum basilicum*) essential oils at a concentration of 1.5% over *Callosobruchus maculatus* in butterbean seeds.

Comment [R13]: Please rewrite this paragraph. It is strongly unclear.

Finally, despite the ineffectiveness of melaleuca essential oil in the treatment of cowpea bean seed, this oil may be useful in the treatment of other seeds and other pathogens. Essential oils present a low risk to the environment, producers and consumers, and hinder the development of pathogen resistance [51]. Thus, further studies on the use of these oils in the management of plant pathogens are needed to make them a viable alternative for farmers.

4. CONCLUSIONS

Under *in vitro* conditions, melaleuca essential oil (*Melaleuca alternifolia*) totally inhibited the mycelial growth of *Alternaria alternata* from 0.2%, had an similar effect to the commercial fungicid Thiram. On the cowpea bean seed treatment, the essential oil had was not able to reduce the incidence of *A. alternata* using the adopted methodology.

Comment [R14]: I very like this part. Maybe you may add some ideas to improve tea-tree EO influence in further studies.

REFERENCES

1. Neves AC. et al. Cultivo do feijão-caupi em sistema agrícola familiar. Teresina- PI: Embrapa Meio Norte, 2011:15. (Embrapa Meio Norte. Circular Técnica, 51). Portuguese.
2. CONAB - Companhia Nacional de Abastecimento. Perspectiva para a Agropecuária. Brasília. 2018;6:1-112. Portuguese.
3. Santos LAC, Silva DMP, Oliveira IA, Pereira CE, Campos MCC. Crescimento de cultivares de feijão-caupi em solo de terra firme e várzea. *Ambiência Guarapuava*. 2017;13(1):261-270. Portuguese.
4. Freire Filho FR. Feijão-caupi no Brasil: Produção, melhoramento genético, avanços e desafios. Embrapa Meio-Norte, Teresina. 2011:84. Portuguese.
5. Machado JC. Tratamento de sementes no controle de doenças. Lavras: LAPS/UFLA/FAEPE. 2000:13.
6. Athayde Sobrinho C, Viana FMP, Santos A. Doenças fúngicas e bacterianas. In: Freire Filho FR, Lima JAA, Ribeiro VQ. Feijão-caupi: avanços tecnológicos. (EDIÇÃO) Brasília, DF: Embrapa Informação Tecnológica, 2005, capítulo 12, p.461-484.
7. Rodrigues AAC, Menezes M. Detecção de fungos endofíticos em sementes de caupi provenientes de Serra Talhada e de Caruaru, estado de Pernambuco. *Fitopatologia Brasileira*. 2002;27(5):532-537. Portuguese.
8. Nechet KL, Halfeld-Vieira BA. Caracterização de isolados de *Rhizoctonia* spp., associados à mela do Feijão-Caupi (*Vigna unguiculata*), coletados em Roraima. *Fitopatologia Brasileira*. 2006;31:505-508.
9. Sinha A, Singh SK, Qaisar J. Seed mycoflora of French bean and its control by means of fungicides. *Tropenlandwirt*. Witzzenhausen. 1999;11(1):59-67.
10. Marques RO, Alves VM, Lima MLP, Arruda GMT. Avaliação sanitária e fisiológica de feijão oriundos de Unaí – MG, Paracatu-MG e Cristalina-GO. *Summa Phytopathologica*. 2006;32(Suplemento). Portuguese.
11. Torres SB, Bringel JMM. Avaliação da qualidade sanitária e fisiológica de sementes de feijão macassar. *Caatinga*. 2005;18(2):88-92. Portuguese.
12. Candido e Silva EK, Melo LGL. Manejo de doenças de plantas: Um enfoque agroecológico. *Revista EDUCamazônia–Educação Sociedade e Meio Ambiente*. 2013;10(1):143-157. Portuguese.
13. Perina FJ. Óleos essenciais e frações majoritárias ativas no controle da mancha marrom de alternaria (*Alternaria alternata*) em tangerina ponkan. 2014. 112f. Tese (Doutorado em Agronomia). Universidade Federal de Lavras, Lavras, MG, 2014. Portuguese.
14. Hillen T, Schwan-Estrada KRF, Mesquinl RM, Cruz MES, Stangarlin JR, Nozaki M. Atividade antimicrobiana de óleos essenciais no controle de alguns fitopatógenos fúngicos *in vitro* e no tratamento de sementes. *Revista Brasileira de Plantas Mediciniais*. 2012;14(3):439-445. Portuguese.

- 342 15. Abdel-Kader MM, El-Mougy NS, Lashin SM. Essential oils and *Trichoderma harzianum* as an integrated control
343 measure against faba bean root rot pathogens. *Journal of Plant Protection Research*. 2011;51(3):306-311.
344
- 345 16. Silva M, Souza HRT, David HMSS, Santos LM, Silva RF, Amaro HTR. Qualidade fisiológica e armazenamento de
346 sementes de feijão-comum produzidas no norte de Minas Gerais. *Revista Agro Ambiente*. 2014;8:97-103. Portuguese.
347
- 348 17. Santos MM, Peixoto AR, Pessoa ES, Nepa HBS, Paz CD; Souza AVV. Estudos dos constituintes químicos e
349 atividade antibacteriana do óleo essencial de *Lippia gracilis* a *Xanthomonas campestris* pv. *viticola* "in vitro". *Summa*
350 *Phytopathologica*, Botucatu. 2014;40(3):277-280. Portuguese.
351
- 352 18. Carson CF, Hammer KA, Riley TV. *Melaleuca alternifolia* (tea tree) oil: A review of antimicrobial and other medicinal
353 properties. *Clinical Microbiology Reviews*. 2006;19(1):50-62.
354
- 355 19. Sagave L, Gressler LT, Flores FC, Silva CB, Vargas APC, Lovato M, Sangioni LA, Pötter L, Botton SA. Atividade de
356 nanoformulações de *Melaleuca alternifolia* e terpinen-4-ol em isolados de *Rhodococcus equi*. *Arquivo Brasileiro de*
357 *Medicina Veterinária e Zootecnia*. 2015;67(1):221-226. Portuguese.
358
- 359 20. Vieira TR, Barbosa LCA, Maltha CRA, Paula VF, Nascimento EA. Constituintes químicos de *Melaleuca alternifolia*
360 (Myrtaceae). *Química Nova*. 2004;27(4):536-539. Portuguese.
361
- 362 21. Souza AD, Roggerio TU, Furlan MR, Aoyama EM. Óleo de melaleuca (*Melaleuca alternifolia* Maiden & Betche, Cheel)
363 no controle de cercosporiose em beterraba. *Revista Brasileira de Plantas Mediciniais*. 2015;17(4):1078-1082. Portuguese.
364
- 365 22. Martins JAS, Sagata E, Santos VA, Juliantti FC. Avaliação do efeito do óleo de *Melaleuca alternifolia* sobre o
366 crescimento micelial *in vitro* de fungos fitopatogênicos. *Bioscience Journal*. 2010;27(1):49-51a. Portuguese.
367
- 368 23. Brum RBCS, Cardon CH, Pereira AS, Cardoso DP, Dos Santos GR. Atividade antifúngica de óleos essenciais sobre
369 fungos fitopatogênicos. *Magistra*. 2014;26(3):361-371. Portuguese.
370
- 371 24. Silva BVS, Paiva GF, Barbieri TPOM, Souza GHS, Gonçalves FJT. Efeito dos Óleos Essenciais Sobre o Crescimento
372 Micelial *in vitro* de *Colletotrichum* spp. *Cadernos de Agroecologia*. 2018;13(2):1-9. Portuguese.
373
- 374 25. Castellani A. Maintenance and cultivation of common pathogenic fungi of man in sterile distilled water. Further
375 Researches. *Journal of Tropical Medicine and Hygiene*. 1967;70:181-84.
376
- 377 26. Concha JM, Moore LS, Holloway WJ. Antifungal activity of *Melaleuca alternifolia* (tea-tree) oil against various
378 pathogenic organisms. *Journal. American Podiatric Medical Association*. 1998;88(10):489-92.
379
- 380 27. Martins ESCS, Santos MS, Barros HMM, Farias MAA. Efeito dos óleos essenciais de citronela, alecrim e erva-cidreira
381 no controle *in vitro* de *Ralstonia solanacearum* em pimentão. *Tecnologia & Ciência Agropecuária*. 2010;4(1):9-
382 13b. Portuguese.
383
- 384 28. Pereira MC, Vilela GR, Costa LMAS, Silva RF, Fernandes AF, Fonseca EWNF, Picolli RH. Inibição do
385 desenvolvimento fúngico através da utilização de óleos essenciais de condimentos. *Ciência e Agrotecnologia*, v. 30, n. 4,
386 p. 731-738, 2006. Portuguese.
387
- 388 29. Bastos CN. Efeito do óleo de *Piper aduncum* sobre *Crinipelis pernicioso* e outros fungos fitopatogênicos. *Fitopatologia*
389 *Brasileira*. 1997;22(3):441-443. Portuguese.
390
- 391 30. Oliveira JA. Efeito do tratamento fungicida em sementes no controle de tombamento de plântulas de pepino (*Cucumis*
392 *sativus* L.) e pimentão (*Capsicum annum* L.). 1991. 111 f. Dissertação (Mestrado em Fitossanidade) – Escola Superior de
393 Agricultura de Lavras, Lavras, 1991. Portuguese.
394
- 395 31. Santos PL. Manejo de *Macrophomina phaseolina* (Tassi) Goid. em sementes de feijoeiro (*Phaseolus vulgaris* L.) com
396 óleos essenciais e antagonistas. 2018. 76fls. Tese (Doutorado em Agronomia) – Universidade Estadual Paulista,
397 Botucatu-SP. Portuguese.
398
- 399 32. Silva MAD, Silva WR. Comportamento de fungos e de sementes de feijoeiro durante o teste de envelhecimento
400 artificial. *Pesquisa Agropecuária Brasileira*. 2000;35(3):599-608. Portuguese.
401
- 402 33. Limonard T. A modified blotter test for seed health. *Netherland Journal of Plant Pathology*. 1996;72(2):319-321.

Formatted: Portuguese (Brazil)

403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462

34. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria; 2018. Disponível em: <https://www.R-project.org/>.
35. Hammer KA, Carson CF, Riley TV. Effects of *Melaleuca alternifolia* (Tea Tree) essential oil and the major monoterpene component Terpinen-4-ol on the development of single and multistep antibiotic resistance and antimicrobial susceptibility. *Antimicrobial Agents and Chemotherapy*. 2012;52(2):909-915.
36. Hammer KA, Carson CF, Riley TV., Antifungal effects of *Melaleuca alternifolia* (Tea tree) oil and its components on *Candida albicans*, *Candida glabrata* and *Saccharomyces cerevisiae*. *Journal of Antimicrobial Chemotherapy*. 2004;53:1081-1085.
37. Azevedo FA, Polydoro DA, Bastianel M, Kupper KC, Stuart RM, Costa FP, Pio RM. Resposta de diferentes genótipos de tangerinas e seus híbridos à inoculação *in vitro* e *in vivo* de *Alternaria alternata*. *Revista Brasileira de Fruticultura*. 2010;32:1-8. Portuguese.
38. Riccioni L, Orzali L. Activity of tea tree (*Melaleuca alternifolia*, Cheel) and thyme (*Thymus vulgaris*, Linnaeus.) essential oils against some pathogenic seed borne fungi. *Journal of Essential Oil Research*. 2011;23(6):43-47.
39. Chutia M, Bhuyan PD, Pathak MG, Sharma TC, Boruah P. Antifungal activity and chemical composition of *Citrus reticulata* Blanco essential oil against phytopathogens from North East India. *LWT - Food Science and Technology*. 2009;42:777-780.
40. Guimarães LGL, Cardoso MG, Sousa PE, Andrade J, Vieira SS. Atividades antioxidante e fungitóxica do óleo essencial de capim-limão e do citral. *Revista Ciência Agronômica*. 2011;42(2):464-472. Portuguese.
41. Barboza HS. Efeito fungitóxico do óleo essencial de alecrim-da-chapada em *Alternaria* sp. 2015. 31fls. Trabalho de Conclusão de Curso (Graduação em Agronomia) – Universidade Federal do SemiÁrido, Mossoró, RN, Brasil. 2015. Portuguese.
42. França KRS, Lima TS, Cardoso TAL, Ugulino ALN, Rodrigues APMS, Mendonça Júnior AF. *In vitro* Effect of essential oil of peppermint (*Mentha x piperita* L.) on the mycelial growth of *Alternaria alternata*. *Journal of Experimental Agriculture International*. 2018;26(5):1-7.
43. Antunes MDC, Cavacob A. The use of essential oils for postharvest decay control. A review. *Flavour Fragrance Journal*. 2010;25:351-366.
44. Russo M, Suraci F, Postorino S, Serra D, Roccotelli A, Agosteo GE. Essential oil chemical composition and antifungal effects on *Sclerotium cepivorum* of *Thymus capitatus* wild populations from Calabria, southern Italy. *Revista Brasileira de Farmacognosia*, v. 23, n. 2, p. 239-248, 2013.
45. Feng W, Zheng X. Essential oils to control *Alternaria alternata* *in vitro* and *in vivo*. *Food Control*. 2007;18:1126-1130.
46. Khalili ST, Mohsenifar A, Beyki M, Zhavah S, Rahmani-Cherati T, Abdollahi A, Bayat M, Tabatabaei M. Encapsulation of Thyme essential oil in chitosan-benzoic acid nanogel with enhanced antimicrobial activity against *Aspergillus flavus*. *Food Science and Technology*. 2015;60(1):502-508.
47. Simões CMO, Spitzer V. Óleos voláteis. In: Gosmann G, Mello JCP, Simões CMO, Schenkel EP. (Org.). *Farmacognosia: da planta ao medicamento*. 5th ed. Porto Alegre: UFRGS/UFSC, p.467-495. 2004. Portuguese.
48. Rozwalka LC. Extratos, decotos e óleos essenciais de plantas medicinais e aromáticas na inibição de *Glomerella cingulata* e *Colletotrichum gloeosporioides* de frutos de goiaba. *Ciência Rural*. 2008;58(2):301-307. Portuguese.
49. Moraes LAS, Ramos NP, Gonçalves GG, Betiol W, Chaves FCM. Atividade antifúngica dos óleos essenciais em sementes de feijão cv. Carioquinha. *Horticultura Brasileira*. 2008;26(2):6261-6266 Portuguese.
50. Wanderley MJA, Silva TMB, Arruda JD, Rocha RB, Cruz CSA, Wanderley PA. Efeito de óleos essenciais sobre *Callosobruchus maculatus* em feijão caupi *Vigna unguiculata* L. Walp. In: 6º Congresso Brasileiro de Plantas Oleaginosas, Óleos, Gorduras e Biodiesel. 2009. Minas Gerais, Montes Claros, 2009. Portuguese.

Formatted: Polish (Poland)

463 51. Derbahlah AS, Hhamza AM, Gazzy AA. Efficacy and safety of some plant extracts as alternatives for *Sitophilus oryzae*
464 control in rice grains. Journal of Entomology. 2012;9:57-67.
465
466
467
468
469
470
471

