1 2	Obstructive Sialadenitis of Submandibular Gland Due to a Nail-Like Fish Bone Foreign Body: A Rare Case Report
3	Running title: Obstructive sialadenitis of submandibular gland
4	
5	Abstract
6	Introduction:
7 8 9	Foreign body-associated sialadenitis of submandibular gland is not often and scarce within the literatures. In this study, a report of piece of Nail-like fish bone foreign body entering the Wharton's duct causing an acute sialadenitis is presented.
10 11	Foreign bodies must be explored and all suspected areas must be examined carefully for avoiding secondary problems and surgeries in the future.
12 13	Foreign bodies in the oral and maxillofacial region are often experienced after trauma and dental treatment.
14	Case report:
15 16 17 18 19 20 21 22 23	We describe a case of obstructive sialadenitis in the submandibular gland caused by penetration of a fish bone in a 68-year-old man. Hhe had swelling and spontaneous pain in the left submandibular region. Radiographic examination didn't show foreign body in the submandibular gland. Initially we diagnosed obstructive sialadenitis in the left submandibular gland and we guess something like salivary stone may be the cause of this swelling so compressing and milking of Wharton duct was done and suddenly a tip of foreign body was appeared. The foreign body measured 1.3cm *3 mm*2mm and was a nail-like object. On pathological examination, the foreign body was found to be a fish bone (cartilage-like organic material).
24	Conclusion
25 26 27 28	This case demonstrated that precise and proper inspection and examination, milking and then paying attention to secretion of salivary gland lead to proper diagnosis and after that suitable treatment, so this could reduce costly assessment and treatment, also lessen bewilderment of the patient.
29 30	Keywords: Obstructive sialadenitis, foreign body, foreign body-associated sialadenitis, submandibular gland, nail-like fish bone
31	
32	1. Introduction
33	Obstructive sialadenitis of the submandibular gland is usually due to sialoliths, but foreign body-

Comment [f1]: Consider to revise this sentence since it is unclear.

associated sialadenitis is not often and rare in the literatures [1-14]. The mechanism of foreign-

- body entry is generally traumatic [4-6, 15], thus reports of a foreign body entering the salivary 35
- gland intraorally through Wharton's or Stensen's duct are not common and usual. [4, 5, 7, 16]. 36
- 37 Foreign body-induced sialoliths are even more rare [6, 8, 9, 17, 18]. As we know,
- 38 sSialoendoscopy, is one of minimally invasive procedure, that has recently been applied for
- 39 direct diagnosis of pathologic features in the ductal system and removal of sialoliths and foreign
- bodies in the duct of the salivary gland but it is used for distal of the salivary's duct [4-6,12,19]. 40
- Here we report a patient has cured with milking and compressing Wharton's duct and suddenly 41
- the foreign body like fish bone-induced sialoliths came out of the duct and after that prescribing 42
- antibiotic. 43

45

2. Case Report

- 46 A 68-year-old male patient was referred to Department of Oral and maxillofacial Medicine and
- surgery with a chief complaint of a swelling in left side of neck since 14 days ago and he was 47
- suffering pain since 2 weeks ago, by the way pPain was increased in intensity while swallowing 48
- and eating meals. (Figure 1A, 1B). The pPatient gave history of fever and malaise, difficulty in 49
- eating and also speaking. He expressed that the swelling was small in size and immediately 50
- increase to present size of 5-6 cm. 51
- The patient also suffered from dDiabetes, hHypertension and he had history of cCardiac 52
- aArrhythmia. He consulted with his dentist and get antibiotic (Cap Amoxicillin 500 mg) every 8 53
- hour and also panoramic view radiography was taken but the dentist couldn't find the cause 54
- (figure 2A). For a short duration partial recovery was done but after that the swelling was 55
- 56 recurred.

- Clinical examination of intraoral revealed that ovoid shape swelling in floor of the mouth and it 57
- 58 measured 5-6 cm in diameter. Extra oral findings reveal that enlargement of lateral neck
- extended from lower border of mandible (5 cm anterior to angle of mandible) to lateral upper 59
- border of thyroid cartilage. (Figure 1A) The border of enlargement was well-defined and regular 60
- border, surface was smooth and skin over the swelling was intact like adjacent tissues. It was 61
- 62 tender on palpation but temperature was not raised. Consistency of swelling was soft and rubbery
- and fluctuation was present but it was not fixed to overlying skin (Figure 1a, 1b). 63
- Intra oral examination showed swelling of Wharton's duct in left floor of the mouth (Figure 2b). 64
- 65 lit was tender and painful on palpation, and consistency of swelling was soft and fluctuant.
- It is important to formulate the differential diagnosis when swelling and mass is seen at the side 66
- of neck since this would help further evaluation of the condition and management of the patient. 67
- After considering all clinical findings following entities were considered in differential 68
- diagnosis—acute submandibular sialadenitis and benign swelling of neck. 69





Figure 1: Swelling of submandibular region that extended to lateral neck, and the patient had tenderness on palpation (a). Asymmetry and swelling of left submandibular region, frontal view of patient (b).

Compressing and Milking of Wharton duct was performed and suddenly the nail-like fish bone foreign body came out of the duct and after that the pus were pushed out and we waited for about 15 minutes to get out of whole pus then we rinsed the orifice and duct (Figure 3a,3b, Figure 4a) High dose of oral antibiotic (c; cap Amoxicillin (2 g) every 6 hours was prescribed for one day and then it was tapered until one week after culture test. We emphasized use antibiotics 2 hours before eating meals for better gGastrointestinal absorption. Also use of adequate hydration and sialagogues, pure honey as a mouth rinse for 3-4 time a day, was advised to him.

The recalled sessions for follow-up was 3 days and 1 week later. Significant improvement was achieved. Foreign body sent for histopathological examination. The report of biopsy was interpreted as a fishbone (Figure 4b, Figure 5a, 5b). Final diagnosis of obstructive submandibular sialadenitis was given. There is no residual or recurrent swelling apparent in the area of intervention after a follow-up period of 6 months.

Comment [f2]: This word seems wrong.

Comment [f3]: Consider to join some of these sentences, since there are too short and a few lack the verb.




Figure 2: (close view) Panoramic view of <u>the patient w</u>Without illustrating opacity of <u>the foreign body in the left</u> submandibular duct (a). Photography of swelling of orifice of Wharton's duct in <u>the left floor of the mouth</u> (compare with right floor of the mouth) (b).




Figure 3: photography (a) illustrating tip of the foreign body in orifice of Wharton's duct that appeared after milking of this orifice . Close view of the nNail-like foreign body (bone fish) that different derbies (???) sSurrounded it around (b)





Figure 4) Photography from the floor of the mouth illustrating pushing out of pus from orifice of Wharton's duct that appeared after removing the foreign body (a). Improvement of swelling of the floor of the mouth after removal of the foreign body and treatment with antibiotic for a period of 2 weeks (b).






Figure 5: The report of biopsy was interpreted as foreign body (fish bone) (a). Microscopic view of $\underline{\text{the}}$ foreign body (fish bone) (b)

3. Discussion

A search of Medline using the key words foreign body, fish bone and submandibular gland revealed that the first case was published ion 1990[10].

Many different kind of things as foreign bodies have been found and reported in salivary glands such as paper clips, feathers ,_toothbrush bristles, spikes of wheat, blades of grass, hairs ,_pencil lead, plastic pen tops, plant material, splinters of wood, pieces of metal, slivers of fingernail, and fish bones [4,6,7,10,11,13,16].

- 106 Two2 reasons and hypotheses considered for entering foreign bodies into the salivary gland: one
- of them is penetrating trauma [4, 6, and 20] and the other reason is retrograde migration [7, 13,
- 108 and 21]
- 109 Perhaps iIn our case because of partial edentulous and inability to complete mastication and also
- 110 lessening of salivary gland's secretion due to presumption of drugs, contingency of entering
- foreign body and retrograde migration will be increased.
- When traumatic injury is a cause and history of obstructive sialadenitis so diagnosis is relatively
- easy [6]. However, in cases without traumatic injury, it is obscure whether the cause is retrograde
- migration for these 4 reasons: (1) there is almost steady salivary flow; (2) the orifice of duct is
- mobile and can twist in all directions; (3) the diameter of the duct at the orifice is miniature; and
- 116 (4) in most submandibular glands, there is a sphincter-like system in the first 3 cm of Wharton's
- duct that prevents the retrograde migration of substances [17,21,22,23]. Findings support the
- possibility that some sialoliths might result from retrograde migration of a fish bone through the
- orifice of Wharton's duct. Fish bones are one of the most common foreign bodies found in the
- pharynx and esophagus, but they are very rarely found in the salivary gland or the duct [4-10, 13,
- 121 14, 24, and 25]. In previous studies, the incidence of a fish bone encompassed by a sialolith
- 122 ranged from 2.8% (12/423) to 4.4% (5/114) of patients with sialoliths of the submandibular
- 123 gland [6, 8, and 26].
- There are interesting statics about fish bone as a foreign body, for example, in one of review of
- 125 English-language literature, migration of a fish bone into the salivary gland was more common
- in men than in women [4-10, 13, 26]. Fish bones were more often present in the submandibular
- gland than the parotid gland [4-10,13,14,26] and were more often located in the left side of the
- submandibular gland than in the right side [4-10,13,14,26]. Stone formation induced by a fish
- bone tended to be obviously related to occupation (fisherman), dietary habit (seafood), and
- history of injury (recollection of a fish-bone injury and subsequent symptoms) [6]. Fish bone-
- induced sialoliths were previously reported to be 3–18 mm in size [5-8, 10, 26].
- Our case was matched to this literature, in our male patient, the nail-like fish bone foreign body
- was a total of 1.2 cm in length and 0.3 cm in diameter in the left submandibular duct.
- 134 In evaluating the patient with sialadenitis, these steps should be taken in the following order:
- 135 1. History, 2. Pphysical examination, 3. Cculture, 4. Laboratory investigation, 5. Radiography,
- and if indicated, 6. Ffine-needle aspiration biopsy.
- 137 There are wide range of approaches for management and treatment of sialadenitis, these include
- 138 conservative medical management to more invasive surgical intervention.
- One management scheme is as follows:
- Acute sialadenitis –
- 141 Medical management (hydration, antibiotics [oral versus parenteral], warm compresses and
- massage, sialagogues);

- 143 Surgical management (consideration of incision and drainage versus excision of the gland in
- 144 cases refractory to antibiotics, incision and drainage with abscess formation, gland excision in
- cases of recurrent acute sialadenitis)
- 146 Conservative therapies for acute management of obstructive sialadenitis, include: hydration,
- analgesia (NSAIDs), sialagogues to stimulate salivary secretion, and regular, gentle gland
- 148 massage.

- As we know the first- line therapy for stones in distal ducts of salivary glands is interventional
- sialadenoscopy, also we can use this approach for removal of foreign bodies.
- 151 By the way, if infection is present, empiric antibiotic therapy should be given after proper
- 152 cultures have been obtained.
- 153 In our case, because of the improvement of sign and symptoms, we continued the same antibiotic
- that we prescribed before for our patient .By the way, with removal of fish bone foreign body,
- most of the pus was pushed out. As we know the foreign body was the main cause.

157 **4. Conclusion**

- 158 This case demonstrated that precise and proper inspection and examination lead to proper diagnosis and
- after that suitable treatment, so this could reduce costly and expensive assessment and treatment, also
- lessen bewilderment of the patient.
- 161 Another important matter is that p, Patients with any form of sialadenitis should be educated as to the
- worthiness of hydration and excellent oral hygiene.
- At the end, mMilking and pay attention to transparency (glassiness) and canescent of secretion of salivary
- gland are helpful for achievement of proper diagnosis.

165 **References**

- 166 1. Taneja M, Taneja MK. Foreign body Wharton's duct. Indian J Otolaryngol Head Neck Surg 2011;
- 167 63:300-301.
- 168 2. Ozturk, Kayhan MD; Erdur, Omer MD; Aksoy, Ceren MD. Foreign Body of Submandibular Gland.
- Journal of Craniofacial Surgery: October 2016 Volume 27 Issue 7 p e600–e601.
- 170 3. Shameeka Thopte, Shams Ul Nisa, Abhijeet Jadhav, Rohan Chaudhari. Sialolithiasis Of
- 171 Submandibular Gland With Acute Suppurative Sialadenitis: A Case Report. World Journal Of Pharmacy
- And Pharmaceutical Sciences 5(4) · March 2016
- 173 4. Gill AS, Kieliszak C, Joshi AS. Sialendoscopy as a management tool in patients with foreign body
- impaction of the salivary gland. Am J Otolaryngol 2016; 37:369–71.
- 175 5. Yamano Y, Uzawa K, Ito H, Tanzawa H. Endoscopically assisted removal of a fish bone penetrating
- the parotid duct: an unusual case. J Oral Maxillofac Surg 2014; 72:1343–9.
- 177 6. Xie L, Zheng L, Yu C, Yang C, Chen Z, Yun B, et al. Foreign body induced sialolithiasis treated by
- sialoendoscopic intervention. J Craniofac Surg 2014; 25:1372–5.

Comment [fg4]: This part is unclear. Are medical and surgical treatment referred to acute sialadenitis? And conservative therapies? Consider to rephrase as bulleted list.

- 179 7. Su YX, Lao XM, Zheng GS, Liang LZ, Huang XH, Liao GQ. Sialoendoscopic management of
- 180 submandibular gland obstruction caused by intraglandular foreign body. Oral Surg Oral Med Oral Pathol
- 181 Oral Radiol 2012; 114:e17–21.
- 182 8. Yu C, Yang C, Zheng L. Sialendoscopic findings in patients with obstructive sialadenitis: long-term
- experience. Br J Oral Maxillofac Surg 2013; 51:337–41.
- 184 9. Sato K, Umeno H. Clinical photographs. Fish bone-induced sialolith. Otolaryngol Head Neck Surg
- 185 2009; 141:539-40.
- 186 10. Abe K, Higuchi T, Kubo S, Oka M. Submandibular sialoadenitis due to a foreign body. Br J Oral
- 187 Maxillofac Surg 1990; 28:50–2.
- 11. Ozturk K, Erdur O, Aksoy C. Foreign body of submandibular gland. J Craniofac Surg 2016; 27:e600-
- 189 601.
- 190 12. Ardekian L, Klain H, Peled M. Obstructive sialadenitis of submandib-ular gland due to foreign body
- successfully treated by sialoendoscopic intervention. J Oral Maxillofac Surg 2009; 67:1337–9.
- 192 13. Derin S, Sahan M, Kule M, Koseoglu S, Celik OI. Fish bone induced sialolith in Warthon duct. J
- 193 Craniofac Surg 2015; 26:e663–664.
- 194 14. Matsuo T. Acute suppurative parotitis caused by a fish bone: a case report. Int J Oral Maxillofac Surg
- 195 1997; 26:54.
- 196 15. Amarbir S.GillBSaChristopher R.KieliszakDObArjun S. Sialendoscopy as a management tool in
- 197 patients with foreign body impaction of the salivary gland. American Journal of Otolaryngology. Volume
- 198 37, Issue 4, July–August 2016, Pages 369-371.
- 199 16. Sivapatha Sundaram Sreetharan, Rajan Philip. Unusual Foreign Body of Parotid Gland Presenting as
- 200 Sialolithiasis: Case Report and Literature Review. Case Reports in Otolaryngology. Volume 2012 (2012),
- 201 Article ID 367349, 3 pages.
- 202 17. Marchal F, Kurt AM, Dulguerov P, Lehmann W. Retrograde theory in sialolithiasis formation. Arch
- 203 Otolaryngol Head Neck Surg 2001; 127:66–8.
- 204 18.Yasufumi KOSUGI, Toshinori Iwai, Shinsuke OHTA, Iwai TOHNAI. A case of an endoscopically
- 205 removed parotid duct sialolith. Nippon Koku Geka Gakkai zasshi· March 2017; 63(3):153-157.
- 206 19.Maria E.PapadakiDMD, MD,Joseph P.McCainDMD,KingKimDMD,Ronald L.KatzDMD§Leonard
- 207 B.KabanDMD, MD||Maria J.TroulisDMD, MSc .Interventional Sialoendoscopy: Early Clinical
- 208 Results. Journal of Oral and Maxillofacial Surgery. Volume 66, Issue 5, May 2008, Pages 954-962.
- 209 20. P Capaccio, Torretta S, Ottaviani F,et al. Modern management of obstructive salivary diseases. Acta
- 210 Otorhinolaryngol Ital. 2007 Aug; 27(4): 161–172.
- 21. Yu-xiongSuMD, DDS^aXiao-meiLaoBDS, DDS^bGuang-senZheng Li-zhongLiangDDS^bXing-
- 212 huaHuangBDS^bGui-QingLiaoMD, DDS^cSialoendoscopic management of submandibular gland
- 213 obstruction caused by intraglandular foreign body. Oral Surgery, Oral Medicine, Oral Pathology and Oral
- Radiology Volume 114, Issue 5, November 2012, Pages e17-e21.
- 215 22. F. Marchal, A.M. Kurt, P. Dulguerov, W. Lehmann. Retrograde theory in sialolithiasis formation
- 216 .Arch Otolaryngol Head Neck Surg, 127 (2001), pp. 66-68.

Formatted: Italian (Italy)

- 217 23. Mahabaleshwara C.H, Jayadeep Nidyalmale, Abhishek P.T, Ashoka G. 'FISH BONE': THE
- 218 REASON BEHIND SUBMANDIBULAR SIALADENTITS A UNIQUE CASE REPORT. International
- 219 Journal of Clinical And Diagnostic Research ISSN 2395-3403 Volume 5, Issue 3, May-June 2017. Intl. J.
- 220 Clin. Diag. Res. 2017; 5(3): I.
- 221 24. Ikenberry SO, Jue TL, Anderson MA, et al; Management of ingested foreign bodies and food
- impactions. Gastrointest Endosc. 2011 Jun73 (6):1085-91. doi: 10.1016/j.gie.2010.11.010.
- 223 25. Marc H Hohman, Wayne J Harsha, K Linnea Peterson. Migration of Ingested Foreign Bodies into the
- 224 Thyroid Gland: Literature Review and Case Report.The Annals of otology, rhinology, and laryngology
- 225 February 2010.119(2):93-8 ·

- 226 26. Yu C, Yang C, Zheng L, Wu D. Endoscopic observation and strategic management of obstructive
- submandibular sialadenitis. J Oral Max-illofac Surg 2010; 68:1770–5.