

Original Research Article

Comparative Bioremediation Potential of *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KY085976 on oil spill dispersant polluted terrestrial soil

ABSTRACT

**Aim:** To assess the bioremediation potential of *Pseudomonas aeruginosa* and *Bacillus megaterium* on oil spill dispersant polluted terrestrial soil.

**Study Design:** The study employs experimental design, statistical analysis of data and interpretation.

**Place and Duration of the Study:** Soil samples were collected from Kegbara-Dere Community, Gokana L.G.A, and were transported to the Microbiology Laboratory for analyses, while, Oil spill dispersant (OSD/LT and OSD/Seacare) were collected from Barker and Hughes Nig Ltd (formally mil park Nigeria limited), all in Rivers state, Nigeria. This investigation study lasted for 28 days and sampling was done every 7day period.

**Methodology:** Seven microcosm were done, three with oil spill dispersant LT (OSD/LT) and three with oil spill dispersant SC (OSD/SC), (*Pseudomonas aeruginosa*, *Bacillus megaterium* and mixt of them) were analysed total hydrocarbons concentration (TPH) Bacteriologically and physicochemical parameters were determined using standard method.

**Methodology:** Collected Soil samples were further polluted with oil spill dispersants and were analysed Bacteriologically and physicochemical parameters were determined using standard method.

**Results:** Nine (9) bacterial genera were isolated from oil spill polluted terrestrial soil including; *Pseudomonas*, *Bacillus*, *Serratia*, *Micrococcus*, *Klebsiella*, *Corynebacterium*, *Listeria*, *Streptococcus*, and *Proteus*. *Pseudomonas*, *Bacillus*, *Serratia*, *Proteus*, and *Micrococcus* species were identified as dispersant utilizers. Addition of (*Pseudomonas aeruginosa* or *Bacillus megaterium*) to oil spill dispersant polluted soils, enhanced decrease in THC (total hydrocarbon of concentration) of the soil. THC for control reduced from 18348.68 to 9111.84 (mg/kg);

TS+OSD/LT+Bm<sub>2</sub>, 18348.68 to 7092.11 (mg/kg); TS+OSD/LT+Pa, 18348.68 to 6263.16 (mg/kg);

TS+OS/LT+Bm+Pa<sub>2</sub>, 18348.68 to 2473.68 (mg/kg); TS+OSD/SC+Bm, 18348.68 to 6421.05 (mg/kg);

Formatted: Font: Italic

Formatted: Font: Italic

**Comment [GP1]:** Results: Nine (9) bacterial genera were isolated from oil spill polluted terrestrial soil including; *Pseudomonas*, *Bacillus*, *Serratia*, *Micrococcus*, *Klebsiella*, *Corynebacterium*, *Listeria*, *Streptococcus*, and *Proteus*. *Pseudomonas*, *Bacillus*, *Serratia*, *Proteus*, and *Micrococcus* species were identified as dispersant utilizers. Addition of (*Pseudomonas aeruginosa* or *Bacillus megaterium*) to oil spill dispersant polluted soils, enhanced decrease in THC of the soil. THC for control reduced from % ;microcosmco with LT an *B. megaterium* .... %; LT *Ps aeruginosa* ...%; .....  
The pH of soil samples ranged from 5.66 to 7.37 across the various set up. Moisture content ranged from 0.03 to 0.4 across the various set up. The bioremediation rates were as follows in all microcosm around 9.27 – 9.46 Log10Cfu/g. .... PUB ranged from 7.02±0.46 Log10Cfu/g to 7.30±0.45Log10Cfu/g; DUB, ranged between 6.95±0.51Log10Cfu/g and 7.08±0.52Log10Cfu/g; TBC ranged between 8.71±0.34 and 9.06±0.41Log10Cfu/g; TPC ranged between 6.85±0.76 Log10Cfu/g and 7.11±0.47Log10Cfu/g across the various set up.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

**Comment [GP2]:** What it is ?  
The reader must understand what their letters mean

**Comment [GP3]:**

TS+OSD/SC+Pa, 18348.68 to 5618.42 (mg/kg); TS+OSD/SC+Bm+Pa, 18348.68 to 5835.53 (mg/kg), from day 1 to day 28. The pH of soil samples ranged from 5.66 to 7.37 across the various set up. Moisture content ranged from 0.03 to 0.4 across the various set up. The bioremediation rates were as follows: THB, (TS(CTRL) 9.27±0.43Log10Cfu/g; TS+OSD/LT+Pa, 9.31±0.46Log10Cfu/g; TS+OSD/LT+Bm, 9.40±0.46Log10Cfu/g; TS+OSD/LT+Bm+Pa, 9.31±0.41Log10Cfu/g; TS+OSD/SC+Pa, 9.36±0.46Log10Cfu/g; TS+OSD/SC+Bm, 9.43±0.46Log10Cfu/g); and TS+OSD/SC+Bm+Pa, 9.36±0.42Log10Cfu/g. PUB ranged from 7.02±0.46 Log10Cfu/g to 7.30±0.45Log10Cfu/g; DUB, ranged between 6.95±0.51Log10Cfu/g and 7.08±0.52Log10Cfu/g; TBC ranged between 8.71±0.34 and 9.06±0.41Log10Cfu/g; TPC ranged between 6.85±0.76 Log10Cfu/g and 7.11±0.47Log10Cfu/g across the various set up.

**Conclusion:** From this study, bioremediation of *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KY085976 should be regarded as a key component in the clean-up strategy for oil spill dispersants pollution

Formatted: Font: Italic

Formatted: Font: Italic

**Keywords:** Bioremediation, Oil spill, Dispersants, OSD/LT, OSD/Seacare, Total hydrocarbon content, Terrestrial soil.

## INTRODUCTION

Bioremediation is the use of microorganisms and their products to remove contaminants from the environment (1). It is also a process, which depends on biological mechanisms to mineralized or transform concentration of pollutants to an innocuous state (2). Bioremediation is among new technologies or approaches that derive its scientific justification from the emerging concept of environmentally friendly chemistry and engineering. This technology is a fast growing and promising remediation option that is increasingly studied and applied for pollutant clean up (3). It is relatively cost-effective and non-invasive and will lead to complete mineralization. As such, the isolation of potentially applicable microorganisms to bioremediation of a variety of oily contaminations has received a lot of ink in the related literature (4,5). However, the main drawback of bioremediation processes in a majority of cases is the slow biodegradation rate of hydrocarbons (6). So many researchers have studied bioremediation, and their studies focused on hydrocarbons on account of frequent pollution of soil and ground water with this particular type of pollutant (7,8,9,10). The technology commonly used for the soil remediation includes mechanical, burying, evaporation, dispersion and washing. Currently, accepted methods for removal of pollutants are incineration, land filling and safe disposal. Though, these technologies are expensive and can leads to incomplete decomposition of contaminants (11,12). The conventional methods to clean-up oil spill from terrestrial and aquatic ecosystems are; mechanical method, chemical method and microbial degradation. The benefits in the use of biological agents is higher than those derived from the use of physical and chemical methods in the restoration of a contaminated environment (13,14,15,16,17). Mechanical and chemical methods generally used to remove hydrocarbon from contaminated sites have limited effectiveness and can be expensive (18,19). Physical and chemical methods to reduce hydrocarbon pollution are expensive (20). In recent years microbial degradation of pollutants is a sustainable way to clean up the contaminated environment (21). The success of oil spill bioremediation depends on one's ability to establish and maintain conditions that favour enhanced oil biodegradation rates in the contaminated environment, such as presence of microorganisms with appropriate metabolic abilities. Where there are frequent oil

spills and difficult terrain for clean up exercise by mechanical means, bioremediation of oil spills become very important in that region (22). Chemicals are used to change the characteristics feature of the oil (23). Dispersants are chemical agents that reduce tension between oil and water interface thereby enhancing the natural process of dispersion by producing large amounts of small droplets of oil that are drawn along into water column. Oil spill dispersant may present a short term effects on marshes or may present a long term effect if not properly handled. The establishment of oil spill dispersant preparedness practices in marshland communities will be very crucial to reduce the impact from oil (24).

Dispersants are most effective when applied immediately after a spill, before the lightest components in the oil evaporates. The use of dispersants in nearshore areas is expected to increase the exposure of aquatic organisms to petroleum (25). Oil pollution, whether acute or chronic, has detrimental effects on agricultural lands and hence significant effects on plant growth (26). Oil spill pollution has become a universal problem in industrialized and developing countries. It has cause a threat to our environment today by imposing a serious health hazard to human health, causes decrease in Agricultural productivity on terrestrial soil and economic loss (27,28,29). On the terrestrial environment, oil spill cause extensive damages ranging from the destruction of terrestrial flora and fauna to biomagnification of the toxic components of the petroleum conversion of arable land to barren soils and the destruction of the aesthetic quality of the environment (30) The oil reduces the soil's fertility such that most of the essential nutrients are no longer available for plants and crop utilization (31). The crude oil spillage in which hydrocarbons are found are large and complex molecules, and persistent in nature and may require a strong reagent to hinder their effect on terrestrial soil. Oil spill pollution with dispersants can be degraded effectively by using microorganism activity. This is possible because microorganisms have enzyme systems to degrade and utilize different hydrocarbon as a source of carbon and energy (32,33,34). Impact of the pollution is dependent on the volume of the hydrocarbons and the impact it has on the environment (35,36). Temperature, the availability of oxygen, pH, moisture content, the kind of microbe, the available nutrients and the type of soil are the conditions which control the breakdown of crude oil pollutants (37,38,39,40,). Hence, the essence for this study is to evaluate and compare the bioremediation potential of *Pseudomonas aeruginosa* and *Bacillus megaterium* on oil spill dispersant polluted terrestrial soil

## MATERIALS AND METHODS

### Study site and sample collection

The soil sampling was carried out at Kegbara-Dere community in Gokana Local Government Area of Rivers state, Nigeria. Kegbara-Dere is situated in the Niger Delta Area of Nigeria, between longitudes 7.01° and 7.07° E; and latitudes 4.08 and 4.2°N. Soil auger was used to collect soil sample from the polluted site and were put in sterile black polyethylene bag and labelled with masking tape, and then immediately taken to the microbiology laboratory, Rivers State University, for microbiological and some physicochemical analyses.

**Collection of Test Chemicals (Oil dispersants):** The oil spill dispersants (OSD) used in the study work and their sources were; OSD/ LT and Seacare, all from Barker and Hughes Nig Ltd (formally mil park Nigeria limited) Port Harcourt.

## Microbiological Analysis

### Serial dilution of Soil samples

The dilution method adopted was the Ten-fold technique in which 1g of soil samples was added into a prepared 9 ml of sterile diluent was pipetted into test tubes containing 9 ml diluent. This was done consecutively until an appropriate dilution was reached (41).

### Source of Microorganisms (*Pseudomonas aeruginosa* and *Bacillus megaterium*)

The method described by [Maryam et al.](#) (22) was adopted. Pure cultures of these organisms were obtained from inoculation and incubation of soil samples using nutrient Agar. Pure cultures were obtained by continuous subculturing (42). Isolates was inoculated into broth culture (43).

### Isolation of Two Test Organisms

The test organisms (*Bacillus megaterium* and *Pseudomonas aeruginosa*) were selected because of their importance as the active hydrocarbon degraders in crude oil polluted environment. They were isolated from the oil-polluted soil samples using the spread plate method (microbiological method). Soil suspensions were prepared by adopting Ten-fold serial dilution. 1g of the soil sample was measured into a test tube and 9ml of sterile distilled water was mixed with the sample. The suspension was properly shaken for 30 seconds to homogenize the solution and this served as the stock solution. Ten-fold serial dilution of all the homogenized mixture was carried out using prepared normal saline as diluents. Seven test tubes containing 9 ml of normal saline was used for the serial dilution. Aliquots of 0.1 ml from  $10^{-6}$  and  $10^{-7}$  dilutions were introduced into duplicated sterile petri dishes using sterile pipette and separately spread plated with flame sterilized bent glass spreader on well-dried Cetrimide agar plate (*Pseudomonas aeruginosa*) and nutrient agar plates. The plates were incubated at 37 °C for 24 to 48 hours. After which bacterial colonies that form during incubation period were picked with sterile inoculating loop and were streaked on freshly prepared well-dried nutrient agar plates. The plates were incubated at 37 °C for 24 hr. Discrete colonies on the plates were aseptically transferred into agar slants and bijou bottles containing 10% (v/v) glycerol, properly labelled and stored as stock cultures for preservation and identification (42, 41)).

### Confirmation of Test Organisms

The confirmation of the isolates was done according to the standard techniques in District laboratory practice in tropical countries (44) and was identified base on the Bergey's manual of Determinative Bacteriology after carrying out the morphological and various biochemical tests.

**Table 3.1 Experimental Design (Bioremediation set-up) for Terrestrial Soil sample.**

SET UP 1	1500g of soil+30ml of Distilled
SET UP 2	1500g of soil+ 20ml of OSD/LT+ 30 ml of Distilled H <sub>2</sub> O+ 50ml of <i>Bacillus megaterium</i>

Comment [GP4]: Missing quote in the text

Formatted Table

SET UP 3	1500g of soil_+20ml of OSD/LT_+30_ml of Distilled H <sub>2</sub> O_+50ml of <i>Pseudomonas aeruginosa</i>
SET UP 4	1500g of soil_+20ml of OSD/LT_+30_ml of Distilled H <sub>2</sub> O_+25_ml of <i>Bacillus megaterium</i> _+25ml <i>Pseudomonas aeruginosa</i>
SET UP 5	1500g of soil_+20ml of OSD/SC_+30ml of Distilled H <sub>2</sub> O_+50ml of <i>Bacillus megaterium</i>
SET UP 6	1500g of soil_+20ml of OSD/SC_+30ml of Distilled H <sub>2</sub> O_+50ml of <i>Pseudomonas aeruginosa</i>
SET UP 7	1500g of soil_+20ml of OSD/SC_+30ml of Distilled H <sub>2</sub> O_+25ml of <i>Bacillus megaterium</i> _+25ml <u>microcosms</u> <i>Pseudomonas aeruginosa</i>

### Soil Preparation and Application of Organisms

The experiment for a proper monitoring was set up for each soil sample and Oil spill dispersants (OSD) in the laboratory. 1500\_g of the sample collected from Kegbara-Dere Gokana L.G.A, was weighed into Seven plastic rubbers. After that, there were controls which were without organisms while others were augmented with organisms (*Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KY085976). 20ml of each of the Oil spill dispersant - OSD/LT and OSD/Seacare liquid detergent was dispensed into each of the rubber containing the soil so as to pollute it and it was then mixed properly using a sterile spatula so as to enable the dispersants mix properly with the soil. 30\_ml of distilled water were also poured into each set sample and properly stirred with a spatula for the organisms to thrive successful and have more oxygen (Table 3.1).

Bioaugmentation was the type of bioremediation carried out in which samples were augmented by adding 50\_ml of broth culture organism (*Pseudomonas aeruginosa* KX828570) to the first set up sample, 50ml broth culture organism (*Bacillus megaterium* KY085976) was added to the second set up sample and then 25ml of each broth culture organism was added to the third set up sample containing 1500g of pollute sample respectively, and they were kept at ambient temperature (28±2<sup>0</sup>C) for 28 days to determine the level of utilization by the organism (*Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KY085976). This method is referred to as ex situ bioremediation, whereby the polluted soil requires excavation and treatment can be carried out in the laboratory. This method of bioremediation can also be carried out on field or polluted sites.

## Media Used and Its Preparation

Nutrient Agar: It encourages the proliferation of organisms without segregating. 28 grams was dissolved in 1000ml of distilled water according to manufacturer's instruction. Oil Agar: The medium was prepared in the laboratory having the following composition of  $K_2HPO_4$  (0.5g),  $MgSO_4$  (0.03g),  $NaCl_2$  (0.3g),  $MnSO_4 \cdot H_2O$  (0.2g),  $FeSO_4 \cdot H_2O$  (0.02g),  $NaNO_3$  (0.03g),  $ZnCl_2$  (0.3g) and Agar agar (15\_g) into 1 litre of distilled water (43). Cetrimide Agar Medium: The preparation of this medium is by dissolving 45.3 gm in 1000ml distilled water, **autoclaved at 15psi (121<sup>0</sup>c) for 15minutes**. Cool to 45-50<sup>0</sup>C, prior to dispense. It is mainly for isolation of *Pseudomonas aeruginosa*. Nutrient Broth: This is a liquid medium used for the general cultivation of wide variety of microorganisms. The broth was prepared by dissolving 13g in 1000ml of distilled water.

## Enumeration and Isolation of bacterial population

**The spread plate technique was used to inoculate the soil samples as described by (45). The suspension was properly shaken for thirty seconds to homogenize the solution and this served as the stock solution. Ten-fold serial dilution of all the homogenized mixture was carried out using sterile distilled water as diluents. The plates were incubated at 37 °C for 24 hours. Counting of colonies followed so as to estimate the microbial load.**

**Values were expressed as colony forming unit per g (Cfu/g). The colonial morphology such as shape, edge, colour, elevation, surface, opacity and their consistency were carried out. while biochemical assay was based on Gram staining reaction, Motility, Catalase, Oxidase, Coagulase, Indole, Methyl red, Citrate, Sugar fermentation tests (43). And sub culturing of bacterial isolates was done to obtain pure culture. Bacterial colonies were picked with sterile inoculating loop and were streaked on freshly prepared well-dried NA plates (24).**

## Enumeration and Isolation of Petroleum Utilizing Bacteria

Petroleum utilizing bacteria hydrocarbons degrading bacteria (PUB) were enumerated as adopted from (46, 47,17) using the vapour phase method, mineral salts medium with crude oil which served as the sole source of carbon. Isolated colonies were further purified by sub-culturing and identified using biochemical tests and microscopy (48). It was done using Oil Agar (Mineral salt agar). Aliquots of 0.1ml from dilutions of  $10^{-4}$  and  $10^{-5}$  were also plated in duplicates on Mineral Salt Agar. Fungosol was added to the Mineral Salt Agar to suppress fungal growth. Spread plate method was used. A filter paper (Whatman No 1) saturated with bonny light crude oil was aseptically placed onto the covers of the Petri dishes and inverted. The culture plates were incubated for 5 to 7 days at 37 °C.

Plates yielding colonies were **afterwards enumerated,enumerated** counted and were later sub-cultured into another plate to obtain pure cultures to be used for biochemical tests. The colonies counted were expressed as the colony forming unit (CFU) per gram of the soil, after applying the appropriate correction factor. The cultural, morphological and biochemical characteristics of the discrete bacterial isolates were compared with the recommendation in Bergey's manual of determinative bacteriology.

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Values were expressed as colony forming unit per g (Cfu/g). The colonial morphology such as shape, edge, colour, elevation, surface, opacity and their consistency were carried out, while biochemical assay was based on Gram staining reaction, motility, catalase, oxidase, coagulase, indole, methyl red, citrate, sugar fermentation tests (43). And sub culturing of bacterial isolates was done to obtain pure culture. Bacterial colonies were picked with sterile inoculating loop and were streaked on freshly prepared well-dried NA plates (24).

### **Enumeration and Isolation of Oil spill dispersant (OSD) utilizing bacteria**

Enumeration of Oil spill dispersant (OSD) utilizing bacteria was done by inoculating 0.1ml aliquot of the dilutions unto mineral salt agar plates containing the OSD (49) Colonies were counted after 48 to 72 hours incubation at ambient temperature. The bacterial colonies on the plates after incubation were counted and sub-cultured onto fresh mineral salt agar plate to obtain pure cultures to be used for biochemical tests.

### **Stock Solution or Culture**

Ten percent glycerol solution was prepared, dispensed in McCartney bottles and autoclaved at 121°C for 15minutes, it was allowed to cool, then the pure cultures were inoculated into each McCartney bottle, until the clear colourless solution turns turbid and were stored in the refrigerator. This served as storage medium for pure cultures for subsequent characterization (42).

### **Identification of test Bacterial Isolates**

Identification of the bacterial isolates was done based on the method of (44).

The cultural, morphological and biochemical characteristics of the discrete bacterial isolates were compared with the recommendation in Bergey's manual of determinative bacteriology (48). The morphological and biochemical test include; gram staining, motility, catalase, coagulase, oxidase, citrate utilization, sugar fermentations, indole production and methyl red tests (24). Morphological identification observed was **colourcolor**, shape, elevation, opacity, margin, size and texture. Microscopy was done under light-microscope to check for the bacterial Grams' reaction, shapes and arrangements. The biochemical test carried out includes motility, catalase, oxidase, citrate utilization, coagulase, indole production and fermentation of the following sugars: glucose, lactose, mannitol, sucrose, and fructose.

### **Physicochemical Analysis**

Moisture Content analysis: It was determined using moisture analyser. 10.g of polluted soil from each of the set up was weighed and was put inside of washed glass petri dish and was placed inside a hot air oven for drying. After which the soils were immediately transferred into the desiccators for cooling. After cooling the soils were then reweighed and the grams gotten was subtracted from the initial 10g of the soil to get the moisture contents (29). The soil pH was obtained with a pH meter (Hannah 8314) stabilized for 15 minutes and calibrated between pH 4 and 7 with standard buffer solution, according to (50). This was determined by weighing 10g of soil sample into the beaker and addition of 10ml of distilled water. This was allowed to stand for 30 minutes and was stir occasionally

with a glass rod. After which Insert the pH meter (previously calibrated) into the partly settled suspension and take the pH reading (51, 52). Soil Temperature: The temperature of the soil was measured ex situ with a mercury thermometer. This was done by inserting the thermometer in each of the set-up rubber. Constant temperature was recorded by allowing the thermometer to remain in the soil samples (53). Total Hydrocarbon Content Analysis (THC): This was done using spectrophotometer. The procedure was undertaken 5 times (interval of 5 days) to form five replicates. During the setup process for spectrophotometric analysis, 10g of soil sample were weighed from each of the setup rubbers containing 1500g of soil sample into sterile beaker and 20ml of xylene was added and shaken properly to extract the oil from the soil and this was allowed to digest for 30 minutes and the extracted oil were sieved with whatman No 1 filter paper into test tube that was transferred into colorimeter curvette and placed in a chamber known as infrared spectrophotometer analyser. The Total Hydrocarbon Content (THC) value was determined by comparism to a calibration curve constructed from dilution of a stock solution of a 1:1 bonny light crude and oil spill dispersant. The Ultraviolet light spectrophotometric measurement was at 420\_nm and Total Hydrocarbon Content (THC) Oil Spill Dispersant (OSD) was at 560\_nm (54, 55)

#### Statistical analysis of data

Analysis of variance (ANOVA) and Duncan test was adopted using the IBM SPSS 20 software to analyze the significant differences. Data obtained were subjected to statistical analysis to determine the significant difference among the data obtained using one-way analysis of variance (ANOVA). A value of  $p > 0.05$  was considered significant while  $p < 0.05$  was considered not significant

### RESULT AND DISCUSSION

*Bacillus megaterium* and *Pseudomonas aeruginosa* helped in remediating the polluted soils caused by oil spill dispersant by reducing pollutant in the soil. Total Hydrocarbon Content (THC – mg/kg) carried on soil samples (oil spill dispersant polluted terrestrial) augmented with bacterial specie (*Bacillus megaterium* and *Pseudomonas aeruginosa*) for 28 days was shown in Fig 1. The THC of the study was determined by decrease in number. The result showed that the total hydrocarbon content decreased with an increase in time, from the day 1 to day 28 of the study, the control values for terrestrial soil ranged from 18348.68mg/kg to 9111.84 mg/kg. While the polluted soil samples of different treatments augmented with *Bacillus megaterium* and *Pseudomonas aeruginosa* were totally different from controls. The range of the soil samples with different treatments from day 1 to day 28 are as follows: TS+OSD/LT+Bm, ranges from 18348.68mg/kg to 7092.11mg/kg, TS+OSD/LT+Pa ranged from 18348.68 to 6263.16mg/kg, TS+OSD/LT+Bm+Pa, ranged from 18348.68mg/kg to 2473.68mg/kg. While TS+OSD/Seacare+Bm, ranged from 18348.68mg/kg to 6421.05 mg/kg, TS+OSD/Seacare+Pa ranged from 18348.68 to 5618.42 mg/kg, TS+OSD/Seacare+Bm+Pa, ranged from 18348.68 mg/kg to 5835.53 mg/kg respectively. At the first day (Day 1) THC was relatively constant, this is because the proper augmentation had not taken place. This indicates that the effect of time on petroleum hydrocarbon bioremediation rate was significant. Moreover, from the results gotten, it was observed that oil spill dispersant (OSD/Seacare) with *Pseudomonas aeruginosa* was more effective than with *Bacillus megaterium*, likewise, oil spill dispersant (OSD/LT) with *Pseudomonas aeruginosa*. But the OSD/LT with mixed consortium showed the highest remediation rate more than OSD/Seacare with mixed consortium. This observation is in line with the findings of (56), who reported that the use of microbial consortium mineralized pollutants more efficiently than pure isolates. Also, (38) states that combination of bacteria is best in the degradation process as to the use of a bacterium. Their percentage (%) bioremediation rate of polluted soils was shown in Fig 3 and there were as follows: controls (TS(CRTL) 50.3%, TS+OSD/LT+Bm 61.3%, OSD/LT+Pa 65.9%, and OSD/LT+Bm+Pa 86.5% Whereas, TS+OSD/Seacare+Bm had 65.0%, OSD/Seacare+Pa 69.4%, OSD/Seacare+Bm+Pa 68.2% respectively. Collectively, the highest percentages of THC in this study were from soil samples treated with oil spill dispersant and organisms, while the least was observed in treatments without oil spill dispersant and organism. This suggests that microorganisms are more abundant in oil spill dispersant polluted soils than unpolluted soils. In addition, the result obtained from mg/kg Dispersant OSD control (TS(CRTL), ranged from 1776.32 to 651.32mg/kg respectively. Whereas, the oil spill

dispersant terrestrial soil augmented with *Bacillus megaterium* and *Pseudomonas aeruginosa* showed higher bioremediation rate OSD/LT+Bm+Pa, (171.05 mg/kg), followed by OSD/Seacare+Pa 513.16mg/kg, OSD/Seacare with mixed consortium (539.47 mg/kg), OSD/LT with *Pseudomonas aeruginosa* 598.65mg/kg, OSD/Seacaer with *Bacillus megaterium* (638.16mg/kg) and OSD/LT with *Bacillus megaterium* (671.05 mg/kg). The result gotten from Dispersant OSD-mg/kg still revealed that OSD/LT with mixed consortium enhanced remediation faster than with pure isolates. These results indicate that the added mixed bacterial culture has enhanced the rate and total extent of crude oil, and oil spill dispersants biodegradation in the soil. Other researchers (57, 58) have reported enhanced crude oil biodegradation in soil caused by inoculation of microbial slurry. Moreover, in nature biodegradation of crude oil typically involves a succession of species of microorganisms within the present consortia. Consequently, study of bioremediation using bacterial consortia is encouraged because such mixed cultures display metabolic ability and superiority to pure cultures (59, 60).




Fig 1: Total Hydrocarbon Content (THC-mg/kg) during bioremediation of Oil Spill Dispersant (OSD/LT and OSD/SC) polluted Terrestrial Soil using bio-augmenting organisms *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KX085976




Fig 2: THC (Dispersant OSD - mg/kg) during bioremediation of Oil Spill Dispersant (OSD/LT and OSD/SC) polluted Terrestrial Soil using bio-augmenting organisms *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KX085976




Fig 3: % Bioremediation of Total Hydrocarbon Content in different remediation Treatment during bioremediation of Oil Spill Dispersant (OSD/LT and OSD/SC) polluted Terrestrial Soil using bio-augmenting organisms *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KX085976




Fig. 4: Soil pH of Different treatments during bioremediation of Oil Spill Dispersant (OSD/LT and OSD/SC) polluted Terrestrial Soil using bio-augmenting organisms *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KX085976




Fig. 5: Soil moisture of different treatments

**Table 2: Physicochemical Parameters on Terrestrial Soil**

Terrestrial Physicochemistry	Temp	pH	Moisture	THC
TS(CRTL)	29.8±0.45 <sup>a</sup>	5.996±0.12 <sup>a</sup>	.034±0.01 <sup>a</sup>	13730.26±0.37 <sup>a</sup>
TS+OSD/LT+Bac	31.8±1.09 <sup>b</sup>	6.012±0.25 <sup>a</sup>	.260±0.09 <sup>b</sup>	11868.516±423 <sup>a</sup>
TS+OSD/LT+Pse	31.2±0.84 <sup>ab</sup>	6.004±0.23 <sup>a</sup>	.240±0.05 <sup>b</sup>	11017.260±454 <sup>a</sup>
TS+OSD/LT+Bac+Pse	31.8±1.09 <sup>b</sup>	6.052±0.25 <sup>a</sup>	.360±0.05 <sup>b</sup>	9932.820±572 <sup>a</sup>
TS+OSD/SC+Bac	30.8±1.30 <sup>ab</sup>	6.030±0.22 <sup>a</sup>	.240±0.05 <sup>b</sup>	9780.260±492 <sup>a</sup>
TS+OSD/SC+Pse	30.8±1.30 <sup>ab</sup>	6.008±0.20 <sup>a</sup>	.240±0.05 <sup>b</sup>	10069.820±493 <sup>a</sup>
TS+OSD/SC+Bac+Pse	31.6±1.95 <sup>b</sup>	6.024±0.23 <sup>a</sup>	.380±0.04 <sup>c</sup>	9489.480±505 <sup>a</sup>
Total	31.11±1.28	5.909±0.37	.215±0.12	11302.521±444


Mean with the same alphabet across Row shows no significant different ( $p > 0.05$ )

Comment [GP5]: The data is in tables or figures, never in both.


### Physicochemical Analysis

It is a known fact that oil spill on polluted soil affects the physical, chemical and biological characteristics of the soil (61). The pH of soil samples ranged from 5.66 to 6.37 across the various set up. The highest soil pH (6.37) was recorded in the treated Terrestrial soil (TS+OSD/LT+Pa) while the lowest soil pH (5.84 to 6.16) was recorded in the Terrestrial soil control (TS(CRTL)). There was no significant difference between the soil sample in soil pH. The soil pH of the soil sample (polluted terrestrial) sites were within the same range, and they were tending from slightly acidic towards neutrality. This result concord with the observation of (62, 63), who indicated that a pH between 5 and 7.8 is favourable for the biodegradation activity of bacteria in the soil. (64) reported similar results on pH of crude oil polluted soils of Niger Delta. Plant grows in soil of pH between the ranges of 3 to 9. The non-significance difference between the soil pH in the experimental set up showed that the bioremediation of the polluted soil did not have any significant effect on soil pH (50). The reduction in pH to slight acidic range in oil polluted soil inoculated with OSD could be attributed to acidic metabolites resulting from oil biodegradation. However, the pH range observed in the present study of terrestrial soil still fall within the pH range suitable for microbial growth indicating that these isolates exhibited optimal growth at pH range of 5.84 to 6.37. (65) reported that the growth of most microorganisms is usually greatest within a pH range of 6 to 8. Soil moisture content ranged from 0.03 to 0.4 across the soil samples. The highest soil moisture (0.4) was recorded in the terrestrial soil (TS+OSD/LT+Bm+Pa and TS+OSD/SC+Bm+Pa) while the lowest soil moisture (0.03) was recorded in the Terrestrial soil (TS(CRTL)). The moisture content result of soil samples in Fig. 6, shows the differences in the moisture content of the different experimental set-up, indicating the mixed consortium with different test chemical; OSD Polluted soil + *Bacillus megaterium*+ *Pseudomonas aeruginosa* (OSD+Bm+Pa)(0.4g/10g and 0.4/10g) having the highest moisture content, followed by single organism application; Polluted soil + *Pseudomonas aeruginosa* (OSD+Pa)(0.2g/10 g and 0.2g/10 g) is equal to Polluted soil + *Bacillus megaterium* (OSD+Bm)(0.2 g/10 g and 0.2 g/10 g), while Control (soil sample without added organism CTRL)(0.03 g/10 g ) has the

lowest. (17, 29) report similar observation on the effect of moisture content on bioremediation potential of bio-stimulating and bio-augmenting agents. Alternatively, this study revealed the effects of different types of augmenting organisms, dispersants and crude oil on the moisture content of the affected soil. The high moisture content observed in the mixed consortium (Polluted soil + *Bacillus megaterium* + *Pseudomonas aeruginosa*) (OSD +Bm+Pa) could be due to its intrinsic moisture retention ability of the augmenting organisms while the control devoid of added organisms has least moisture content. These attributes (high moisture content) enhances the growth of microorganisms up to day 28 which was evident in their higher percentage bioremediation. The temperature reading for the soil sample in different set up ranged from 29°C to 30°C for control, 30°C to 33°C for oil spill dispersant (OSD/LT), while that OSD/SC range from 29°C to 34°C respectively. The highest temperature in day 28 was observed in (TS+OSD/SC+Bm+Pa), followed by (TS+OSD/LT+Bm+Pa). The temperature values obtained for the different oil spill dispersant polluted soil during the investigation study fall within the mesophilic range. This indicates that the temperature of the different oil spill dispersant polluted soils supported mesophilic bacteria throughout the investigation period. Similar result was reported by [Obir and Nwaubeta](#), (53).




Formatted: Font: Times New Roman, 12 pt, Font color: Black




Formatted: Font: Times New Roman

Figure 5a. Total Heterotrophic Bacterial count (THB) during bioremediation of Oil Spill Dispersant (OSD/LT and OSD/SC) polluted Terrestrial Soil using bio-augmenting organisms *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KX085976




Comment [GP6]: Change this graphic to bars

Figure 6. Dispersant utilizing bacterial count (DUB (Log<sub>10</sub>Cfu/g) during bioremediation of Oil Spill Dispersant (OSD/LT and OSD/SC) polluted Terrestrial Soil using bio-augmenting organisms *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KX085976




Comment [GP7]: Change this graphic to bars

Figure 7. Total Pseudomonas count (TPC) during bioremediation of Oil Spill Dispersant (OSD/LT and OSD/SC) polluted Terrestrial Soil using bio-augmenting organisms *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KX085976



Comment [GP8]: Change this graphic to bars

Figure 8. Petroleum utilizing bacterial count (PUB) during bioremediation of Oil Spill Dispersant (OSD/LT and OSD/SC) polluted Terrestrial Soil using bio-augmenting organisms *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KX085976



Comment [GP9]: Change this graphic to bars

Figure 9. Total Bacillus count (TBC) during bioremediation of Oil Spill Dispersant (OSD/LT and OSD/SC) polluted Terrestrial Soil using bio-augmenting organisms *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KX085976

Table 3: Microbial count during bioremediation process

Terrestrial Microbiology	THB	PUB	DUB	TPC	TBC
TS(CRTL)	9.27±0.43 <sup>a</sup>	7.02±0.46 <sup>a</sup>	6.95±0.51 <sup>a</sup>	6.85±0.76 <sup>a</sup>	8.71±0.34 <sup>a</sup>
TS+OSD/LT+Bm	9.40±0.46 <sup>a</sup>	7.30±0.46 <sup>a</sup>	7.08±0.52 <sup>a</sup>	6.65±0.38 <sup>a</sup>	9.06±0.41 <sup>a</sup>
TS+OSD/LT+Pa	9.31±0.46 <sup>a</sup>	7.28±0.45 <sup>a</sup>	7.04±0.53 <sup>a</sup>	7.09±0.45 <sup>a</sup>	8.63±0.29 <sup>a</sup>
TS+OSD/LT+Bm+Pa	9.31±0.41 <sup>a</sup>	7.29±0.44 <sup>a</sup>	7.05±0.52 <sup>a</sup>	7.11±0.47 <sup>a</sup>	8.83±0.39 <sup>a</sup>
TS+OSD/SC+Bm	9.43±0.46 <sup>a</sup>	7.29±0.43 <sup>a</sup>	7.00±0.51 <sup>a</sup>	6.67±0.44 <sup>a</sup>	9.04±0.39 <sup>a</sup>
TS+OSD/SC+Pa	9.36±0.46 <sup>a</sup>	7.28±0.45 <sup>a</sup>	6.99±0.54 <sup>a</sup>	6.94±0.42 <sup>a</sup>	8.79±0.77 <sup>a</sup>
TS+OSD/SC+Bm+Pa	9.36±0.42 <sup>a</sup>	7.30±0.45 <sup>a</sup>	7.02±0.53 <sup>a</sup>	6.85±0.46 <sup>a</sup>	8.80±0.39 <sup>a</sup>

Mean with the same alphabet across Row shows no significant different ( $p > 0.05$ )

KEYS: OSD/LT= Oil spill dispersant /LT, OSD/SC= Oil spill dispersant /Seacare, TS=Terrestrial soil, Bm= Bacillus megaterium. Pa= Pseudomonas aeruginosa, THC= Total hydrocarbon content, THB= Total heterotrophic bacteria, PUB= Petroleum utilizing bacterial, DUB= Dispersant utilizing bacterial, TBC= Total Bacillus count, TPC= Total Pseudomonas count.

### Bacteriological Analysis

It was observed that there was an increase in all the count from the soil polluted with oil spill dispersants when compared with control. Total heterotrophic bacterial population ranged from 8.498  $\log_{10}$  cfu/g to 9.528  $\log_{10}$  cfu/g. The highest count 9.528  $\log_{10}$  cfu/g was observed in soil sample (terrestrial) polluted with oil spill dispersant (OSD/SC+Bm+Pa), while the lowest THB count 8.498  $\log_{10}$  cfu/g was recorded in control (TS(CRTL)). The highest THB count was due to increase in hydrocarbon content, which is in agreement with (66, 67). The total petroleum utilizing bacterial count for terrestrial soil ranged from 6.201  $\log_{10}$  Cfu/g to 5.528  $\log_{10}$  Cfu/g. This increase is due the presence of hydrocarbon availability in the soil. This concord with the findings of [Obire and Nwaubeta \(533\)](#), who reported gradual increase in microbial population in contaminated soil. The inoculation of the soil samples on a Mineral salt agar (MSA) supplemented with the test chemical reveal that only 5 out of the 9 bacterial genera were able to degrade and utilize the two test chemicals as their sole carbon and energy source. This is likely to be as a result of the metabolic ability possessed by these oil-degrading bacteria such as *Bacillus spp* and *Pseudomonas spp*. Which enables the utilization of hydrocarbon present in the crude oil as source of carbon and energy (68). Similarly, the result obtained in this study is in line with the report [for Okpokwasili and Nnubia of \(69\)](#) who demonstrated that *Bacillus spp* and *Pseudomonas spp* are of high predominance in hydrocarbon polluted soil as a result of their ability to utilize it. There is an increased and decreased in total dispersant utilizing bacterial in soil samples used as inoculum. The increase could be due to the presence and activities of dispersants utilizers in the soil because, as they utilizes the dispersants, they keep multiplying day by day. The count ranged from 6.045  $\log_{10}$  cfu/g to 7.391  $\log_{10}$  cfu/g. The highest count 7.391  $\log_{10}$  cfu/g was observed in the soil sample polluted with oil spill dispersant (OSD/LT+Bm), while lowest DUB count 6.045  $\log_{10}$  cfu/g was observed in control (TS(CRTL)). This suggests that at the introduction of the treatments, the concentrated dispersants were still very high, which inhibited bacterial growth, as it lethal to it. This observation concord with [author the findings of \(70, 71, 24,72\)](#) that oil spill dispersants support mild increases (stimulation) and decreases (inhibition) in the growth dispersants degraders than hydrocarbon-

degraders and supported the growth of indigenous seawater bacteria confirming that the bacteria could utilize the nutrients available within the dispersants even at low concentrations. [Olusola and Benjamin](#). (73) studied the biodegradability of three dispersants; Pars 1, Pars 2 and Gamlen OD4000. The study showed that, the highest growth of microorganisms was documented for either Pars 1 or Pars 2. Pars dispersants 1 showed more degradability in the first 24 h compared to others, and has more adaptability to the aquatic ecosystem. A consortium of micro-and macro-organisms known as biomass is able to use the dispersant as food. *Pseudomonas aeruginosa* KX828570, *Bacillus megaterium* KY085976, *Micrococcus* sp, *Serratia* sp, and *Proteus* sp were the most predominant dispersant utilizing bacteria found in this study. And this is in line with the findings of (72). Other organisms present in the crude oil polluted soil such as *Streptococcus* spp., *Klebsiella* spp., *Listeria* spp., and *Escherichia coli* were not able to utilize dispersants products, their presence may be due to poor hygienic practices in the community. With these, it shows that some of these isolates were not able to cope nutritionally. Similarity in test chemicals utilization by isolates could be as a result of adequate level of carbon source and phosphate components which has been reported by other investigators to play a role in the test chemical degradation (74, 75). Most of the THB isolated and identified across the two samples belonged to genera; *Pseudomonas aeruginosa* KX828570, *Bacillus megaterium* KY085976, *Micrococcus* spp, *Serratia* spp, *Klebsiella* spp, *Corynebacteria* spp, *Listeria* spp, *Bacillus* spp, and *Proteus* spp. This study reported that *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KY085976 exhibited the best growth and had the highest levels of petroleum degradation. It was observed from the study that these organisms helped in remediating the pollutant caused by oil spill dispersants in the soil with time.

### **Conclusion and Recommendation**

The selection of proper microbial strains is the key step to a successful bioaugmentation. For a pollutant to be eliminated, it is very important to select microbial inoculant isolated from contaminated sites. Bioremediation using *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KY085976 on oil spill dispersants pollution can improve the soil status. Therefore, based on the present research, it can be concluded and recommended that bioremediation of *Pseudomonas aeruginosa* KX828570 and *Bacillus megaterium* KY085976 should be regarded as a key component in the clean-up strategy for oil spill dispersants pollution. Since OSD/LT with mixed consortium are more degradable than OSD/Seacare, its use is more preferable during clean-up of oil spill polluted terrestrial

### **References**

1. Onuoha, S.C, Chukwura, E.I and Fatokun, K. (2014). Stimulated biodegradation of spent lubricating motor oil in soil amended with animal droppings. *American Journal of Biological Science*. 2(1): 19-27
2. Azubuikwe, C.C., Chikere, B.C and Okpokwasili, C.G (2016). Bioremediation techniques-classification based on site of application: principles, advantages, limitations and prospects. *World Journal of Microbiology and Technology*. 32(11): 180.

3. Dadrasnia, A., Salmah, I., Emenike, C.U., and Shahsavari, N. (2014). Remediation of oil contaminated media using organic material supplementation. *Petroleum Science Technology*. 33(9): 1030–1037.
4. Malik, Z.A. and Ahmed, S. (2012). “Degradation of petroleum hydrocarbons by oil field isolated bacteria consortium,” *African Journal of Biotechnology*. 1, 650-658,
5. Jingqiu, Liao, Jie. Wang, Yi. Huang, (2015). Bacterial Community Features Are Shaped by Geographic Location, Physicochemical Properties, and Oil Contamination of Soil in Main Oil Fields of China. *Microbial Ecology*. 70: 380-389.
6. Arezoo, D and Salmah B. I, 2015. Bio-Enrichment of Waste Crude Oil Polluted Soil: Amended with Bacillus 139SI and Organic Waste. *International Journal of Environmental Science and Development*, 6.
7. Frutos, F.J.G., Escolano, O., Garcia, S., Mar Babin, M., and Fernandez, M.D (2010). Bioventing remediation and ecotoxicity evaluation of phenanthrene-contaminated soil. *Journal of Hazard Materials*. 183: 806-813.
8. Sui, H and Li, X (2011). Modelling for volatilization and bioremediation of toluene-contaminated soil by bioventing. *Chinese Journal of Chemical Engineering*. 19: 340-348.
9. Kim, S., Krajmalnik-Brown, R., Kim, J.O and Chung, J (2014). Remediation of petroleum Hydrocarbon-contaminated sites by DNA diagnosis-based bioslurping technology. *Science Total Environment*. 497: 250-259.
10. Firmino, P.I.M., Farias, R.S., Barros, A.N., Buarque, P.M.C., Rodriguez, E., Lopez, A.C and dos Santos, A.B (2015). Understanding the anaerobic BTEX removal in continuous-flow bioreactors for ex situ bioremediation process. *Journal of Chemical Engineering*. 281: 272-280.
11. Nilanjana, D and Preethy, C (2011). Microbial Degradation of Petroleum Hydrocarbon Contaminants: An Overview. *Biotechnology Research International*. Article ID 941810, 13 pages.
12. Pratima, S., Jyoti, S., Shipra, D and Mahendra, K (2014). Bioremediation of Oil Spill. *Journal of Bioscience and Technology*, 5(6):571-581.
13. Van Gestel, K., Margret, J., Swings, J., Coosemans, J., Ryckebore, J., (2001). “Bioremediation of diesel oil contaminated soil by composting with biowaste”. *Environmental pollution*, 125(5): 361 – 368.
14. Gogoi, B. K, Dutta, N. N., Krishnamohn, T. R. (2003). A case study of bioremediation of petroleum hydrocarbon contaminated soil at a crude oil spill site. *Advances in Environmental Research*, 7 (4), 767-782.
15. Nano, G., Borroni, A., Rota, R., (2003). Combined Slurry and Solid-phase bioremediation of diesel contained soils. *Hazardous Materials* 100 (4): 91 – 94.
16. Morelli, A., Fillipi, S. and Zhang, X. H. (2005). Peripheral regulatory mechanism in erection. *International Journal of Androl*. 28(2):23-27
17. Nrior, R. R and Echezolom, C (2016). Assessment of percentage bioremediation of petroleum Hydrocarbon contaminated soil with biostimulating agents. *Journal of International Society of Comparative Education, Science and Technology (ICEST)* 3(1):203-215.

18. Hii, Y.S., Theemlaw, A.H. and Shazili, N.A.M. 2009. Biodegradation of Tapic Blended Crude Oil in Marine Sediment by a Symbiotic Bacteria. *International Biodeterioration and Biodegradation* (63); 142-150.
19. Nrior, R. R, Douglas S. I, and Ojong M.T, 2017. Effect of Waste Water on Biodegradability of Drilling Fluid. *International Journal of Development Research*. Vol. 7(12) pp.17872-17876.
20. Erdogan E.E, Sahin F, Karaca A. (2012). Determination of petroleum-degrading bacteria isolated from crude oil-contaminated soil in Turkey. *African Journal of Biotechnology* 11, 48534859.
21. Diaz, E., (2008). *Microbial Biodegradation: Genomics and molecular Biology*. 1<sup>st</sup> Edition, Caister Academic Press, UK, ISBN: 978-1-90455-17-2, pp:402.
22. Maryam L. Riskuwa-Shehu, and Udemé J.J. Ijah, (2016). "Enhanced Removal of Crude Oil in Soil by Mixed Culture of *Bacillus megaterium* UL05 and *Pseudomonas aeruginosa* UL07." *International Journal of Environmental Bioremediation & Biodegradation*, vol. 4(1): 8-12. doi: 10.12691/ijebb-4-1-2.
23. Vergetis, E (2002). Oil Pollution in Greek sea and spill confrontation means methods, National Technical University of Athens, GREECE.
24. Uzoigwe, C. I. and Okpokwasili, G. C. (2012). Biodegradation of oil spill dispersants in natural aquatic ecosystem. *International Journal of Physical Sciences*, 7(38), 5477-5484.
25. Milinkovitch, T. J. G., M. Théron, H. Thomas-Guyon. (2011). Toxicity of Dispersant Application: Biomarkers responses in gills of Juvenile golden grey mullet (*Lizaaurata*). *Environmental Pollution* 159: 2921-2928.
26. Agbogidi OM, Eruotor PG, Akparabi S O. (2007). Effects of Time of Application of Crude Oil to Soil on the Growth of Maize (*Zea mays* L.) *Research Journal of Environmental Toxicology*.1 (3): 116- 123.
27. Kadaf, A. A. (2012). Environmental Impact of Oil Exploration and Exploitation in the Niger Delta Region of Nigeria. *Global Journal of Science Frontier Research Environment and Earth Sciences* (12);22-24.
28. Tebyanian, H., Hassanshahian, M., and Kariminik, A (2013). Hexadecane-degradation by *teskumurella* and *stenotrophomonas* strains isolated from hydrocarbon contaminated soils. *Jundishapur Journal of Microbiology*. 6 (7), e9182.
29. Nrior, R. R and Onwuka, N. F (2017). Comparative Bioaugmentation Potential of *Pencillium chrysogenum* and *Candida tropicalis* Crude Oil Contaminated Marshland Muddy Soil. *Journal of Environmental Science, Toxicology and Food Technology*. 11 (10) pp 57-64.
30. Obire, O. and Anyanwu, E.C. (2009) Impact of Various Concentration of Crude Oil on Fungal Populations of Soil. *International Journal of Environmental Science and Technology*, 6, 211-218. <http://dx.doi.org/10.1007/BF03327624>
31. Abii, T.A; Nwosu, P. C (2009).The Effect of Oil-Spillage on the Soil of the Eleme in Rivers State of the Niger Delta Area of Nigeria. *Res Journal of Environmental Science* 3(3): 318-320.

32. Trejo-Hernandez, M.R., Oritz, A., Okoh, A.I., Morales, D. and Quintero, R. 2007. Biodegradation of Heavy Crude Oil Maya Using Spent Compost and Sugarcane Bagasse. (68); 848-855.
33. Nrior, R.R., Odokuma, L.O. and Tete, E (2017a). Ultimate Biodegradation of Industrial Detergent Used in the Upstream Sector of the Nigeria Petroleum Industry in Freshwater, Blackish and Marine Water. *International Journal of Ecotoxicology and Ecology*. 2 (4): 134-144.
34. Nrior, R.R., Ogbonna, D.N. and Alabo, A.E. 2017b. Biodegradation of Drilling Fluid Used in the Upstream Sector of the Nigeria Petroleum Industry in Marine Water Environment. *International Journal of Waste Resources*. 7 (4): 302-306.
35. Colwell, R.R and Walker, J.D (1977). Ecological aspects of microbial degradation of petroleum in the marine environment. *Critical Review of Microbiology*. 5:423-445.
36. Obire, O (2018). Microbes: Indicators and Remediators of Polluted Environments. An Inaugural Lecture, 54: 41-90
37. Sabele, J., Vinas, M., Solanas, A. M., (2004). "Laboratory Scale bioremediation experiments on hydrocarbon contaminated soil." *International Journal of Biodeterioration*., 54(3): 19 – 25.
38. Ghazali, F. M., Rahman, R. N. Z. A., Salleh, A. B., Basri, M., (2004). "Biodegradation of hydrocarbons in hydrocarbons in soil by microbial consortium" *International Journal of Biodeterioration and Biodegradation*; 54(5): 61 – 67.
39. Walter, M., Boyd-Wilson, K.S.H., McNaughton, D., Northcott, G., (2005). "Laboratory Trials on the bioremediation of aged pentachlorophenol residue". *Journal of International Biodeterioration & Biodegradation*., 55(3): 121 – 130.
40. Atlas, R. M. and Bartha, R. (2006). "Fate and effect of polluting petroleum in the marine environment" *Residue Review*. 49 (1): 49 – 83.
41. Williams, J.O (2018). Assessment of the physicochemical and microbiological quality of palm oil effluent and soil in Aluu. *Asian Journal of Biology*. 6(3):1-11.
42. Amadi, E.N, Kabar-Klin, D.B, Kpormon, L.B and Robinson, V.K.K (2014). Microbial flora and nutritional composition of adult Palm -Wine Beetle (*Rhychophorus phenicus*). *International Journal of Current Microbiology and Applied Sciences* 3(11):189-192.
43. Nrior, R. R. and Jirigwa, C. C (2017). Comparative bioremediation potential of *Mucor racemosus* and *Paecilomyces variotii* on crude oil spill site in Gio Tai, Ogoni land. *Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT)* e-ISSN: 2319-2402, p- ISSN: 2319-2399. Vol. 11(10): 49-57
44. Cheesbrough, M. 2005. District laboratory practices in tropical countries. Cambridge University Press, United Kingdom. 30-41
45. Nrior, R. R. and Odokuma L.O. (2015) Ultimate Biodegradability potential of Trichloroethylene (TCE) used as Degreaser in marine, brackish and fresh water *Journal of Environmental Sciences, Toxicology and Food Technology (IOSR-JESTET)* www.iosrjournals.org 9:80-89. DOI: 10.9790/240209728089
46. Hamamura, N., Olson, S.H., Ward, D.M. and Inskip, W.P. (2006) Microbial population dynamics associated with crude oil biodegradation in diverse soils. *Applied and Environmental Microbiology*, 72: 6316-6324.

47. Chikere, B.O and Chijioko-Osuji, C.C. (2006). Microbial diversity and physiochemical properties of a crude oil- polluted soil. *Nigerian Journal of Microbiology* 20 (2): 1039-1046.
48. Holt, J.G., Kreig, N.R., Sneath, P.H.A., Stanley, J.T. and Willams, S.T. (1994). *Bergey's Manual of Determinative Bacteriology*-Ninth Edition. Lippincott, Williams & Wilkins, Baltimore
49. Odokuma L.O, Okpokwasili GC (1992). Role of Composition in Degradability of Oil Spill Dispersants, Waste. Management. pp.590- 660.
50. Tanee, FBG and Albert, E (2011). Post-Remediaton Assessment of Crude Oil Polluted Site at Kegbara-Dere Community, Gokana L.G.A. of Rivers State, Nigeria. *Journal of Bioremediation and Biodegradation*, 2:122. doi:10.4172/2155-6199.1000122.
51. Musa, J.J (2014). Effect of Domestic Waste Leachates on Quality Parameters of Groundwater. *Leonardo Journal of Science*, 24: 28-38.
52. Edori, O.S and Iyama, W.A (2017). Assessment of physicochemical parameters of soils from Selected Abattoirs in Port Harcourt, Rivers State, Nigeria. *Journal of Environment and Analytical Chemistry*, 4: 194.
53. Obire, O. and Nwaubeta, O. (2002). Biodegradation of refined petroleum hydrocarbons on soil physicochemical and bacteriological characteristics. *Journal of Applied Science and Environmental Management*. (1) 43-46
54. UNEP (2004). United Nations Environmental Programmes, Analytical Methods for water quality. Burlington publishing house, Ontario, Canada, pp 160. Available at <http://www.gemswater.org> (Accessed 9th June, 2009)
55. Osuji, L.C., and Ezebuio, P.C. (2006). Hydrocarbon contamination of typical mangrove floor in Niger Delta, Nigeria. *International Journal of Environmental Science Technology* 3(3):313-320.
56. Silva-Castro, G.A, Uad, I, Gonzalez-Lopez, J, Fandino, C.G, Toledo, F.L and Calvo, C (2012). Application of selected microbial consortia combined with inorganic and oleophilic fertilizers to recuperate oil-polluted soil using land farming technology. *Clean Technology and Environmental Policy*. 14: 719-726
57. Bento, F.M., Camargo, F.A.O., Okeke, B.C., Frankenberger, W.T (2005)., "Comparative bioremediation of soils contaminated with diesel oil by natural attenuation, biostimulation and bioaugmentation," *Bioresource of Technology*. 96, 1049-1055.
58. Vinithini, O., Sudhakar, S. and Ravikumar, R (2015). "Biodegradation of petroleum and crude oil by *Pseudomonas putida* and *Bacillus cereus*," *International Journal of Current Microbiology and Applied Science*. 4(1). 318-329.
59. Hamme, J.D., Odumeru, J.A., Ward, O.P (2000). "Community Dynamics of a Mixed-Bacterial Culture Growing on Petroleum Hydrocarbons in Batch Culture," *Canadian Journal of Microbiology*. 46, 441-450,
60. Al-Wasify, R.S., Hamed, S.R (2014). "Bacterial biodegradation of crude oil using local isolates," *International Journal of Bacteriology*
61. Tanee, F.B.G and Kinako, P.D.S (2008). Comparative studies of biostimulation and phytoremediation in the mitigation of crude oil toxicity in tropical soil. *Journal of Applied Sciences and Environmental Management* 12: 143-147.

62. Kaboré-Ouédraogo P.W, Savadogo P.W, Ouattara C.A.T, Savadogo A, Traoré AS. (2010). Etude de la Bio-dépollution de Sols contaminés par les Hydrocarbures au Burkina Faso. *Journal de la Société Ouest-Africaine de Chimie* 30, 19-28.
63. Tanee, F.B.G, and Albert, E (2011) Biostimulation potential of sawdust on soil parameters and cassava (*Manihot esculenta*; Crantz) yields in crude oil polluted tropical soil. *Advances in Environmental Biology* 5: 938-945.
64. Eweis, J.B., Ergas, S.J., Chang, D.P.Y., Schroeder, E.D., (1998) "Bioremediation principles," McGraw-Hill Companies, Inc., pp 296,
65. Okerentugba, P.O. and Ezeronye, O.U (2003). Petroleum degrading potential of single and mixed microbial cultures isolated from Rivers and Refinery Effluents in Nigeria. *African Journal Biotechnology*, 2(9): 288-292.
66. Eze, V.C, Onwukar, C.E and Orok, F.E (2014). Microbiological and physiochemical characteristics of soil contaminated with petroleum products in Umuahia, Abia State, Nigeria. *Journal of Applied and Environmental Microbiology*. 2 (6)281-286.
67. Stephen E, Emmanuel O.E, Okpanachi OS, Emmanuel S, Temola OT, et al. (2013) In vitro study of biodegradation of spent lubricating oil by *Aspergillus niger*. *Nature and Science* 11(10): 40-44.
68. Ijah U.J, Abioye O.P (2003) Assessment of physiochemical and microbiological properties of soil, 30 months after kerosene spill. *Journal Research Science Management* 1(1): 24-30.16.
69. Okpokwasili G.C, Nnubia C (1995). Effects of oil spill dispersants and Drilling fluids on substrate specificity of marine bacteria, *Waste. Manage.* 7:515-520.
70. Swannell, R. P. J., Daniel, F. (1999). Effect of dispersants on oil biodegradation under simulated marine conditions. In: Proceedings of the 1999 International Oil Spill Conference, Seattle, Washington. American Petroleum Institute, Washington, DC pp. 166-176.
71. Nrior, R.R. and Wosa, C. (2016). Biodegradation of Oil Spill Dispersant in Brackish Water Ecosystem in the Niger Delta, Nigeria. *Journal of International Society of Comparative Education, Science and Technology (ICEST)*, 3:187-201.11
72. Baghbaderani A.Z, Emtayazjoo M, Porsafa P, Mehrabian S, Bijani S, Farkhani D, Mirmoghtadee P (2012). Effects of three types of oil dispersants on biodegradation of dispersed crude oil in water
73. Olusola, A.O and Benjamin, A.O (2009). Biodegradation of synthetic detergents in wastewater. *African Journal of Biotechnology*, 8 (6): 1090-1109.
74. Okpokwasili, G.C and Olisa, A.O (1991). River-Water Biodegradation of Surfactants in Liquid Detergents and Shampoos. *Water Research* 25: 1425-1429.

UNDER PEER REVIEW

