

Original Research Article

Prevalence and Distribution of Oral Leukoplakia in Patients Attending Oral Medicine Department at Dentistry college in Tishreen University

Abstract

BACKGROUND: To find out the prevalence and distribution of oral leukoplakia in patients who are visiting the Department of Oral Medicine at Tishreen University is necessary to assess oral health and identify the risks of malignant transformation.

OBJECTIVES: The aim of this study was to find out the prevalence and distribution of oral leukoplakia in the patients who visited the Department of Oral Medicine at Tishreen University.

MATERIALS AND METHODS: The study was conducted on 500 patients of the Tishreen Oral Medicine Department at Tishreen University. The number of males was 348 and females 152. The number of who drink alcohol was 117 and non-alcoholic 383. The number of smokers was 279 and non-smokers 221. The average age of the sample was 52 years.

Results: We found that the percentage of leukoplakia in the sample was 2.8%. There was a statistically significant correlation between leukoplakia and smoking, drinking alcohol, Increase in age and sex, and no relation was found with general diseases.

CONCLUSIONS: Increased incidence and associated risk factors (smoking, drinking alcohol, increasing age and sex) require dentists to carefully examine Oral mucosa for early detection of precancerous changes and therefore early treatment.

Leukoplakia, smoking, drinking alcohol

Introduction :

Leukoplakia is "The lesion is often a white lesion on the mucous membrane of the mouth that can not be classified as any other disease". [1] As such, it is not a specific disease in itself, where there is a clinical similarity and variable tissue manifestations. [2] Which are strongly attached to the mucosa and associated with an increased risk of cancer [4,3], the lesion has clear and

variable edges over time [5,3]. Advanced models have developed red spots and there are no other symptoms. [5] Mucosa sometimes though to other parts of the gastrointestinal tract or urinary tract Genitals may be affected [6,7,8].

leukoplakia is a descriptive term that should be launched only after excluding other possible causes. The cause of the episode is not known but the risk factors include smoking chewing tobacco, excessive drinking alcohol, viruses and chronic irritation and the use of nuts [9,3]. It is a pre-cancerous lesion where tissue biopsy generally shows an increase in correlations with or without abnormal cells [5,3] and is mixed with lichen planus , hyperkeratosis, and white candidiasis [3].

Treatment recommendations depend on the clinical appearance and histological examination of the lesion. When abnormal cells are present, simple surgical removal is one possible solution. In other cases, monitoring for periods of three to six months may be sufficient. [3] People are advised to stop smoking and reduce alcohol intake. [3] In half the cases, When smoking continues, 66% of cases Increase thick and white. [5] These cases are more common with age and usually do not occur until after 30. [3] Rates may be as high as 8% in men over the age of 70. Several studies have been conducted on their prevalence and risk factors in several studies in different communities to determine the risk rate, which increases the predictability and ease of treatment

materials and methods :

– The sample consists of 500 patients who visite the Department of Oral Medicine at the Faculty of Dentistry at Tishreen University, who are over 16 years of age. The number of males is 348 and females 152, and 14 cases have been diagnosed as leukoplakia.

- 57 – The number of smokers 279 and non-smokers 221.
- 58 – The number of who drink alcohol was 117 and non-alcoholic 383.
- 59 – The number of people with systemic diseases 101 and the number of non-
- 60 infected 399.
- 61 – The average ages were 52 years, while the age of those infected was between
- 62 49–62 years.
- 63 – A research form was designed in which the researcher recorded the patient's
- 64 personal information (age and gender), Smoking (intensity, duration), Drinking
- 65 Alcohol (Quantity, Frequency, Duration).
- 66 – The existence of general diseases through the use of indirect and directed
- 67 questions.
- 68 – WHO standards for clinical diagnosis to leukoplakia were adopted.
- 69 – The statistical SPSS program was used to analyze the results.

70

71

72 3. Results:

73 3.1 Prevalence leukoplakia of the sample:

Statistics		
leukoplakia		
N	Valid	500
Mean		.028
Std. Deviation		.165
Sum		14

74

Table (3.1) Some descriptive statistics for the rate of leukoplakia

75 The previous table shows some metadata for the variable variable. The sample
76 in question was 500. The number of individuals who had leukoplakia 14 was
77 2.8% and a standard deviation 0.16.

78 3.2 Relationship between leukoplakia and age:

79 The average ages were 52 years, while the age of those infected was
80 between 49 – 62. To test whether there was a relationship between leukoplakia
81 and the age, we used the Point Biserial Correlation Coefficient, which is
82 expressed in Pearson Correlation, next one:

83

Correlations			
		leukoplakia	age
leukoplakia	Pearson Correlation	1	.115*
	Sig. (2-tailed)		.010
	N	500	500
age	Pearson Correlation	.115*	1
	Sig. (2-tailed)	.010	
	N	500	500

84 Table (3.2) Correlation is significant at the 0.05 level (2-tailed).
85 Note from the table that the correlation coefficient value is 0.115. The correlation
86 is linear in the sense that the longer the age, the greater the probability of a
87 coating. Although this coefficient is relatively small, the correlation is significant or
88 significant at the significance level of 0.05 (Sig = 0.01 <0.05).

89 3.3 Relationship between leukoplakia and sex:

90 The number of males who do not have a diploma is 334 and the number
91 of females is 152. However, it should be noted that those who have a class are
92 male only.

93 To study the relationship between sex and class, a correlation coefficient can be
 94 used.

95 * Crosstabulation sex

Count		sex		Total
		male	female	
leukoplakia	Non-leukoplakia	334	152	486
	leukoplakia	14	0	14
Total		348	152	500

96 Table (3.3): Some descriptive statistics on the rate of sex-related relationship

97

Symmetric Measures

	Value	Approximate Significance
Phi	.112	.012
N of Valid Cases	500	

98 Table (3.4) Relationship between leukoplakia and sex using Phi coefficient

99 Note from the above table that the value of the Fay correlation coefficient is
 100 0.112, ie the correlation is positive or negative, which is a significant correlation
 101 (Sig = 0.012 < 0.05).

102 3.4 The relationship between leukoplakia and smoking:

103 leukoplakia was distributed as follows:

104 There are two non-smokers and twelve smokers .

105 To study the relationship between smoking and class, a correlation coefficient
 106 was used.

107 * Smoking Crosstabulation

Total	
Non-smokers	smokers

Leukoplakia	Non-leukoplakia	219	267	486
	leukoplakia	2	12	14
Total		221	279	500

Table (3.5): The relationship between leukoplakia and smoking

Symmetric Measures

	Value	Approximate Significance
Phi	.154	.008
Cramer's V	.154	.008
N of Valid Cases	500	

Table (3.6): The relationship between leukoplakia and smoking using the laboratory Fay
Note from the above table that the value of the coefficient of Fay correlation is 0.154, ie the correlation is positive or negative, which is a significant correlation (Sig = 0.008 <0.05). In the sense that those who are increasing their smoking are more likely to have a class.

3.5 The relationship between leukoplakia and drinking alcohol:

Six people in the sample do not drink alcohol and eight drink . To study the relationship between drinking alcohol and leukoplakia can be used Chi-Square test.

* Alcohol consumption Crosstabulation

		Total		
		Non-alcoholic	alcoholic	
leukoplakia	Non-leukoplakia	377	109	486
	Leukoplakia	6	8	14
Total		383	117	500

Table (3.7) Relation between drinking alcohol and leukoplakia

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)
Pearson Chi-Square	14.890 ^a	2	.001
Likelihood Ratio	9.971	2	.007
Linear-by-Linear Association	13.800	1	.000
N of Valid Cases	500		

Table (3.8) Relation between drinking alcohol and leukoplakia using Chi-Square Test

From the table above, the value of the Chi-Square test index is 14.89 and the test is significant at the significance level of 0.05 (Sig = 0.001 < 0.05). That is, there is a link between drinking alcohol and leukoplakia.

3.6 The relationship between leukoplakia and systemic diseases:

The distribution leukoplakia among sample is as follows: Thirteen people are without systemic diseases and only one is with. To study the relationship between leukoplakia and systemic diseases, the correlation coefficients can be used coefficient of Phi

Crosstabulation

Count		Systemic diseases		Total
		without	with	
leukoplakia	Non-leukoplakia	386	100	486
	leukoplakia	13	1	14
Total		399	101	500

(3.9)

Symmetric Measures

	Value	Approximate Significance
Phi	-.055	.220
N of Valid Cases	500	

(3.10)

Note from the above table that the value of the coefficient of Phi correlation is equal to 0.055, ie, the correlation is weak and is insignificant at the level of 0.05 (Sig = 0.22 > 0.05). It is not clear that those who have leukoplakia have a disease.

4.Discussion:

The prevalence of leukoplakia in our study was 2.8% while it was 0.9% in a study by Reichart et al [10] While a high proportion was observed in Zhang et al. 9.18% [11] and 9.3% in a study conducted by Kumars et al [12] among the tribal population of Kundam province. In a study by Granero et al [13] in Mallorca it was 5,1%and It was 22% at Patil s et al [14].

The difference between prevalence rates in different studies is explained by a number of factors, including: sample size, the nature of the studied society, common habits (smoking and drinking alcohol) and the age of the studied sample, where we see a significant increase in prevalence in studies conducted on older persons. [14] The nature and climate of the region may also play a role.[12]

We found in our study that there was a positive correlation between the prevalence of the leukoplakia and the increase in age. Reichart et al. [15] agreed with us because he studied the German elders to a similar result while R Chandran et al. [16] Kassab et al [17] disagreed with us in a study conducted at the Lebanese University found no difference in the distribution of oral lesions among age groups. Several studies [15] [10] have found a positive correlation between age and leukoplakia. This may be explained by histological changes that occur with increase in age, as well as by prolonged use of oral habits (smoking, drinking alcohol).

In our study, we found that only males were affected by leukoplakia, indicating their association with sex. PA Reichart et al. [10] agreed that males are more affected than females 1.6% to 0.2% .

A study conducted in Budapest by J.Banoczy et al. [10] Where the ratio of males to females was 3.2% to 1% and It also reached the same conclusion Sujathy et al.[19] and Patil S et al. [14] In a study carried out by Cebeci Ar et al. in Ankara [52], the number of men was four times greater than that of women. These results may explain the different oral habits of the sexes (smoking and drinking alcohol) and may be the cause of occupational stress [21] and sex there are no studies to prove a direct relationship.

We found a positive correlation between prevalence Other authors such as Madiyal et al. similar Femopase FI et al. [22] Gary et al.[23] Saraswathi et al. [24] Zhang et al. [11] and Mathewall et al [25]. have also found The agreement between studies on the presence of such a relationship may explain the effect of nicotine on the oral mucous and the changes it causes in mucous membranes. We found that there was a positive correlation between drinking alcohol and leukoplakia, and we agreed with that, Zhang et al. [11], Saraswathi et al [26], and Sujathd et al.[19] and Rooban et al.[27]

While Cebeci Ar et al. [20] did not find a relationship between drinking alcohol and the risk of developing oral lesions. Explanation of the effect of drinking alcohol on the oral mucous where the excessive use of high alcohol, which contains (more than 25%) to the presence of gray board [28].

In our study, there was no relationship between the prevalence of leukoplakia and the presence of systemic diseases , Agreed with us Cebeci Ar et al. [20] This may be due to the low age of the sample and the nature of the studied society while Reichart et al. [15] disagreed with us This may be because he studied German elders with a high proportion of systemic diseases as a result of age.

Conclusions:

The prevalence and distribution of oral leukoplakia were influenced by a range of factors (smoking, drinking alcohol, sex and age), but no association

was found with systemic disease, which should prompt dentists to examine the oral mucous in the most high risk factor groups for early detection of pre-cancerous lesions .

Reference

199

1. Neville BW; Damm DD; Allen CM; Bouquot JE. (2002). Oral & maxillofacial pathology (2. ed.). Philadelphia: W.B. Saunders. pp. 337–345. ISBN 0–7216–9003 .
2. Greenberg MS, Glick M (2003). Burket's oral medicine diagnosis & treatment (10th ed.). Hamilton, Ont.: BC Decker. pp. 87,88,90–93,101–105. ISBN 1–55009–186–7 .
3. Villa, A; Woo, SB (26 October 2016). "Leukoplakia–A Diagnostic and Management Algorithm". Journal of Oral and Maxillofacial Surgery. 75: 723–734. doi:10.1016/j.joms.2016.10.012. PMID 27865803 .
4. Scully, C; Porter, S (Jul 22, 2000). "ABC of oral health. Swellings and red, white, and pigmented lesions". BMJ (Clinical research ed.). 321 (7255): 225–8. doi:10.1136/bmj.321.7255.225. PMC 1118223 . PMID 10903660 .
5. Neville, Brad W.; Damm, Douglas D.; Chi, Angela C.; Allen, Carl M. (2015). Oral and Maxillofacial Pathology (4 ed.). Elsevier Health Sciences. pp. 355–358. ISBN 9781455770526. Archived from the original on 2017–09–10 .
6. Banfalvi, Gaspar (2013). Homeostasis – Tumor – Metastasis. Springer Science & Business Media. p. 156. ISBN 9789400773356. Archived from the original on 2017–09–10 .
7. Wein, Alan J.; Kavoussi, Louis R.; Novick, Andrew C.; Partin, Alan W.; Peters, Craig A. (2011). Campbell–Walsh Urology: Expert Consult Premium Edition: Enhanced Online Features and Print, 4–Volume Set. Elsevier

Health Sciences. p. 2309. ISBN 9781416069119. Archived from the original on 2017-09-10 .

8. Montgomery, Elizabeth A.; Voltaggio, Lysandra (2012). Biopsy Interpretation of the Gastrointestinal Tract Mucosa: Volume 1: Non-Neoplastic (2 ed.). Lippincott Williams & Wilkins. p. 10. ISBN 9781451180589 .

9. Underner, M; Perriot, J; Peiffer, G (January 2012). "[Smokeless tobacco]". Presse Médicale. 41 (1): 3–9. doi:10.1016/j.lpm.2011.06.005. PMID 21840161 .

10. PA Reichart* and H Kohn, Oral Dis dec,2(4):291–4 1996

11. Zhang X1, Li C, Song Y, Reichart PA Oral Maxillaofal Surg dec,14(4):195–202 2008

12. Kumar S1, Muniyandi M , Asian Pac J Canser Rev . 16(4):15–8 2015

13. Granero Fernandez M1, Lopez-Jornet P2 Aust Dent J mar,62(1) :47–51 .2016

14. Patil S1, Doni B2, Maheshwari S3 Can Geriatr J mar31 ,18(1) :11–40 2015

15. Reichart PA et al, Oral Dis nov,9(6):302–4 1996

16. R Chandran, S Meer, L Feller Oral Dis sep,19(6):592–7 2012

17. El Toun S1, Cassia A1, Bouchi N2, Kassab I2. Int J Dent 4030134 2018

18. Department of Conservative Dentistry (Director: Prof. J. Bánóczy, M.D., C.Sc.), Faculty of Dentistry, Semmelweis Medical University, Budapest, Hungary 2005

19. Sujatha D1, Hebbar PB, Pai A.2012 The Oxford Dental College and Hospital, Bangalore, India

20. Cebeci AR1, Gülşahi A, Kamburoglu K, Orhan BK, Oztas B Med Oral Patol Oral Cir Bucal jun1,(416):e272–7 2009

21. Dagli RJ1, Kumar S, Mathur A, Balasubrimanyam G, Duraiswamy P, Kulkarni S Med Oral Patol Oral Cir Bucal nov1,13(11):e687–92 2008

- 249 22.Femopase FL1, Binagui MV, López de Blanc S, Gandolfo M, Lanfranchi HE.
250 Acta Odontol Latinoam 10(2):89–99 1997
- 251 23. GARY CHADMARTIND.D.S., M.P.H.1JOHN P.BROWNB.D.S.,
252 PH.D.2CLAYTON W.EIFLERPH.D.3GLEN D.HOUSTOND.D.S.,
253 M.S.D.1999
- 254 24. Splieth CH1, Sümrig W, Bessel F, John U, Kocher T Quintessence Int.V104
255 2007
- 256 25.Mathewal , Indian J Dent Res apr–jun,19(2):99–103 2008
- 257 26.Saraswathi TR1, Ranganathan K, Shanmugam S, Sowmya R, Narasimhan
258 PD, Gunaseelan R Indean J Dent Res sep,17(3):121–5 .2006
- 259 27.Rooban Thavarajah Arch Oral Biol jun,51(6):512–9 2006
- 260 28.Højgaard, AD; Jessen, AL (Aug 26, 1991). "[Bladder leukoplakia]". Ugeskrift
261 for læger. 153 (35): 2408–9. PMID 1949238 .
262