

Industrialisation scenario at Sreepur of Gazipur, Bangladesh and physico-chemical properties of wastewater discharged from industries

ABSTRACT

This study was conducted to explore the present trend and pattern of industrial growth with spatial distribution of industries and seasonal extent of physicochemical characteristics of wastewater at Sreepur of Gazipur, Bangladesh. The wastewater samples were collected from 5 locations in three seasons viz. pre-monsoon, monsoon and dry season. A total of 120 medium to large industries were surveyed, among those 52 were in red category, 53 were in orange-B category, 13 were in orange-A and only 2 were in green category. In 1995, there were only three industries, which gradually increased to a total of 29 in 2005. Among them, 11 were in red category and 18 were in orange-B category. But from 2006-2010, a total 59 industries were developed and most of them were in red and orange-B categories. Similarly, during the period of 2011 to March 2013, a total 16 industries were developed, among them 8 industries were in red and 3 were in orange-B category. Major types of wastewater discharging industries were textile, dyeing, washing and printing. Among the surveyed industries about 33% didn't have any effluent treatment plant (ETP). The mean value of pH, EC, DO, BOD, COD and TDS of wastewater were 7.28, 2.64, 1.62 mg L⁻¹, 82.0 mg L⁻¹, 217.31 mg L⁻¹ and 1380 mg L⁻¹, respectively during pre-monsoon; 6.7, 1.15, trace, 8.0 mg L⁻¹, 152.4 mg L⁻¹ and 539.58 mg L⁻¹, respectively during monsoon and 7.7, 1.82, 0.74 mg L⁻¹, 48.8 mg L⁻¹, 204.8 mg L⁻¹ and 993.6 mg L⁻¹, respectively during dry season. Average DO concentrations in all seasons and sites were significantly low, while BOD and COD contents were higher in pre-monsoon and dry seasons than the DoE permissible limit. The study concluded that the area is now a hub of polluting industries which are mostly liable to pollute the surrounding environment.

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

Keywords: Industrialisation, Spatial distribution, Wastewater, Sreepur, Gazipur

1. INTRODUCTION

During the last two decades, Bangladesh has experienced a dramatic expansion in small and medium level industries, particularly in garments and textile sector, which have boosted the economy of the country. Undoubtedly, industrialisation plays a significant role to accelerate economic growth and employment status, increase in incomes and standard of living of the people. On the contrary, with the rise in number of industries and expansion of urban areas, the agricultural and residential places are under tremendous pressure in Bangladesh. Therefore, the residents' peoples of such areas are now suffering from various forms of environmental and social hazards. Ironically, environmental degradation in such areas persistently continued despite multiple designated government agencies that are

29 equipped with various conservation laws, codes and planning documents in hand during the
30 past couple of years.

31 Once upon a **time**, Sreepur of Gazipur district has a unique topographical position with rich
32 biodiversity and ecological habitats. But now-a-days farmlands are surrounded by boundary
33 walls and used for different industrial purposes. Beautiful water **bodies came the** carrier of
34 dark, filthy and foul smelled channel. Canals became narrowed down and the polluted water
35 spreading over the farmlands during heavy rain in the rainy season. Furthermore, irrigation
36 practices with these industrial wastewater adds significant quantities of different
37 contaminants including toxic metals which is ultimately damaging the soil quality [1-6].
38 Consumption of agricultural commodities produced in such contaminated soil can cause
39 serious health problems to the **people** [7-9].

40 However, there are **scanty inclusive** research for the Sreepur area in context of industrial
41 pollution. Some of the researches are done sporadically along with areas of other **Upazila's**
42 of same district, without **providing inclusive** result especially for this area [10]. Therefore,
43 detailed systematic field researches on industrialisation scenario and their consequences on
44 water pollution were inadequate or missing. Considering the fact stated **above**, this work was
45 conducted to assess industrialisation scenario, their **categorisation** as environmental
46 pollution sources and physico-chemical properties of wastewater discharged from different
47 industries of Sreepur Upazila of Gazipur district.

48
49

50 **2. MATERIAL AND METHODS**

51

52 **2.1 Description of the Study Area**

53 According to physiographic features Sreepur is an area which evolved during Pleistocene
54 period having area of 465.25 km². The Upazila is located at the north-eastern part of Gazipur
55 district, which lies between 24°01' to 24°20' N latitude and 90°18' to 90°33' E longitude [11].
56 Geologically, the Gazipur cluster lies on the southern corner of Madhupur tract with its
57 average thickness of about 10 **m, which** consists of over consolidated clayey silt and is
58 underlain by the Pleistocene Dupi Tila formation. The rocks encountered here are much
59 younger in geologic age and ranges between Oligocene and Recent time. The basin has got
60 the record of rapid subsidence and sedimentation [12]. Jamindari system was there like
61 other parts of the then Bengal. "Bhawal Raja" estate was there for long time. By virtue of
62 **this**, Jamindari system a number of people of this place historically owned handsome
63 amount of land [13].

64

65 **2.2 Data Collection about the Industries**


66 Data of the industries in the study area **were** collected on the basis of the following pre-
67 structured format, viz. serial no., name of the industry, type of industry, category on the basis
68 of ECR, installation of ETP (yes/no), location, GPS point, establishment year and area
69 covered. In case of any query or **clarification**, industry personnel were asked to reply and
70 sometimes it **was** discussed also with people living nearby industry. Some of the information
71 collected on the basis of oral statement and some of the data collected black and white
72 provided by the industry personnel. Distribution of different types of industries in the study
73 area along with the sampling sites are shown in Fig. 1.

74

75 **2.3 Water Sampling and Processing**

76 **A total of** 5 wastewater samples were collected from the study area during three seasons
77 viz. pre-monsoon, monsoon and dry from different points of the canal following the sampling
78 techniques as outlined by APHA [14]. The collected water samples were stored in 500 mL
79 preconditioned clean, high density plastic bottles and use for the analysis of physicochemical
80 parameters. During collection of water samples, bottles were well rinsed using the same
81 water. All **the** water samples were filtered through Whatman No.1 filter paper to remove

82 unwanted solids and suspended material. After filtration, 3-4 drops of nitric acid were added
83 to the samples to avoid any fungal and other microbial growth. In the laboratory, the samples
84 were kept in a clean, cool and dry place. The locations of the sampling sites have been
85 presented in Fig. 1.
86



87
88 **Figure 1:** Distribution of different types of industries and sampling sites in study area
89
90

91 **2.4 Analytical Methods**

92 The collected wastewater samples were analysed for various physicochemical parameters,
93 which included: pH, electrical conductivity (EC) and total dissolved solids (TDS) were
94 measured within a few hours by using a pH meter (Jenway 3505, UK) and a conductivity
95 meter (SensIONTM+EC5, HACH, USA), respectively. Dissolve oxygen (DO) was determined
96 by Azide modification method, where 2 ml of MnSO₄, 2 ml alkali iodide azid and 2 ml of
97 conc. H₂SO₄ were added as outlined by APHA [14]. Biochemical oxygen demand (BOD) was
98 also determined by Azide modification method, where the samples were kept in a BOD
99 incubator at 20°C for 5 days. The differences between 5 days DO and initial DO was treated
100 as BOD of the water sample. Chemical oxygen demand (COD) was measured by close
101 reflux method using COD vials and measured the concentration by means of a photometer
102 as outlined by APHA [14].
103
104

105 3. RESULTS AND DISCUSSION

106

107 3.1 Spatial Distribution of Industries

108 Over the recent years, Sreepur is experiencing immense pressure of new industrial and
109 commercial establishments. But most of the development activities have done
110 indiscriminately violating the extent environmental laws and ignoring overall public
111 convenience. In the absence of any land zoning system or strict monitoring of land use
112 policy both land developers and entrepreneurs are exploiting the farm lands and using those
113 lands for industrial or commercial purposes.

114

115 In the study area, industrial densities are high in three unions of Sreepur upazila. Among
116 these three unions, industries are mainly located in five mouzas namely- Sreepur, Kewa,
117 Maona, Mulaid and Danua. Most of the industries were developed along the Dhaka-
118 Mymensingh high way and Gorgoria Masterbari-Sreepur road. Major types of industries are
119 textile, dyeing, washing and printing. There were also other types of readymade garments
120 (RMG) industries such as, garments, spinning, sweaters, etc., but they don't release any
121 liquid waste to the surrounding environment or into the canal.

122

123 From Feb-March, 2013, a detailed survey of industries was carried out in the study area. A
124 total of 120 medium to large industries was surveyed in the study area which are shown in
125 Table 1. Among the industries surveyed, 52 were in red category and 53 were in orange-B
126 category and 13 were in orange-A and only 2 were in green category industries (categorised
127 on the basis of ECR [15]). Actual position and type of the major industries are depicted in
128 Fig. 1.

129

130 3.2 Development Scenario and Density of Industries at the Study Area

131 Among the 120 industries surveyed, at least 20 textile dyeing and washing industries were
132 close to the water sampling points. These industries and others also discharge their
133 wastewater to the nearby canal through the pipeline or drain close to each of the sampling
134 points. This pipeline or drain either constructed by the individual industry up to the canal or
135 joined the individual pipeline/drain to a common pipeline/drain by which water ultimately
136 goes to the canal. Different clusters of industries close to the sampling points are shown in
137 Fig. 2. The pipeline or drain networking system so far identified in the field are also shown in
138 Fig. 2 with arrow marks.

139

140 It can be seen from Table 1 that till 1995, there were only three industries in the study area.
141 But, the number of industries gradually increased from 1996 and since then to 2005, a total
142 of 29 industries developed, among them 11 were in red category and 18 were in orange - B
143 categories. But, from 2006 to 2010 the number of industries massively increased in the study
144 area. During this time, a total of 59 industries were developed in the study area, which were
145 mostly in red and orange - B category. During this period 30 red category industries
146 established against 23 orange - B category industries. As the survey was done till March
147 2013, therefore the number of industries from 2011 to March 2013 was not big enough
148 compared to previous time due to the short period of time. This time, a total of 16 industries
149 were developed and among them, 08 industries were in red and 03 industries were in
150 orange - B categories. Therefore, it is a pity to say the study area is now a hub of polluting
151 industries (Fig. 2) which are mostly liable to pollute the environment of the study area. It is to
152 be noted here that out of 120 industries the year of establishment of 13 industries was not
153 known. Most of the industries (49.17%) developed during the period 2006-2010. The
154 majority of the red category and orange-B category industries discharge the liquid waste by
155 their individual pipeline or a common pipeline involving other industries which finally
156 connected to the nearby canal.

157

158
159
160
161
162

Table 1. Detailed information about the industrial establishment in the study area (up to 2013)

SL No.	Name of the Industry	Type	Category (according to ECR [15])	Presence of ETP (Yes/No)	Mouza	GPS point	Estab. year	Area in Acre
1	Unilliance Group	Composite	Red	Yes	Sreepur	24 ⁰ 11'39" 90 ⁰ 21'59.3"	2008	20
2	Hams Garments Ltd.	Knit composite	Red	Yes	Sreepur	24 ⁰ 11'33.8" 90 ⁰ 27'09.1"	2010	2
3	Aman Textile Ltd.	Composite	Red	Yes	Sreepur	24 ⁰ 11'21.3" 90 ⁰ 26'29.5"	2008	13.2
4	AnwaraMannan Textile Mills Ltd.	Spinning	Red	No	Kewa	24 ⁰ 11'02.1" 90 ⁰ 26'30.4"	2012	11.22
5	Ishraque Spinning Mills Ltd.	spinning	Orange-B	No	Sreepur	24 ⁰ 10'49.1" 90 ⁰ 26'14.3"	2006	36.3
6	Chittagong Denim Mills Ltd.	Denim/ Fabrics	Red	Yes	Kewa	24 ⁰ 11'22.6" 90 ⁰ 26'22.4"	2007	7.26
7	Fakruddin Textile Mills Ltd.	Textile composite	Red	Yes	Kewa	24 ⁰ 11'22.8" 90 ⁰ 26'22.5"	2007	13.53
8	Power Mann Limited	Transformer	Red	No	Kewa	24 ⁰ 11'36.0" 90 ⁰ 26'09.0"	2001	1
9	Fashion Makers Ltd.	Fashion	Green	-	Kewa	24 ⁰ 11'44.1" 90 ⁰ 24'10.3"	2010	1.65
10	Skynet Power Company Ltd.	Textile division	Red	No	Kewa	24 ⁰ 11'48.1" 90 ⁰ 26'12.1"	2008	1.81
11	Denimach Washing Ltd.	Washing	Red	Yes	Kewa	24 ⁰ 11'12.1" 90 ⁰ 25'52.1"	2006	17.82
12	Denimach Ltd.	Oven bottom	Orange-B	No	Kewa	24 ⁰ 11'16.2" 90 ⁰ 25.5'51"	2006	3.96
13	Crystal Industries Private Bangladesh Ltd.	Sweater, Jumper	Orange A	No	Sreepur	24 ⁰ 11'17.0" 90 ⁰ 25'47.2"	2010	4.68
14	Mita Textiles Ltd.	Spinning/ Yarn	Orange-B	-	Kewa	24 ⁰ 11'17.0" 90 ⁰ 25'47.3"	1992	19.18
15	How Are You Textile Industries Ltd.	Textile	Red	Yes	Kewa	24 ⁰ 11'11.7" 90 ⁰ 25'40.9"	2006	2.8
16	Meghna Knit Composite Ltd.	Knit composite	Red	Yes	Kewa	24 ⁰ 11'10.2" 90 ⁰ 25'31.2"	2006	3
17	Onetex Ltd.	Yarn/ Dyeing	Red	Yes	Kewa	24 ⁰ 11'04.7" 90 ⁰ 25'22.6"	2003	55
18	Eco-Cotton Mills Ltd.	Cotton	Orange-B	-	Kewa	24 ⁰ 11'04.9" 90 ⁰ 25'18.2"	1996	39.6
19	Sabnam Textile Mills Ltd.	Textile	Orange-B	-	Kewa	2401101.5 9002519.3	1996	2
20	Pandora Sweaters Ltd	Sweater	Orange-B	Yes	Kewa	2401115.0 9002527.9	2005	1
21	Your Fashion Sweater Ltd.	Sweater	Green	-	Kewa	24 ⁰ 11'16.7" 90 ⁰ 25'13.4"	2008	1
22	Welldone Apparel Ltd.	Sweater	Orange- A	-	Kewa	24 ⁰ 11'17.0" 90 ⁰ 24'58.0"	2011	11
23	Perfetti Van Melle Bangladesh Pvt. Ltd.	Candy, Gum	Orange-B	Yes	Kewa	24 ⁰ 10'53.4" 90 ⁰ 24'55.2"	2009	11.38
24	Westeria Textiles Ltd.	Textile	Red	Yes	Kewa	24 ⁰ 10'53.4" 90 ⁰ 25'01.9"	2007	1.23
25	Synergey Textile Ltd.	Textile	Red	No	Kewa	24 ⁰ 10'45.9" 90 ⁰ 25'06.0"	2012	1
26	Integrated Textile Resources Ltd.	Printing	Orange- A	Yes	Kewa	24 ⁰ 10'50.6" 90 ⁰ 25'10.6"	2012	15

SL No.	Name of the Industry	Type	Category (according to ECR [15])	Presence of ETP (Yes/No)	Mouza	GPS point	Estab. year	Area in Acre
27	Dignity Textile Mills Ltd.	Textile	Red	Yes	Kewa	24°10'48.7" 90°25'09.0"	2011	4
28	Argon Denims Ltd.	Denim/ Fabrics	Red	No	Kewa	24°11'25.7" 90°24'57.5"	2008	8
29	Colour and Fashion Industries Ltd.	Sweater	Red	No	Sreepur	24°10'37.9" 90°26'25.7"	2010	2.86
30	Uniglory Cycle Components Ltd.	By-cycle	Red	Yes	Maona	24°10'32.5" 90°25'33.5"	2003	5.61
31	SM Knit Wears Ltd.	Knit composite	Red	Yes	Maona	24°35'35.4" 90°25'27.5"	2000	21.12
32	SM Knitting Industries Ltd.	dyeing	Red	-	Maona	24°10'34.4" 90°25'28.1"	2000	11.68
33	Meghna Cycles Ltd.	By-cycle	Red	Yes	Maona	24°10'37.9" 90°25'24.7"	2010	9.24
34	Aswad Composite Mills Ltd.	Composite	Red	-	Kewa	24°11'48.7" 90°24'45.0"	2008	16.5
35	Shekhor Sweaters	Sweater	Orange-A	-	Kewa	24°11'50.1" 90°24'48.4"	2010	8.25
36	Phoenix Home Textiles Ltd.	Bed sheet	Red	Yes	Kewa	24°11'50.3" 90°24'49.1"	2013	6.6
37	Taqwa Fabrics Ltd.	Knit composite	Red	Yes	Kewa	24°11'44.5" 90°24'25.2"	2009	24.42
38	X Ceramics Ltd.	Tiles	Orange-B	Yes	Kewa	24°11'32.7" 90°24'12.6"	2010	4.95
39	Abcott Industries Ltd.	Cotton (medical)	Orange-B	-	Kewa	24°11'33.3" 90°24'25.2"	2013	13.86
40	Shaharish Composite Towel Ltd.	Home textile	Red	No	Kewa	24°11'43.8" 90°24'15.6"	2012	2.92
41	Knit Horizon Ltd.	Knitting & dyeing	Red	No	Kewa	24°12'01.3" 90°24'02.4"	2011	4.12
42	KSS Knit Composite Ltd.	Knit composite	Red	Yes	Sreepur	24°12'02.1" 90°24'08.1"	2010	4
43	Crown Wool Wear Ltd.	Yarn/ Dyeing	Red	Yes	Maona	24°13'36.1" 90°23'37.5"	2007	12
44	Yasmin Spinning Mills Ltd.	Spinning	Orange-B	-	Maona	24°12'43.1" 90°25'02.6"	2001	15
45	SQ Celsius Ltd.	Sweater	Orange-B	Yes	Kewa	24°12'38" 90°25'02.2"	2002	20
46	Noman Weaving Mills Ltd.	Weaving	Orange-B	-	Maona	24°12'57.7" 90°24'49.1"	2006	22.37
47	Reedisha Knitex Ltd.	Knit & dyeing	Red	Yes	Dhanua	24°15'32.1" 90°23'50.5"	2004	33
48	Hongkong Shanghai Manjela	Spinning	Orange-A	-	Dhanua	24°15'31.1" 90°23'46.4"	1990	14.08
49	Nayanpur Hatchery (Kazi Farms Group)	Hatching	Orange-B	-	Dhanua	24°15'29.6" 90°23'41.2"	*NK	1.81
50	Confidence Knit Wear Ltd.	Knitting	Orange-A	-	Dhanua	24°15'19.7" 90°23'20.8"	2011	5.28
51	Salvo Alkali Chemi- Industry Ltd.	ca Chemical	Red	Yes	Dhanua	24°15'18.4" 90°23'18.9"	2004	7.5
52	Brac Seeds/Feeds	Seed, feed meal	Orange-B	-	Dhanua	24°16'16.0" 90°23'56.7"	2000	4
53	ML Steel Mills Ltd.	Tubes, Furniture	Orange-B	-	Dhanua	24°16'08.1" 90°23'46.5"	2010	42
54	RAK Ceramics (Bangladesh)	Tiles	Orange-B	Yes	Dhanua	24°15'56.2" 90°23'48.1"	2002	7.1
55	RAK Pharmaceu- ticals Private Ltd.	Medicine	Orange-B	Yes	Faridpur	24°15'46.2" 90°23'51.5"	2008	*NA
56	KEA Printing and Packaging Ltd.	Packaging	Orange-A	-	Faridpur	24°15'44.6" 90°23'52.1"	2007	2.3

SL No.	Name of the Industry	Type	Category (according to ECR [15])	Presence of ETP (Yes/No)	Mouza	GPS point	Estab. year	Area in Acre
57	Roshawa Spinning Mills Ltd.	Spinning	Orange-B	-	Dhanua	24°15'45.4" 90°23'50.4"	*NK	8.25
58	Otto Spinning Ltd.	Spinning	Orange-B	-	Faridpur	24°15'26.6" 90°23'56.9"	1998	2.3
59	M and U Cycles Ltd.	By- cycle	Red	No	Dhanua	24°15'24.5" 90°23'55.9"	*NK	85.8
60	HA-MEEM Denim	Denim	Red	Yes	Maona	24°15'12.0" 90°24'00.8"	2007	0.99
61	Century Spinning Mills Ltd.	Spinning	Orange-B	-	Uttarpara	24°15'07.8" 90°24'02.0"	*NK	6.5
62	MIR Ceramics Ltd.	Tiles	Orange-B	Yes	Uttarpara	24°15'06.3" 90°24'06.9"	2003	14.85
63	Jaber Spinning Mills Ltd.	Spinning	Orange-B	-	Uttarpara	24°14'58.3" 90°24'02.2"	2006	0.34
64	Shamsuddin Knitwear Ltd.	Knitting	Orange-A	-	Uttarpara	24°14'59.8" 90°24'02.9"	2013	*NA
65	Summit Uttaranchal Power Company Ltd.	33 MW Power generation plant	Red	No	Uttarpara	24°14'55.3" 90°24'03.1"	*NK	*NA
66	Monica Fashion Ltd.	Garments	Orange-A	-	Mulaid	24°14'42.6" 90°24'08.0"	*NK	9
67	Sufia Cotton Mills Ltd.	Cotton	Orange-B	-	Mulaid	24°14'39.5" 90°24'07.0"	2000	33
68	Nice Denim Mills Ltd.	Denim	Red	No	Uttarpara	24°14'39.6" 90°24'08.5"	2013	6.6
69	Zarba Textile Mills Ltd.	Cotton	Orange-B	-	Uttarpara	24°14'26.9" 90°23'45.7"	2007	13.2
70	Asia Composite Mills Ltd.	Composite	Red	-	Uttarpara	24°14'23.3" 90°23'45.7"	2006	2.64
71	Ashfaq Textiles Ltd.	Textile	Orange-B	No	Uttarpara	24°14'21.1" 90°23'52.3"	2002	5
72	Premiaflex Plastics Ltd.	Plastic materials	Orange-B	--	Uttarpara	24°13'29.5" 90°24'27.1"	2008	6.6
73	Uniglory Cycles Industries Ltd.	By-cycle	Red	No	Mulaid	24°14'20.9" 90°24'13.4"	2009	1.74
74	Meghna Associates Ltd.	Steel Rim	Red	-	Mulaid	24°14'18.0" 90°24'14.1"	2012	5.61
75	Ismail Textile Mills Ltd.	Cotton	Orange-B	No	Mulaid	24°14'21.8" 90°24'31.4"	2008	3
76	Ekota Composite Mills Ltd.	Composite	Red	No	Mulaid	24°14'21.6" 90°24'31.3"	2000	5
77	Haseen Apparels and Knit Composite	Yarn/ Dyeing	Red	Yes	Mulaid	24°14'15.8" 90°24'13.1"	2008	2.51
78	Blue Seal Composite Textile	Composite	Red	Yes	Mulaid	24°14'15.6" 90°24'21.8"	2006	*NA
79	Talha Spinning Mills and Saad-Saan Textile Mills	Spinning	Orange-B	-	Mulaid	24°14'12.3" 90°24'16.5"	NK	5.135
80	Siraj Cycles Industries Ltd.	Tire, tube etc.	Red	No	Mulaid	24°14'08.4" 90°24'18.5"	2010	0.7342
81	Adib Dyeing Ltd.	Knitting/ Dyeing	Red	Yes	Mulaid	24°14'07.0" 90°24'17.5"	2004	3
82	The Welltex Ltd.	Sweater	Orange-A	-	Mulaid	24°14'04.5" 90°24'19.2"	2005	9.9
83	Super Meat Ltd.	Meat processing	Orange-B	-	Mulaid	24°14'02.5" 90°24'19.7"	*NK	1.326
84	Paradise Spinning Mills Ltd.	Spinning	Orange-B	-	Mulaid	24°13'56.2" 90°24'24.3"	*NK	3.63

SL No.	Name of the Industry	Type	Category (according to ECR [15])	Presence of ETP (Yes/No)	Mouza	GPS point	Estab. year	Area in Acre
85	Anwara Knit Composite Ltd.	Composite	Red	Yes	Mulaid	24°13'45.0" 90°24'27.7"	2006	0.41
86	Golden Times Sweater & Dyeing Ltd.	Sweater and dyeing	Orange-B	No	Mulaid	24°13'35.2" 90°24'35.4"	2009	8
87	Viyellatex Spinning Ltd.	Spinning	Orange-B	-	Mulaid	24°13'56.5" 90°24'57.4"	2005	4.62
88	Badar Spinning Mills Ltd.	Spinning	Orange-B	-	Mulaid	24°13'49.3" 90°25'00.3"	*NK	6.6
89	Noman Textile Mills Ltd.	Cotton	orange-B	-	Mulaid	24°13'45.4" 90°24'52.8"	*NK	1.62
90	ABL Design and Fashions	Knit	Orange-B	Yes	Kewa	*RNT	2006	2.8
91	Greenfield Composite Ltd.	Composite	Red	Yes	Kewa	*RNT	NK	21.5
92	Great Wall Ceramic Ind. Ltd.	Tiles	Orange-B	Yes	Kewa	24°11'31" 90°25'39"	2006	15+5.85
93	DIRD Composite Textile	Composite	Red	Yes	Doladia	*RNT	2007	35
94	ACI Formulation Ltd.	Agro-chemicals	Red	Yes	Gojaria	*RNT	1998	10.125
95	Alpha Agro Ltd.	Agro-chemicals	Red	Yes	Atlora	*RNT	2000	4.62
96	FS Sweater Ltd.	Sweater	Orange-B	Yes	Kewa	*RNT	2002	22
97	Nestle Bangladesh Ltd.	Food item	Orange-B	Yes	Baroipara	*RNT	1998	33
98	Paramount Textiles Ltd.	Dyeing	Red	Yes	Gilarchala	24°11'33" 90°25'36"	2009	*NA
99	Health Care Pharmaceuticals Ltd.	Medicine	Orange-B	Yes		*RNT	2000	*NA
100	DaDa Zipper	Dyeing	Red	Yes	Dhanua	*RNT	2008	13.2
101	Padma Paper Mills Ltd.	Dyeing	Red	Yes	Satiabari	*RNT	2008	0.75
102	Organic Health Care Ltd.	Medicine	Orange-B	No	Kewa	24°11'30" 90°25'33"	2007	4.29
103	Vintage Denim Ltd.	Denim/Fabrics	Orange-A	No	Gilarchala	24°11'33" 90°25'29"	2008	26.4
104	Zubair Spinning Mills Ltd	Spinning	Orange-B	-	Gilarchala	24°11'38" 90°25'30"	2008	2.64
105	Miracle Industries Ltd.	Plastic bag	Orange-B	No	Gilarchala	24°11'45" 90°25'28"	1993	2.3
106	Bangladesh Master Pack Ltd.	Plastic bag	Orange-B	-	Gilarchala	24°11'46" 90°25'28"	2007	2.64
107	Package stone Ltd.	Lebel medicine	Orange-B	-	Kewa	24°11'55" 90°25'24"	2003	2.38
108	Sk Sweaters Ltd	Sweater	Orange-A	-	Sreepur	24°12'00" 90°25'30"	2012	14
109	Spring Knit Wears Ltd	Sweater	Orange-A	-	Ansar rd.	24°12'05" 90°25'49"	*NK	3.63
110	Noman Home Textile Mills Ltd.	Fabrics	Red	-	Ansarrd	24°12'03" 90°26'02"	2009	9.5
111	New Hope Feed Mill Bangladesh Ltd.	Cattle/poultry Feed	Orange-B	-	Bhangnahati	24°12'29" 90°27'14"	2008	23
112	Nourish Poultry and Hatchery Ltd.	Poultry	Orange-B	-	Patka	24°11'38" 90°29'28"	2001	2
113	Gentry Pharmaceuticals Ltd.	Medicine	Orange-B	-	Bhangnahati	24°12'27" 90°27'12"	2013	6.6
114	CRC Textile Mills Ltd.	Yarn	Orange-B	-	Bhangnahati	24°12'35" 90°26'48"	2008	*NA

SL No.	Name of the Industry	Type	Category (according to ECR [15])	Presence of ETP (Yes/No)	Mouza	GPS point	Estab. year	Area in Acre
115	Markup Cotton (Square Group)	Cotton	Orange-B	-	Ujilabo	24 ⁰ 12'43" 90 ⁰ 27'05"	2006	13.2
116	Sarah Composite Mills Ltd.	Jute goods	Orange-B	-	Kewa	24 ⁰ 12'37" 90 ⁰ 26'27"	2012	3.63
117	Nikki Thai Aluminium Industries Ltd.	Aluminium	Red	-	Kewa	24 ⁰ 12'49" 90 ⁰ 26'03"	2009	5
118	Nakib Spinning Mills Ltd.	Spinning	Orange-B	-	Kewa	24 ⁰ 12'58" 90 ⁰ 25'34"	2007	8.58
119	MM Spinning Mills Ltd.	Spinning	Orange-B	-	Kewa	24 ⁰ 13'09" 90 ⁰ 25'18"	2006	40
120	Out Pace Spinning Mills Ltd.	Spinning	Orange-B	-	Kewa	24 ⁰ 13'11" 90 ⁰ 25'13"	2009	*NA
Total land								1223.75

163 *RNT = Reading Not Taken; *NA = Not Available; *NK = Not Known and Estab. = Establishment

164

165 3.2 Development Scenario and Density of Industries at the Study Area

166 Among the 120 industries, at least 20 textile dyeing and washing industries were close to the
 167 water sampling points. These industries and others also discharge their wastewater to the
 168 nearby canal through the pipeline or drain close to each of the sampling points. This pipeline
 169 or drain either constructed by the individual industry up to the canal or joined the individual
 170 pipeline/drain to a common pipeline/drain by which water ultimately goes to the canal.
 171 Different clusters of industries close to the sampling points are shown in Fig. 2. The pipeline
 172 or drain networking system so far identified in the field are also shown in Fig. 2 with arrow
 173 marks.

174


175 It can be seen from Table 1 that till 1995, there were only three industries in the study area.
 176 But, the number of industries gradually increased from 1996 and since then to 2005 a total of
 177 29 industries developed, among them 11 were in red category and 18 were in orange - B
 178 categories. But, from 2006 to 2010 the number of industries massively increased in the study
 179 area. During this time, a total of 59 industries were developed in the study area, which were
 180 mostly in red and orange - B category. During this period, 30 red category industries
 181 established against 23 orange - B category industries. As the survey was done till March
 182 2013, therefore the number of industries from 2011 to March 2013 was not big enough
 183 compared to previous time due to the short period of time. This time, a total of 16 industries
 184 were developed and among them, eight (8) industries were in red and three (3) industries
 185 were in orange - B categories. Therefore, it is pity to say study area is now a hub of polluting
 186 industries (Fig. 2) which are mostly liable to pollute the environment of the study area. It is to
 187 be noted here that out of 120 industries the year of establishment of 13 industries was not
 188 known. Most of the industries (49.17%) developed during the period 2006-2010. The
 189 majority of the red category and orange-B category industries discharge the liquid waste by
 190 their individual pipeline or a common pipeline involving other industries which finally
 191 connected to the nearby canal.

192

193 3.3 Status of Effluent Treatment Plant (ETP) of the Existing Industries

194 Among the surveyed industries, 68 needed effluent treatment plant (ETP), but during the
 195 survey it was found 45 installed ETP of different capacity and 23 didn't install ETP.
 196 Therefore, about 33% of surveyed industries didn't have ETP. Installation of ETP has been
 197 made mandatory in liquid waste generating industries by the Department of Environment
 198 (DoE) and DoE is not supposed to issue any environmental clearances to industries running
 199 without ETP. The monitoring team of DoE penalizes the violator following the "polluter pay
 200 principle" and DoE also has made a big change in the law through amending the

201 Environmental Conservation Act, 1995 in the year 2010 and also enacted the Environment
 202 Court Act, 2010 repealing the Environment Court Act, 2000. Amid these manifold stringent
 203 measures **there were still** many industries without ETP. Although some of the industries
 204 have ETP, but either that was not fully functioning or not up to the proper capacity. But there
 205 was also a limitation during the survey, it was not possible to see how many industries ETP
 206 was fully functional or had optimum capacity.
 207 During the period 13 July 2010 to 29 January 2013 through enforcement drive of DoE **some**
 208 **industries of Sreepur area were** penalized for non-compliance of environmental rules and
 209 regulations. **Twelve (12) effluent releasing industries were visited and it was** found that four
 210 (4) had no ETP and **five (5) either had ineffective or closed ETPs**. Among the rest of the
 211 industries, two were dumping solid wastes **on** the nearby agricultural lands and the other
 212 industry was discharging liquid wastes to the environment without mentioning status of ETP
 213 [16]. From the above instances, it can be assumed that in the long run, a good result from
 214 the concept of ETP installation can only be ensured by proper monitoring and environmental
 215 audit by the government. But, the reality is that the success of this initiative could not be
 216 **made** possible alone by the government, the integrity and sincerity of the industrial owner or
 217 the management is most important.
 218
 219



220
 221 Figure 2: Cluster of some industries nearby sampling points along with the direction of
 222 linking drain/ pipeline

223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275

3.4 Physicochemical Characteristics of Wastewater Samples

3.4.1 pH of water sample

Average pH of canal waters for the sites was comparatively high in both pre-monsoon (7.28 ± 0.29) and dry (7.70 ± 0.35) seasons than monsoon season (6.70 ± 0.58) (Table 2). The range of pH was 6.78-7.52 in pre-monsoon season, 6.40-7.30 in monsoon season and 7.30-8.70 in dry season. The higher pH value in pre-monsoon and dry season was because of high base saturation with low volume of water during this period. On the other hand, the pH was slightly low during monsoon season in all sites due to dilution effect. During bleaching and mercerizing processes in textile production processes wastewater produce high pH [17]. The seasonal variation of pH values obtained in this study supports with some other studies. Haque [18] reported that the maximum pH has observed in the winter and minimum in the rainy season.

The pH variation is primarily caused by different kinds of dye stuff used in the dyeing process in different industries. In textile dyeing industries H_2O_2 and NaOH is used as bleaching and kier agents. Higher pH approaches in effluents owing to the waste composition of textile mills such as: NaOCl, NaOH, Na_2SiO_3 , surfactants, sodium phosphate [19]. A study conducted by Moniruzzaman et al. [20] on the water of Buriganga river found that the pH of water was slightly alkaline from December to April (7.4 to 7.6) and the highest average pH value found during the month of April (pH = 7.6). This is due to high base saturations with low volume of water during dry season. On the other hand, the pH of water was slightly lower in wet season from June to October (7.3 to 7.4) than dry season due to dilution effect and the lowest average pH value found during the month of August (pH = 7.3). But all these pH values at different times of year were within the permissible limit. Although there was minor seasonal variation in pH, but all the values were within permissible limit of DoE standard for inland surface water (6.0-9.0) and irrigation water (6.5-8.5).

3.4.2 Electrical conductivity (EC)

Electrical conductivity (EC) is an estimate of the total amount of dissolved ions in water. The EC of water is an indicator of salinity and hazard gives the total salt concentration in water [21]. The mean EC value was comparatively high in pre-monsoon (2.64 ± 0.53 ds/m) and dry (1.82 ± 0.66 ds/m) season than monsoon season (1.15 ± 0.43 ds/m) (Table 2). The range was 2.16-3.30 ds/m in pre-monsoon season, 0.25-1.8 ds/m in monsoon season and 1.14-2.27 ds/m in dry season. In pre-monsoon and dry season EC values in all sites and in monsoon season EC values in two sites were beyond DoE permissible limit (1.2 ds/m). Such a high value of EC is not suitable for aquatic life and irrigation purposes. On the other hand, the EC value was relatively low during monsoon season due to dilution effect, although at three points (1, 2 and 3) EC values were within permissible limit during monsoon season (0.25 , 0.77 and 1.2 mg L^{-1}) but in another two points EC (4 and 5) values were beyond permissible limit (1.8 and 1.74 mg L^{-1}). The lower value of EC of three points might be because of upstream site where accumulation of ions was less than downstream points. The sites 4 and 5 were just adjacent to the industrial effluent releasing point. Therefore, the effluent got less time for effect of dilution and it might be another reason of higher value of EC in these two sites. Another reason might be addition of urban or construction wastes as these two sites were located just adjacent to the roadside where some construction activities also occurred. Furthermore, at site 4 there was an end point of municipal drainage line. Through this line, many pollutants also come out which add to the canal water. Apart from the above reasons these two sites were located downstream level. Therefore, accumulation of substances from different upstream flows to the downstream could be the reason of higher values recorded. The higher values of EC recorded indicates that a large amount of

276 ionic substances were released from the different industries in the study area. A difference in
277 the conductivity in effluent, wastewater or surface water is mainly as a result of difference in
278 the concentration of charged solutes [22]. Haque [18] reported that high tide and winter
279 season have shown the maximum values of EC, and low tide and rainy season have shown
280 the minimum value in the Sundarban area.
281

UNDER PEER REVIEW

282

283 **Table 2: pH, EC and TDS of wastewater discharged from industries of Sreepur, Gazipur at different seasons**

Site	pH of water					EC of water (dS m ⁻¹)					TDS of water (mg L ⁻¹)				
	Pre- monsoon n	Monsoon	Dry	Mean ±SD	Range	Pre- monsoon	Monsoon	Dry	Mean ±SD	Range	Pre- monsoon	Monsoon	Dry	Mean ±SD	Range
1	7.52	7.30	8.70	7.84 ±0.75	7.30 – 8.70	2.96	0.25	2.27	1.83 ±1.40	0.25– 2.96	1512	114.9	1286	970.96 ±749.93	114.9 - 1512
2	6.78	6.60	7.60	6.993 ±0.53	6.6 0– 7.60	2.74	0.77	1.95	1.82 ±0.99	0.77 – 2.74	1384	339	1068	930.33 ±535.92	339 - 1384
3	7.31	6.40	7.60	7.106 ±0.62	6.40 – 7.60	2.04	1.2	2.01	1.75 ±0.47	1.20 – 2.04	1123	580	1080	927.67 ±301.85	580 - 1123
4	7.36	6.50	7.30	7.053 ±0.48	6.50 – 7.36	3.30	1.8	1.74	2.28 ±0.88	1.74 – 3.30	1763	855	933	1183.67 ±503.23	855 - 1763
5	7.42	6.70	7.30	7.14 ±0.38	6.70 – 7.42	2.16	1.74	1.14	1.68 ±0.51	1.14 – 2.16	1118	809	601	842.67 ±260.13	601 - 1118
Average	7.28	6.70	7.70	-	-	2.64	1.15	1.82	-	-	1380	539	993	-	-
SD	0.29	0.35	0.58	-	-	0.53	0.66	0.43	-	-	273.32	313.97	253.13	-	-
Range	6.78 – 7.52	6.40 – 7.30	7.30 – 8.70	-	-	2.16 - 3.30	0.25- 1.8	1.14 – 2.27	-	-	1118 - 1763	114.9 - 809	601 - 1286	-	-
Standard (DoE)	6.0 – 9.0 (Inland surface water) 6.5 – 8.5 (irrigation water)					1.2 dS m⁻¹					2100 mg L⁻¹				

284

285

286

287

Table 3: DO, COD and BOD of wastewater discharged from industries of Sreepur, Gazipur at different seasons

Site	DO of water (mg L ⁻¹)					COD of water (mg L ⁻¹)					BOD of water (mg L ⁻¹)				
	Pre- monsoon n	Monsoon	Dry	Mean ±SD	Range	Pre- monsoon	Monsoon	Dry	Mean ±SD	Range	Pre- monsoon	Monsoon	Dry	Mean ±SD	Range
1	2.81	trace	0.65	1.73 ±1.53	Trace - 2.81	469.14	51.0	288.0	269.38 ±209.69	51.0 - 469.14	50	trace	42	46.0 ±5.66	trace - 50.0
2	1.11	trace	0.47	0.79 ±0.45	trace - 1.11	74.07	56.0	152.0	94.02 ±51.01	56.0 - 152.0	72	trace	38	55.0 ±24.04	trace - 72.0
3	1.12	trace	0.14	0.63 ±0.69	trace - 1.12	74.07	362.0	271.0	235.69 ±147.17	74.07 - 362.0	60	12	70	47.33 ±31.00	12.0 - 70.0
4	1.08	trace	0.85	0.96 ±0.16	trace - 1.08	98.88	184.0	119.0	133.92 ±44.53	98.77 - 184.0	144	4	23	57.0 ±75.95	4.0 - 144.0
5	1.96	trace	1.59	1.775 ±0.26	trace - 1.96	370.37	109.0	194.0	224.45 ±133.32	109.0 - 370.37	84	trace	71	77.5± 9.19	trace - 84.0
Average	1.61	trace	0.74	-	-	217	152.4	204.8	-	-	82	8	48.8	-	-
SD	0.76	-	0.54	-	-	188.35	128.83	73.44	-	-	36.93	5.66	21.04	-	-
Range	1.11 - 2.81	trace	0.14 - 1.59	-	-	74.07 - 469.14	51.0 - 362.0	119.0 - 288.0	-	-	50 - 144	trace - 12	23 - 71	-	-
Standard (DoE)	4.5 – 8.0; ≥ 5 (for irrigation purposes)							200			≤ 10 for irrigation; ≤ 6 for fishing; ≤ 2 for drinking				

288

289

290

291 **3.4.3 Total dissolved solids (TDS)**

292 The TDS values were also comparatively higher in pre-monsoon ($1380 \pm 273.32 \text{ mg L}^{-1}$) and
293 dry season ($993.6 \pm 253.13 \text{ mg L}^{-1}$). Similar to EC, it was relatively lower during monsoon
294 season ($539.58 \pm 313.97 \text{ mg L}^{-1}$). The range was $1118\text{-}1763 \text{ mg L}^{-1}$ in pre-monsoon season,
295 $114.9\text{-}809 \text{ mg L}^{-1}$ in monsoon season and $601\text{-}1286 \text{ mg L}^{-1}$ in dry season (Table 2). The
296 higher TDS value in pre-monsoon and dry season was because of high base saturation with
297 low volume of water during dry and pre-monsoon time. On the other hand, the TDS was
298 lower during monsoon season in all sites due to dilution effect. But, TDS during monsoon
299 season in two sites (4 and 5) was comparatively higher than other three sites as the sites
300 were just adjacent to the industrial effluent releasing point. Therefore, the effluent got less
301 time for effect of dilution. Another reason might be the addition of urban or construction
302 wastes as mentioned earlier. Furthermore, at site 4 there was an end point of municipal
303 drainage line. Through this line many pollutants also came out which adds to the canal
304 water. Apart from above reasons, these two sites were located downstream level. Therefore,
305 accumulation of substances from different upstream flows to the downstream, this could be
306 the reason for higher values of TDS. The result supports the studies done by Haque [18], he
307 found that TDS value increased in the order: rainy season < summer < winter. High TDS
308 elevates the density of water, influences osmo-regulation of fresh water organisms and utility
309 of water for drinking and irrigation purposes. Primary sources for elevated TDS level water
310 pollution discharge from industrial and sewage line, particularly during dry and pre-monsoon
311 season with low water level. Textile, dyeing and printing processes release huge amount of
312 suspended solids and dissolved solid which are mixed in the wastewater during desizing,
313 dyeing and printing stages [17, 23]. Although there was seasonal variation of TDS, all
314 the values were within permissible limit of DoE standard (2100 mg L^{-1}) of Bangladesh.

315

316 **3.4.4 Dissolve oxygen (DO)**

317 The average DO values of wastewater samples were 1.61 ± 0.76 ; $0.74 \pm 0.54 \text{ mg L}^{-1}$ and
318 trace in pre-monsoon, dry and monsoon season, respectively. The range was $1.11\text{-}2.81 \text{ mg}$
319 L^{-1} in pre-monsoon season, $0.14\text{-}1.59 \text{ mg L}^{-1}$ in dry season (Table 3). Adequate DO is
320 necessary for good quality water. As DO levels in water drops below 5.0 mg L^{-1} , aquatic life
321 is put under stress. The lower the concentration, the greater the stress [24]. DO
322 concentrations in all sampling sites were significantly lower than the DoE permissible limit
323 ($4.5\text{-}8.0$ for inland surface water and ≥ 5 for irrigation water) and unsuitable for drinking,
324 fisheries and irrigation purposes. This may be due to high organic and microbial activities
325 with low volume of water. High amount of organic wastes are discharged from textile and
326 dyeing industries into the canal. The dye effluent disposed into the canal water reduces the
327 depth of penetration of sunlight into the water environment, which in turn decreases
328 photosynthetic activity and dissolved oxygen (DO). From the above DO values, it is clear
329 that the water is completely unsuitable for drinking, fishing and irrigation purposes in all
330 seasons in all sampling sites. This result is at par with the findings reported by Zakir et al.
331 [25] for the Mayur river water of Khulna, Bangladesh. Textiles and dyeing mills of the study
332 area release a huge amount of BOD and COD wastes, which consume the DO of water. In
333 natural waters, DO concentration is greatest at 0°C and decreases with increasing
334 temperature. Again, solubility of oxygen decreases with increasing salinity of water [26].

335

336 **3.4.5 Biological oxygen demand (BOD)**

337 BOD is a direct measure of the oxygen uptake in the microbiologically mediated oxidation of
338 organic matter. In other words, it measures the amount of oxygen consumed by an organic
339 compound undergoing decomposition [27]. The BOD average in the study area is relatively
340 higher in pre-monsoon ($82 \pm 36.93 \text{ mg L}^{-1}$) and dry season ($48.8 \pm 21.04 \text{ mg L}^{-1}$) than
341 monsoon season ($8 \pm 5.66 \text{ mg L}^{-1}$). The range was $50\text{-}144$; $23\text{-}71$ and $4\text{-}12 \text{ mg L}^{-1}$ in pre-
342 monsoon, dry and monsoon season, respectively (Table 3). Different steps are followed in

343 the textile processing before the cloth is taken for **bleaching**- it is subjected to kier boiling to
344 remove natural impurities, such as grease, wax, fats, etc. Chemicals used are caustic soda,
345 soda ash, sodium silicate and sodium peroxide. The effluent water from this process is
346 brown in colour and highly alkaline and high in both BOD and COD [28]. Freeman et al. [23]
347 reported that the major pollution indicator parameters for textile wet finishing effluents were
348 the COD, BOD, TDS, suspended solids (SS), colour and heavy metals levels. Wynne et al.
349 [29] stated that textile effluents are highly coloured and saline, contain non-biodegradable
350 compounds, and are high in BOD and COD. Ahmed et al. [30] reported that tannery and
351 textile industries use organic substances as raw materials and high levels of dissolved
352 organic matter consume large amounts of oxygen and increase BOD level, which undergoes
353 anaerobic fermentation processes leading to formation of ammonia and organic acids. High
354 base saturation with low volume of water during dry and pre-monsoon time was the reason
355 behind to increase the BOD in the study area. On the other hand, the BOD is slightly low
356 during monsoon season due to dilution effect. Overall, the BOD value is higher in all sites in
357 all 3 seasons and beyond DoE permissible limit (4.5-8.0 for Inland surface water, ≤ 10 for
358 irrigation and ≤ 6 for fishing). The determined values were not suitable for irrigation, fishing
359 and drinking purposes, **although** some farmers of the area use the canal water in their lands
360 for irrigation purposes.

361

362 **3.4.6 Chemical oxygen demand (COD)**

363 Average COD value was higher in pre-monsoon season ($217 \pm 188.35 \text{ mg L}^{-1}$) and dry
364 ($204.8 \pm 73.44 \text{ mg L}^{-1}$) season than monsoon season ($152.4 \pm 128.83 \text{ mg L}^{-1}$). The range
365 was $74.0\text{-}469.14 \text{ mg L}^{-1}$ in pre-monsoon season, $119.0\text{-}288.0 \text{ mg L}^{-1}$ in dry season and 51.0-
366 362.0 mg L^{-1} in monsoon season (Table 3). COD and BOD are often used to estimate the
367 total quantity of organic matter present in water. Textile industries release a lot of chemical
368 oxygen demanding wastes. The COD levels obtained from garment washing show that
369 detergents, softeners and impurities on the fabrics contributes a significant portion of the
370 COD. The highest COD levels were obtained on dyeing indicating that in addition to fabric
371 impurities removed during scouring or desizing and the contribution of detergents and
372 softeners. Dyes contain high concentrations of salts, and exhibit high BOD/COD values [17].
373 Among the sampling **sites**, 1 and 5 sites during pre-monsoon season and 3 site during
374 monsoon season had COD values excessively higher than other. **At each** of these sites,
375 heavy construction activities were going on during sampling and such activities could
376 **contribute** different types of pollutants into the canal water which might be the reason of
377 increasing COD. Increase **in** organic loadings due to construction activities would increase
378 COD and reduce DO levels [31]. According to Firdissa et al. [32], the mean COD value of
379 effluent from selected industries was significantly above the maximum permissible limit value
380 and effluent with high COD load are released from **beverage followed** by paint, food, soap,
381 tannery, textile and pharmaceutical industry. Sivakumar et al. [33], calculated a ratio of COD
382 : BOD for effluent samples collected from 3 different textile dyeing and bleaching industries,
383 resulting in 1.87, 1.90 and 1.84, respectively. This indicates that these effluents are high in
384 recalcitrant and hardly degradable compounds and may not undergo more than 50%
385 substrate biodegradation, as it is known that organic matter with 50-90% substrate
386 biodegradation has a COD : BOD ratio between 2 and 3.5 [33]. However, on the basis of
387 COD **value**, the canal water in the study area was not suitable for any domestic uses and
388 also not fit for irrigation purposes.

389

390

391 **4. CONCLUSION**

392

393 This study has explored trend and pattern of industrial growth with spatial distribution of
394 industries and seasonal variation in physicochemical properties of wastewater. Total 120
395 medium to large industries was surveyed in the study area of which 52 were in red category

396 and 53 were in orange-B category and 13 were in orange-A and only 2 were in green
397 category. The study revealed that these industries discharge their wastewater into the
398 nearby canal through the pipeline or drain. Number of industries massively increased in the
399 study area during the period 2006-2010 and most of them were in red and orange-B
400 categories. Although installations of effluent treatment plants (ETP) has been made
401 mandatory in liquid waste generating industries by the Department of Environment (DoE),
402 Bangladesh, about 33% of the industries were found to be without ETP. The pH of
403 wastewater collected from the study area was slightly alkaline in pre-monsoon and dry
404 season and near neutral during monsoon season. Average EC values were much higher
405 than DoE standard in pre-monsoon and dry season. TDS value was higher in pre-monsoon
406 and dry season, but comparatively lower in monsoon season although the values were
407 within permissible limit of DoE standard. DO level in wastewater in all seasons was much
408 lower than the DoE standard. BOD and COD values were comparatively higher in pre-
409 monsoon and dry season, and in both seasons average values were much higher than the
410 DoE standard of Bangladesh. Finally, the study results inferred that the area is now a hub of
411 polluting industries which are mostly liable to pollute the surrounding environment.
412 Therefore, to overcome the present situation integrated action plan is necessary.
413 Enforcement and monitoring from the government side alone will not give a concrete solution
414 although political commitment of the ruling government is very important. Thus, action
415 together by the government agencies, non-government organisations and community people
416 will give a fruitful result to make the situation tolerable.

417
418

419 **COMPETING INTERESTS DISCLAIMER:**

420

421 Authors have declared that no competing interests exist. The products used for this
422 research are commonly and predominantly use products in our area of research and
423 country. There is absolutely no conflict of interest between the authors and
424 producers of the products because we do not intend to use these products as an
425 avenue for any litigation but for the advancement of knowledge. Also, the research
426 was not funded by the producing company rather it was funded by personal efforts
427 of the authors.

428
429

430

431 **REFERENCES**

432

- 433 1. Akter A, Zakir HM, Atiqur Rahman, Rumana Yesmeen, Rahman MS. Appraisal of
434 surface water quality for irrigation collected from Sadar upazila of Jamalpur district,
435 Bangladesh. Arc Agric Environ Sci. 2018; 3(3):216-225.
436 DOI:10.26832/24566632.2018.030302
- 437 2. Yesmeen R, Zakir HM, Alam MS, Mallick S. Heavy metal and major ionic contamination
438 level in effluents, surface and groundwater of an urban industrialised city: a case study
439 of Rangpur city, Bangladesh. Asian J Chem Sci. 2018; 5(1):1-16. DOI:
440 10.9734/AJOCS/2018/45061
- 441 3. Zakir HM, Islam MM, Hossain MS. Impact of urbanization and industrialization on
442 irrigation water quality of a canal- a case study of Tongi canal, Bangladesh. Adv Environ
443 Res. 2016; 5(2):109-123. <https://dx.doi.org/10.12989/aer.2016.5.2.109>

- 444 4. Hossain MA, Zakir HM, Kumar D, Alam MS. Quality and metallic pollution level in
445 surface waters of an urban industrialized city: a case study of Chittagong city,
446 Bangladesh. *J Ind Safety Engg.* 2017; 4(2):9-18.
- 447 5. Zakir HM, Islam MM, Arafat MY, Sharmin S. Hydrogeochemistry and quality assessment
448 of waters of an open coal mine area in a developing country: a case study from
449 Barapukuria, Bangladesh. *Int J Geosci Res.* 2013; 1(1):20-44.
- 450 6. Hossain MS, Zakir HM, Rahman MS, Islam MM. Toxic metallic contamination in
451 wastewater of some industrial areas of Mymensingh town, Bangladesh. *Adv Archit City
452 Environ.* 2015; 1(3):7-13.
- 453 7. Aysha MIJ, Zakir HM, Haque R, Quadir QF, Choudhury TR, Quraishi SB, Mollah MZI.
454 Health risk assessment for population via consumption of vegetables grown in soils
455 artificially contaminated with arsenic. *Arch Cur Res Int.* 2017; 10(3):1-12.
456 <https://dx.doi.org/10.9734/ACRI/2017/37612>
- 457 8. Haque R, Zakir HM, Aysha MIJ, Supti Mallick, Rahman MS. Heavy metal uptake pattern
458 and potential human health risk through consumption of tomato grown in industrial
459 contaminated soils. *Asian J Adv Agril Res.* 2018; 5(4):1-11, [https://dx.doi.org/
460 10.9734/AJAAR/2018/40169](https://dx.doi.org/10.9734/AJAAR/2018/40169)
- 461 9. Zakir HM, Aysha MIJ, Mallick S, Sharmin S, Quadir QF, Hossain MA. Heavy metals and
462 major nutrients accumulation pattern in spinach grown in farm and industrial
463 contaminated soils and health risk assessment. *Arch Agric Environ Sci.* 2018; 3(1):95-
464 102, [https://dx.doi.org/ 10.26832/ 24566632.2018.0301015](https://dx.doi.org/10.26832/24566632.2018.0301015)
- 465 10. Zakir HM, Sumi SA, Sharmin S, Mohiuddin KM, Kaysar S. Heavy metal contamination in
466 surface soils of some industrial areas of Gazipur, Bangladesh. *J Chem Bio Phy Sci.*
467 2015; 5(2):2191-2206.
- 468 11. NWPB (The National Web Portal of Bangladesh). District list. Gazipur district. The
469 People's Republic of Bangladesh. 2019. Available: <http://sreepur.gazipur.gov.bd/>
- 470 12. DoE & LGED (Department of Environment & Local Government Engineering
471 Department). Limited Environmental & Social Impact Assessment and Environmental &
472 Social Management Framework. Bangladesh: Dhaka Environment and Water Project.
473 Vol. 1: Main Text. 2010. 1-92 pp.
- 474 13. Sheikh RA. Sreepurer Itihas O Kristhi (in Bengali) (History and Culture of Sreepur),
475 Prokash Printing and Packaging 33/1 Sonargaon road, Dhaka, Bangladesh. 1993.
- 476 14. APHA (American Public Health Association). Standard Methods for the Examination of
477 Water and Wastewater, WEF and AWWA, 20th Edition, USA. 1998.
- 478 15. ECR (The Environment Conservation Rules). Ministry of Environment and Forest. Govt.
479 of the People's Republic of Bangladesh, Dhaka, Bangladesh. August 27, 1997. pp. 179-
480 226.
- 481 16. DoE (Department of Environment). Monitoring and Enforcement Division, DoE, Govt. of
482 the People's Republic of Bangladesh, Dhaka, Bangladesh. Personal communication with
483 director. 2013.
- 484 17. Ramesh BB, Parande AK, Raghu S, Prem Kumar T. Cotton textile processing: waste
485 generation and effluent treatment. *The J Cotton Sci.* 2007; 11:141-153.
- 486 18. Haque MR. Effect of industrial effluent on the mangrove ecosystem of the Sundarbans.
487 Ph.D. thesis, Department of Chemistry, Jahangirnagar University, Savar, Dhaka,
488 Bangladesh. 2004.

- 489 19. Islam MM, Mahmud K, Faruk O, Billah MS. Textile dyeing industries in Bangladesh for
490 sustainable development. *Int J Environ Sci Develop*. 2011; 2(6):428-436.
- 491 20. Moniruzzaman M, Elahi SF, Jahangir MAA. Study on Temporal variation of physico-
492 chemical parameters of Buriganga river water through GIS (Geographical Information
493 System) Technology. *Bangladesh J Sci Ind Res*. 2009; 44(3):327-334.
- 494 21. Brady NC, Weil RR. *The Nature and Properties of Soils*. 12th Ed., Pearson Education,
495 Inc. New Delhi, India. 2002. pp. 261-269.
- 496 22. Trivedi PR, Raj G. *Environmental Water and Soil Analysis*. Akashdeep Publication
497 House, New Delhi, India. 1992. p. 72.
- 498 23. Freeman N, Daniel IO, Pardon K, Kuipa EM, Belaid M. Characterization of effluent from
499 textile wet finishing operations. *Proceedings of the World Congress on Engineering and
500 Computer Science 2009, Vol I, October 20-22, 2009, San Francisco, USA*. 2009.
- 501 24. DEP (Department of Environmental Protection). *Surface Water Quality Standards*.
502 Chapter 62-302, Tallahassee, FL-32399, USA. 2010.
- 503 25. Zakir HM, Sattar MA, Quadir QF. Cadmium pollution and irrigation water quality
504 assessment of an urban river: A case study of the *Mayur* river, Khulna, Bangladesh. *J
505 Chem Bio Phy Sci*. 2015; 5(2):2133-2149.
- 506 26. Boyed CE. *Water Quality Mangement for Pond Fish Culture*. Elsevier Science
507 Publishers B.V. London - Tokyo – New York. 1992.
- 508 27. Hounslow AW. *Water Quality Analysis: Analysis and Interpretation*. CRC Press. 1995.
- 509 28. Azeez PA. Environmental implications of untreated effluents from bleaching and dyeing.
510 In: *Ecofriendly technology for waste minimization in textile industry*. Ed: Senthilnathan S.
511 Centre for Envrionment Education, Tirupur Field Office, Tirupur & Environment Cell
512 Division, Public Works Department WRO, Coimbatore. 2001. pp: 5-11.
- 513 29. Wynne G, Maharaj D, Buckley C. *Cleaner production in the textile industry –Lessons
514 from the Danish experience*. South African Dyers and Finishers Association, Natal
515 Branch Pollution Research Group, School of Chemical Engineering, University of Natal,
516 Durban, South Africa. 2001.
- 517 30. Ahmed MK, Monika D, Islam MM, Akter MS. Physico-Chemical Properties of Tannery
518 and Textile Effluents and Surface Water of River Buriganga and Karnatoli, Bangladesh.
519 *World Appl Sci J*. 2011; 12(2): 152-159.
- 520 31. Environmental impact statement Yalobusha river watershed demonstration erosion
521 control project, Yazooobasin, Mississippi (Final report), US army corps of engineers,
522 Vicksburg district. 2002.
- 523 32. Firdissa B, Solomon Y, Soromessa T. Assessment of the status of industrial waste water
524 effluent for selected industries in Addis Ababa, Ethiopia. *J Natural Sci Res*. 2016;
525 6(17):1-10.
- 526 33. Sivakumar KK., Balamurugan C, Ramakrishnan D, Leena Hebsi Bhai. Assessment
527 studies on wastewater pollution by textile dyeing and bleaching industries at Karur,
528 Tamilnadu. *Rasayan J Chem*. 2011; 4(2):264-269.
- 529
530
531
532