

1
2
3 **Palynological and paleoecological characterization of Upper Eocene-Lower**
4 **Miocene deposits of the southeastern part of the onshore sedimentary basin**
5 **of Côte d'Ivoire (West Africa)**
6
7
8

9 **ABSTRACT**

10
11 Sedimentary rocks cuttings from two boreholes in Bingerville and Assinie (Côte d'Ivoire)
12 were the subject of this study.

13 Sands and clays were collected from the Bingerville well and sands, green clays and
14 limestones from the Assinie well.

15 The main objective of this work is to make an inventory of the plant species that existed at the
16 time of the deposition of sediments on both sides of the lagoon fault based on palynomorph
17 fossils.

18 Paleovegetation consisted of freshwater species such as (determined spores *Verrucatosporites*
19 *usmensis*, *Laevigatosporites ovatus*, *Polypodiaceiosporites regularis*, and *Deltoidospora*
20 *delicata*), which thrived in a coastal wetland environment under a tropical climate with
21 alternating warm and humid periods. Palynostratigraphic analyzes point to the age of the
22 Upper Eocene and the Lower Miocene for the studied samples.
23
24

25 *Keywords: palynomorphs; paleovegetation ; Miocene; Eocene; Bingerville; Assinie.*
26
27

28 **1. INTRODUCTION**

29 The basin of Côte d'Ivoire in which this study is located, is part of a large set of coastal basins
30 bordering the west Atlantic coast from southern Morocco to beyond Angola [1].

31 Cenozoic deposits, contain glauconites and remains of marine organisms, evidence of a
32 transgressive sea, along with pollen grains and spores derived from the land.

33 Palynological studies on the ivorian sedimentary basin began in 1960 with the work of [2],
34 devoted to the Cretaceous deposits.

35 Several other authors contributed to the palynostratigraphical study of the ivorian basin,
36 sometimes on Paleogene and Neogene deposits [3, 4, 5, 6], sometimes Cretaceous [7, 8].

37 Many unpublished dissertation studies (DEA) dissertations have also provided data on the
38 biostratigraphy of Paleogene and Neogene age deposits [9, 10, 11] and upper Cretaceous age
39 [12, 13, 14].

40 The present study was undertaken to date the formations of these two wells made in the
41 Ivorian onshore basin on both sides of the Lagoons fault in order to contribute to the
42 paleobotanic reconstruction of the region which remains enigmatic.
43
44

45 **2. PRESENTATION OF THE STUDY AREA**

46 The study area (Fig. 1) is located southeast of the Ivorian sedimentary basin on both sides of
47 the lagoon fault. Two wells made at Bingerville (P1) and Assinie (P2), the geographical
48 coordinates and depths of which are given in Table 1 below are concerned to this study.

49 The geological history of the sedimentary basin of Côte d'Ivoire is linked to the opening of the
50 South Atlantic, the consequence of which is the dislocation of Gondwana, which intimately
51 united South America and Africa. This story recently recalled by [15] indicates that this basin
52 is characterized by two distinct domains.

53
54 a) a continental domain or onshore basin area affected by a major "lagoon fault" along the
55 coast from west to east. This accident has a vertical discharge of several thousand meters
56 (4000 - 5000 m).


57 (b) a marine domain or offshore basin known only through oil drilling. This offshore basin is
58 subdivided into two margins including the margin of Abidjan and that of San-Pedro.

59

60 **Table 1. Coordinates of the wells**

Site	Location	Longitude (w)	Latitude (N)	Depth in meter
Bingerville	P1	03° 52' 53,8"	05° 20' 06,8"	120
Assinie	P2	03° 24' 02,3"	05° 08' 54,8"	180

61



62
63 **Fig. 1. Location of wells**

64
65 **3. MATERIALS AND METHODS**

66
67 The **studied** materials consisted of twenty-five (25) cuttings from two water wells located at
68 Bingerville (10 samples) and Assinie (15 samples). Each sample was palynologically
69 prepared as practiced in paleobotany laboratories [3].

70 **Procedure** consists of destroying all the mineral phases of the sediment with strong acids
71 (30% HCl and 70% HF) and preserving the organic phase generally consisting of
72 sporopollenic materials.

73 A final attack with nitric acid (HNO₃) 68% cold **in order** to clear the palynological material
74 and organic matter content. After this last attack, the residue is sieved on a 10 µm single-use
75 cloth and then the sporopollenic residue obtained is mounted between the blade and the
76 coverslip using a special resin.

77 Using a biological microscope, observations are made to identify the palynomorphs contained
78 in the slides. These palynomorphs made it possible to date the formations studied and to

79 characterize the paleoenvironment of the region. Paleobotanical analysis is based on the
 80 ecological importance and different botanical affinities of the determined sporomorphs.

81
 82

83 **4. RESULTS**


84
 85 **4.1 Lithological analysis of the wells**

86 **4.1.1 Lithology of the Bingerville well**

87

88 The lithology of cuttings from the well (P1) located in Bingerville shows from the bottom tot
 89 he top: coarse white sand (120 - 97m); sandy variegated clays (97 - 92m); coarse sands (92 -
 90 86 m); compact variegated clays and dark clays (86-44 m); reddish-brown sands (44 - 39 m)
 91 testifying to a strong presence of ferric oxide; very compacted dark clays (39 -25 m) and
 92 yellow-orange laterite clays (25-2 m) (Fig. 2).

93



94

95 **Fig. 2. Schematic lithological column of the P1 well (after [16])**

96

97
98
99
100
101
102
103
104
105
106

4.1.2 Lithology of the Assinie well

The lithological analysis of the cuttings of the Assinie well (P2) shows from older to younger horizons : glauconitic limestones of greenish-gray color with shell debris (180-164 m); intensively green clays, rich in glauconites (164 - 65 m), sandy clays (65-47m); coarse orange-yellow sands, with rare shelly debris (47 - 23 m); medium to fine grained shellfish sands, of a light yellow color rich in bivalve debris (23 - 2 m) (Fig. 3).




Fig. 3. Schematic lithological column of the P2 well (after [16])

4.2 Qualitative and quantitative analysis of palynomorpha from the P1 and P2 wells

The palynomorphs of the well P1 are composed mainly of spores and pollen grains (85%) and scarce dinocysts (15%). The state of conservation of these palynomorphs is excellent. The palynological material of the well P2 is composed of spores and pollen grains (73%) as well as dinocysts (27%). This quantitative study has made it possible to observe many fossil palynomorphs, some of which are of stratigraphic interest.

117 Table 2. Palynomorph Count Sheet for the well P1

DEPTH METER	TOTAL DINOCYSTS	DINOCYSTS				SPORE AND POLLEN GRAIN																		
		TOTAL SPORE AND POLLEN				<i>Batiacospharea</i> sp.	<i>Lingulodinium machaerophorum</i>	<i>Selenopemphix quanta</i>	<i>Operculodinium centrocarpum</i>	<i>Cupressacites hiatipites</i>	<i>Verrucosporites usmensis</i>	<i>Retitricolporites irregularis</i>	<i>Polyadipollenites microreticulatus</i>	<i>Striatopollis catantambus</i>	<i>Retitricolporites</i> sp.	<i>Psilatricolporites crassus</i>	<i>Verrucosporites complanatus</i>	<i>Psilatricolporites laevigatus</i>	<i>Monocolpopollenites</i> sp.	<i>Inaperturopollenites</i> sp.	<i>Magnaperiporites spinosus</i>	<i>Monosulcites</i> sp.	<i>Retimonocolpites irregularis</i>	<i>Laevigatosporites ovatus</i>
30	15									2	6	1	1		1			2	1					1
34	12									1	3	2	2		1			1	1					1
42	23										13	1	2	1	2			1	1					2
47	17	1				1				5	1	1	3	3			1	2						1
53	20	2				2				6	2			2	2	1	3	2	1					1
59	16	1				1				3	2			1	1	2	1	2	1					3
64	16	4	2			2				4	3			1	1	1	1	1	1	1				2
70	20	6	5			1				2	3			3	2	1	1	2	2	2				2
75	18	7	3	1		3				3	1			1	2	1	2	1	2	1	1	1	1	2
94	23	11	6	1		2	2			4	1			4	1	3	2	1	1	2	1	1	1	2
TOTALS		16	2	12	2					3	49	17	6	4	19	9	9	15	14	8	6	2	2	17

118

119

120 Table 3. Palynomorph Count Sheet for the well P2

DEPTH METER	TOTAL DINOCYSTS	DINOCYSTS							SPORE AND POLLEN GRAIN																
		TOTAL SPORE AND POLLEN							<i>Brevicolporites molinae</i>	<i>Laevigatosporites ovatus</i>	<i>Margatricolporites ranvoftii</i>	<i>Deltoispora delicata</i>	<i>Cingulatisporites</i> sp.	<i>Tricolporites</i> sp.	<i>Leiotriletes adriensis</i>	<i>Boculatisporites</i> sp.	<i>Retitricolporites</i> sp.	<i>Verrucosporites usmensis</i>	<i>Pachydermites dietericii</i>	<i>Retitricolporites irregularis</i>	<i>Spinizonocolpites echinatus</i>	<i>Cicatricosporites dorogensis</i>	<i>Polydiacoccolporites regularis</i>	<i>Monipites</i> sp.	<i>Retitricolporites</i> sp.
52	13																								
60	12																								
64	20																								
71	22	2																							
76	14	1																							
82	13	1																							
94	12	1	1	2	1	1	1	1																	
103	17	2	1	1	1	2	1	1																	
112	17	1	2	1	1	1	1	1																	
121	20	1	1	2	2	1	2	1																	
130	25	2	3	1	2	2	1	1																	
139	18	1	1	1	3	3	1	1																	
144	16	1	1	1	2	2	2	2																	
152	17	2	3	2	2	1	1	1																	
165	15		2	1	1	1	1	1																	
TOTALS		15	15	12	15	14	11	10																	

121

122

123 4.3 Palynostratigraphy

124

125 ➤ Well P1

126 Palynological analysis of the Bingerville well (P1) revealed two stages, defined by
127 associations composed mainly of spores and pollen grains and rare dinocysts (Fig. 4).

128

129 - The **upper horizon** ranges from 25 m to 51 m and is characterized by the following
130 spores and pollen grains: *Cupressacites hiatipites*, *Laevigatosporites ovatus*,
131 *Polyadopollenites microreticulatus*, *Psilatricolporites laevigatus*, *Striatopollis*
132 *catatumbus*, *Retitricolporites irregularis*, *Verrucatosporites usmensis*, *Retitriporites* **sp.**
133 and *Monocolpopollenites* **sp.**

134

135 - The **lower horizon** ranges from 51 m to 120 m is marked by species of spores and pollen
136 grains such as: *Psilatricolporites crassus*, *Verrustephanocolporites complanatus*,
137 *Retitricolporites irregularis*, *Verrucatosporites usmensis*, *Retimonocolpites irregularis*.
138 These spores and pollen grains are associated with the following dinocysts:
139 *Selenopemphix quanta*, *Batiacasphaera* **sp.**, *Spiniferites ramosus* and *Cleistosphaeridium*
140 *flexuosum*.

141

142 ➤ Well P2

143 Palynological analysis of the P2 well also highlighted two stages as well (Fig. 5).

144

145 - The **upper horizon** range from 47 to 85 m is revealed by the palynological association
146 composed of spores and pollen grains such as *Laevigatosporites ovatus*, *Leiotriletes*
147 *adriennis*, *Polypodiaceosporites regularis*, *Polypodiisporites speciosus*, *Cingulatisporites*
148 **sp.**

149 - The **lower horizon** extends from 85 to 180 m and is marked by spores and pollen grains
150 such as *Pachydermites diderixii*, *Retitricolporites irregularis*, *Spinizonocolpites*
151 *echinatus*, *Cicatricosporites doregensis*, *Margotricolporites rauwolfii*, *Verrucatosporites*
152 *usmensis*. To these spores and grains of pollen are associated dinocysts such as
153 *Cometodinium obscurum*, *Spiniferites ramosus*, *Operculodinium* **sp.**,
154 *Batiacasphaera* **sp.**, *Cordosphaeridium inodes*, *Isabelidium* **sp.** and *Lingulodinium*
155 *machaerophorum*.

156





Fig. 4. Vertical distribution of the main Bingerville palynomorphs (P1)

157
158
159
160
161
162



163

164 Fig. 5. Vertical distribution of the main Assinie palynomorphs (P2)

165 **4.4 Paleobotanical characterization**

166

167 The paleobotanical study of these two wells shows the presence of pollen grains from the
 168 Arecaceae (*Retitricolporites irregularis*, *Monocolpopollenites* sp.), Fabaceae (*Striatopollis*
 169 *catatumbus*), Schizeaceae (*Inaperturopollenites* sp.), Pelliceria (*Psilatricolporites crassus*),
 170 Nypa (*Spinizonocolpites echinatus*, *Retimonocolpites irregularis*), Apocynaceae
 171 (*Margotricolporites rauwolfii*, *Brevitricolporites molinae*). These pollen grains are associated
 172 with spores of Polypodiaceae (*Laevigatosporites ovatus*, *Verrucatosporites usmensis*,
 173 *Polypodiaceosporites regularis*), Schizeaceae (*Cicatricosporites dorogensis*, *Leiotriletes*
 174 *adriennis*), to Cyatheaceae (*Deltoidospora delicata*) and to Lygodium (*Crassoretitriletes*
 175 *vanraadshooveni*).

176 Palynoflora consists of angiosperm pollen grains typical for tropical rainforests and coastal
 177 swamps (*Pachydermites diderixii*, *Retitricolporites irregularis* and *Striatapollis catatumbus*),
 178 ancestors of the present-day palm trees of the genus Nypa (*Spinizonocolpites echinatus*,
 179 *Retimonocolpites irregularis*), fern spores basically hygrophilous freshwaters that develop in
 180 moist, swampy areas (*Laevigatosporites ovatus*, *Verrucatosporites usmensis*,
 181 *Polypodiaceosporites regularis*).

182 This palynoflora indicates a tropical paleoclimate with alternating warm and humid periods.
183 The association of coastal marine ecosystems (*Cordosphaeridium inodes*, *Spiniferites*
184 *ramosus*) with this paleovegetation indicates a coastal marine ecosystem in this area.

185
186

187 5. DISCUSSION

188

189 5.1 Palynostratigraphy

190

191 Palynological analysis revealed lower Miocene and upper Eocene age of the studied samples.
192 Lower Miocene age has been identified through associations of *Cupressacites hiatipites*,
193 *Laevigatosporites ovatus*, *Polyadopollenites microreticulatus*, *Psilatricolporites laevigatus*,
194 *Striatopollis catatumbus*, *Retitricolporites irregularis*, *Verrucatosporites usmensis*,
195 *Leiotriletes adriennis*, *Polypodiaceoisporites regularis*, *Retitriporites sp.*

196 Our results are consistent with those of [17, 18, 19], who used some of these sporomorphs
197 respectively in Soudan and Côte d'Ivoire to determine the lower Miocene age of palynomorph
198 assemblage.

199 The species *Crassoretitriletes vanraadshooveni* extends from Miocene to Pliocene in Nigeria
200 [20] and from the Middle Miocene to the Pleistocene in Venezuela [21]. As for
201 *Verrucatosporites usmensis*, it characterizes the Eocene to Pleistocene interval in Nigeria and
202 Borneo [20, 22].

203 *Laevigatosporites ovatus* is known from the Neogene in Burundi [23] and Paleogene in
204 Nigeria [24].

205 *Striatopollis catatumbus* characterizes the Paleocene-Pleistocene interval in Nigeria [20] and
206 the Pleistocene-Eocene range in Venezuela [21].

207 *Brevicolporites molinae* marks the Oligocene and the Lower Miocene in Cameroon [22] and
208 the Miocene in Soudan [17].

209 The species *Retitriporites sp.* is a good marker of the Upper Oligocene and the Lower
210 Miocene in Soudan [17]. However, the absence of *Lejeunecysta* (good marker of the
211 Oligocene in Côte d'Ivoire) [7] in this interval restricts this age to the lower Miocene.

212 The Upper Eocene age was determined due to the associations of *Psilatricolporites crassus*,
213 *Verrustephanocolporites complanatus*, *Retitricolporites irregularis*, *Verrucatosporites*
214 *usmensis*, *Retimonocolpites irregularis*, *Pachydermites diderixii*, *Spinizonocolpites*
215 *echinatus*, *Cicatricosporites dorogensis*, *Margotricolporites rauwolfii*.

216 Results can be compared [25, 26, 27, 28] who described such palynomorph assemblage from
217 the Upper Eocene in the Cameroun Basin. To these spores and pollen grains are associated
218 dinocysts such as *Cometodinium obscurum*, *Spiniferites ramosus*, *Operculodinium*
219 *centrocarpum*, *Batiacasphaera sp.*, *Cordosphaeridium inodes*. [29], considers the species
220 *Cordosphaeridium inodes* as an indicator of the Eocene in Germany, while [30] attributes it to
221 the Middle Oligocene in Australia.

222 The species *Spinizonocolpites echinatus* last appears in the Upper Eocene as stated in many
223 works [20, 27, 31, 32, 33] in Nigeria, Cameroun, Soudan and Ghana.

224 *Psilatricolporites crassus* characterizes the Upper Paleocene and Lower Eocene. In
225 Cameroun, [27] identified it in the Lower Eocene and Middle Eocene. In Nigeria this species
226 has been used by [20] to characterize the late Pliocene-Pleistocene interval. In South America,
227 this species characterizes the Lower to Middle Eocene [31, 34].

228 The species *Pachydermites diderixi* present in this stage characterizes the Eocene and
229 Miocene in Cameroon [27], Oligocene and Miocene in Soudan [17].

230 However, the presence in this stage of *Lingulodinium machaerophorum*, an Eocene marker in
231 Egypt [35] and *Cordosphaeridium inodes* known from the Maastrichtian to Upper Eocene [7,
232 24, 32, 36, 37] restricts this age to the Upper Eocene.

233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276

5.2 Paleocology

Paleobotanically, our work is in agreement with results of [19], considering the assemblage composed of *Verrucatosporites usmensis*, *Retitricolporites irregularis*, *Laevigatosporites ovatus*, *Leiotriletes adriennis*, *Pachydermites diderixii*, *Polypodiaceoisporites regularis* as a characteristic of tropical hot and humid climate.

The presence of the pollen grain *Brevitricolporites molinae* (Apocynaceae) typical of tropical forests [22] is confirmed in our work.

In addition, the results of [38] in conformity with ours reveal that fern spores such as *Laevigatosporites ovatus*, *Leiotriletes adriennis*, and *Verrucatosporites usmensis* indicate a humid tropical climate. This author also states that the species *Psilatricolporites crassus* is a pollen grain from mangrove vegetation which has been verified by our work.

The results of [39] reported by [40] indicate, as in our work, that Polypodiaceae (*Polypodiaceoisporites regularis*) are derived from tree ferns that indicate a thick and closed tropical forest.

For [40, 41], the genus *Striatopollis catatumbus* encountered in our formations is a species of freshwater and coastal swamps. These results are verified by our work. These authors also claim that they can be found in the coastal plains as well as in tree savannas.

Similarly, our work is verified by results [42]. They claim that dinocysts such as *Operculodinium centrocarpum*, *Spiniferites ramosus*, *Cordosphaeridium inodes* and *Batiacasphaera* sp. indicate a marine depositional environment near the coast.

6. CONCLUSION


The palynostratigraphic and paleoecological study the plant fossil from the two wells of Bingerville and Assinie reveal the age and the depositional environment of the studied sample.

Dark, variegated sand and clays occur in the Bingerville well, while bioclastic sand, glauconite green clay and limestone in the assinie well.

Green clays contain remains of marine organisms, evidence of a transgressive sea at this time. The palynostratigraphic analyzes revealed a palynoflora characterizing the Upper Eocene and the Lower Miocene. Paleovegetation reveals the presence of species that develop in a mangrove environment with moist, lowland, partly marshy forest in a tidal estuarine coastal environment.

277

278



279

280 *1.Cupressacites hiatipites*; 2.*Laevigatosporites ovatus* ;3.*Polyadopollenites microreticulatus* ; 4.
281 *Psilatricolporites laevigatus* ;5. *Striatopollis catatumbus* ;6. *Retitricolporites irregularis*; 7. *Verrucatosporites*
282 *usmensis* ;8. *Retitriporites* sp. ; 9. *Monocolpolleniites* sp.

283


284 **Fig. 6. Spores and pollen grains from the Lower Miocene of Bingerville (from [4])**

285

286

287


288



289
 290 1. *Brevicolporites molinae* ; 2. *Cingulatisporites* sp. ; 3. *Verrucatosporites usmensis* ; 4. *Laevigatosporites*
 291 *ovatus* ; 5. *Leiotriletes adriennis* ; 6. *Polypodiaceoisporites regularis* ; 7. *Baculatisporites* sp. ; 8.
 292 *Margotricolporites rauvolffii* ; 9. *Striatopollis catatumbus*
 293

294 **Fig. 7. Spores and pollen grains from the Lower Miocene of Assinie (from [4])**

295
 296
 297
 298




299

300 1. *Psilatricolporites crassus* ; 2. *Inaperturopollenites* sp. ; 3. *Verrustephanocolporites complanatus* ; 4.
 301 *Magnaperiporites spinosus* ; 5. *Verrucatosporites usmensis* ; 6. *Monosulcites* ; 7. *Retimonocolpites irregularis* ;
 302 8. *Laevigatosporites ovatus* ; 9. *Retitricolporites irregularis* ; 10. *Retitriporites* sp. ; 11. *Monocolpopollenites* ;
 303 12. *Retitricolpites* sp.

304 **Fig. 8. Spores and pollen grains from the Upper Eocene of Bingerville (from [4])**

305


306



307
 308 1. *Verrucatosporites usmensis* ; 2. *Baculatisporites* sp. ; 3. *Polypodiaceoisporites regularis* ; 4. *Tricolpites* ;
 309 5. *Retitripites* sp. ; 6. *Spinizonocolpites echitanus* ; 7. *Laevigatosporites ovatus* ; 8. *Momipites* sp. ; 9.
 310 *Deltoidospora delicata* ; 10. *Cingulatisporites* sp. ; 11-13. *Pachydermites diderixii* ; 12. *Retitricolporites*
 311 *irregularis* ; 14. *Cicatricosporites dorengensis*.

312 **Fig. 9. Spores and pollen grains of the Upper Eocene of Assinie (from [4])**

313



315

316

317 1. *Lingulodinium machaerophorum* ; 2. *Operculodinium centrocarpum* ;3. *Selenopemphix quanta* ; 4

318

318 *Batiacasphaera* sp.;

319

320 **Fig. 10. Dinocysts of the Upper Eocene of Bingerville (from [4])**

321

322

323

324

325


326

327

328

329

330



331
 332 1. *Comotodinium obscurum*; 2. *Isabelidinium* sp.; 3-8. *Operculodilium centrocarpum*; 4. *Batiacasphaera* sp.; 5-
 333 9. *Spiniferites ramosus*; 6. *Cordosphaeridium inodes*; 7. *Lingulodinium machaerophorum*
 334

335 **Fig. 11. Dinocysts of the upper Eocene of Assinie (from [4])**

336

337

338

339

340 **APPENDIX**

341

342 **Spore and pollen grains**

- 343 *Baculatisporites* sp. (Jaramillo & Dilcher, 2001)
344 *Brevicolporites molinae* (Schuler & Doubinger 1970) Salard-Cheboldaeff 1978
345 *Cicatricosisporites doregensis* (Potonié&Gelletich, 1933)
346 *Cingulatisporites* sp.
347 *Cupressacites hiatipites* (Wodehouse,1933) Krutzsch, 1971
348 *Deltoidospora delicata* (Sah, 1967)
349 *Inaperturopollenites* sp.
350 *Laevigatosporites ovatus* (Wilson & Webster, 1947)
351 *Leiotriletes adriennis* (Krutzsch, 1959)
352 *Magnaperiporites spinosus* (Gonzalez, 1967)
353 *Margotricolporites rauwolfii* (Salard-Cheboldaeff, 1978)
354 *Monocolpollenites* sp.
355 *Monosulcites* sp.
356 *Pachydermites diderixii* (Germeraad, & Muller, 1968)
357 *Polyadopollenites microreticulatus* (Salard, 1974)
358 *Polypodiaceoisporites regularis* (Zhang, 1981)
359 *Psilatriporites* sp.
360 *Psilatricolporites crassus* (Van der Hammen & Wijmstra 1964)
361 *Psilatricolporites laevigatus* (Van der Hammen and Wijmstra, 1964)
362 *Retimonocolpites irregularis* (Van der hammen & Wijmstra 1964)
363 *Retitricolpites* sp.
364 *Retitricolporites irregularis* (Van de Hammen & Wijmstra, 1964)
365 *Retitriporites* sp.
366 *Spinizonocolpites echinatus* (Muller, 1968)
367 *Striatopollis catatumbus* (González Guzmán, 1967) Ward, 1986
368 *Tricolpites* sp.
369 *Verrucatosporites usmensis* (Van der Hammen, 1956) Germeraad et al., 1968
370 *Verrustephanocolporites complanatus* (Salard-Cheboldaeff, 1978)

371

372

373

374

375 **Dinocyst**

- 376 *Batiacasphaera* sp. (Jaramillo & Dilcher, 2001)
377 *Cometodinium obscurum* (Deflandre & Courteville, 1959) Monteil, 1991
378 *Cordosphaeridium inodes* (Klumpp, 1953) Eisenack, 1963
379 *Isabelidinium* sp.
380 *Lingulodinium machaeropharum* (Deflandre and Cookson, 1955) Wall, 1967
381 *Operculodinium centrocarpum* (Deflandre & Cookson, 1955) Wall, 1967
382 *Selenopemphix quanta* (Bradford, 1975) Harland, 1981
383 *Spiniferites ramosus* (Ehrenberg, 1838) Mantell, 1854

384

385

386

387

388 **REFERENCES**

389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435

1. Aka K. Quaternary sedimentation on the margin of Côte d'Ivoire: modeling test. PhD thesis Nat. Sc., Abidjan University. 1991; 233.
2. Jardine S, Magloire L. Palynology and stratigraphy of Cretaceous basins of Senegal and Ivory Coast. Memory of the Office of Geological and Mining Research. 1965; 187-245.
3. Bie GR. Evolution of the microflora of the sedimentary basin of Côte d'Ivoire (Abidjan margin) during the cenozoic: palynostratigraphy, paleobotany, evolution of depositional environments and maturation of organic matter. PhD Thesis, Univ. Felix Houphouët Boigny. 2012; 218.
4. Gbangbot JMK. Stratigraphic characterization of aquifers of subsurface formations of the Ivory Coast lagoons region. Modeling test of Tertiary deposit environments. PhD Thesis, Univ. Felix Houphouët Boigny. 2012; 196.
5. Assale FYP. Sedimentological, palynological, geochemical and palaeoenvironmental characterization of sedimentary formations associated with the lagoon fault (eastern Côte d'Ivoire onshore basin). PhD Thesis, Univ. Felix Houphouët Boigny. 2013; 441.
6. Behi ZDA. Litho-biostratigraphy of neogenic deposits associated with the lagoon fault (Ivorian onshore basin): palaeoecological and palaeoenvironmental reconstruction. PhD Thesis, Univ. Felix Houphouët Boigny. 2017; 210.
7. Digbehi ZB, Doukoure M, Tea YJ, Yao KR, Yao NJP, Kangah KD, Tahi I. Palynostratigraphy and Palaeoenvironmental Characterization and Evidence of Oligocene in Terrestrial Sedimentary Basin, Bingerville area, Southern Ivory Coast, Northern Gulf of Guinea. "African Journal of Environmental Science and Technology". 2012; 28-42.
8. Guede KE. Comparative study of palynoflora (dinoflagellate cysts) at the Cretaceous-Paleogene (K-Pg) and Paleocene-Eocene (P-E) passages of northwestern Morocco and southwestern Côte d'Ivoire: Systematic, Biostratigraphy, Palaeoenvironments and Paleobiogeography. Ph.D. thesis, Mohammed V University of Rabat, Morocco. 2016; 341.
9. Kangah KD. Palynostratigraphy of the K1-1X well, Cretaceous-Tertiary passage. Memory of DEA. 1997; 57.
10. Ennin TM. Sedimentary and palynological study of four wells in the Brégbo region (South-East of Abidjan). DEA of Earth Sciences option Marine Geology and Sedimentology, UFR STRM, University of Abidjan. 2003; 58.
11. Doukoure M. Biostratigraphy of tertiary deposits in the Bingerville area. DEA of Earth Sciences option Marine Geology and Sedimentology, UFR STRM, Univ. Cocody (Abidjan). 2006; 59.

- 436 12. Zahoui DHB. Sedimentological and biostratigraphic characterization of oligostéginid
437 limestones in the Ivorian sedimentary basin. DEA Earth Sciences option, Marine Geology,
438 UFR STRM, Univ. Cocody (Abidjan). 2003; 70.
439
- 440 13. Toe Bi KKK, Yao NJP, Kesse TM, Digbehi ZB. Sedimentological and hydrodynamic
441 characterization of the lower Miocene sandy formations of the Eboinda region (South-East of
442 Côte d'Ivoire). European Scientific Journal. 2016; 12 (9): 192-211.
443
- 444 14. Guede KE. Palynostratigraphic and palaeoenvironmental characterization of the
445 Cretaceous-Tertiary and Eocene passage formations in the study of the DINO-1X offshore
446 well. DEA earth sciences option Marine Geology and Sedimentology, UFR STRM,
447 University of Cocody, Abidjan. 2009; 78.
448
- 449 15. Bamba MK, Digbehi ZB, Sombo BC, Goua TE & N'Da LV. Planktonic foraminifera,
450 biostratigraphy and palaeoenvironment of Albo-Turonian deposits in Côte d'Ivoire, West
451 Africa. Journal of Paleobiology, Geneva. 2011; 30 (1): 1-11.
452
- 453 16. Gbangbot JMK, Digbehi ZB, Yao NJ, Monde S, Yao NA. Lithostratigraphy of the
454 subsurface deposits of the Bingerville and Assinie Regions, South and South-East, Lower
455 Ivory Coast. Comparison test of deposit environments during the Tertiary. European Journal
456 of Scientific Research. 2012 ; 86 (1) : 41-52.
457
- 458 17. Eisawi A, Schrank E. Upper Cretaceous to Neogene palynology of Melut Basin, Southeast
459 Sudan. Palynology. 2008; 32 (1): 101-129.
460
- 461 18. Toe Bi KKK. Evolution and characterization of subsurface sediments and microflora in
462 the Eboinda region (southern zone of the lagoon fault): sedimentology, geochemistry,
463 biostratigraphy, palaeoenvironment and paleobiogeography. PhD Thesis, Univ. Felix
464 Houphouet Boigny. 2016 227.
465
- 466 19. Behi ZDA, KE Guede, Toe Bi KKK, Kouassi KA, Digbehi ZB. Palynostratigraphy and
467 Paleobotany of Lower Miocene Deposits in South-East Côte d'Ivoire, West Africa. Isee. Sci.
468 Technol. 2018; 32: 331-349.
469
- 470 20. Germeraad JH, Hopping CA, Muller J. Palynology of Tertiary sediments from tropical
471 areas. Review of Palaeobotany and Palynology. 1968 6: 189-348.
472
- 473 21. Lorente M. Palynology and palynofacies of the Upper Tertiary in Venezuela.
474 Dissertationes Botanicae (Gebrüder Borntraeger, Berlin). 1986; 99: 1-222.
475
- 476 22. Salard-Cheboldaeff M. Maestrichtian and Tertiary palynoflora of the coastal sedimentary
477 basin of Cameroon, pollen and spores. National Museum of Natural History. 1978; 215-260.
478
- 479 23. Sah SCD. Palynology of an Upper Neogene profile from Rusizi Valley (Burundi). Royal
480 Museum of Central Africa, Tervuren, Belgium. Annals Series 8 in Sciences.Géologique.
481 1967; 57: 173.
482
- 483 24. Bankole SI, Schrank E, Erdtmann BD. Palynology of the Paleogene Oshosun Formation
484 in the Dahomey Basin, southwestern Nigeria. Revista Española de Micropaleontología 2007;
485 39 (1-2): 29-44.
486

- 487
488 25. Salard-Cheboldaeff M. Paleopalynology of the coastal sedimentary basin of Cameroon in
489 relation to stratigraphy and palaeoecology. PhD thesis in Natural Sciences, Univ. Pierre and
490 Marie Curie, Paris VI. 1977; 262.
491
492 26. Salard-Cheboldaeff M. Maestrichtian and tertiary palynology of Cameroon. Qualitative
493 study and vertical distribution of the main species. Review of Palaeobotany and Palynology.
494 1979 ; 28 : 325-388.
495
496 27. Salard-Cheboldaeff M. Maestrichtian and Tertiary Palynology of Cameroon. Botanical
497 results. Review of Palaeobotany and Palynology. 1981; 32 : 401- 439.
498
499 28. Salard-Cheboldaeff M. Intertropical African palynostratigraphy from Cretaceous to
500 Late Quaternary times. Journal of African Earth Sciences. 1990; 11 (1-2) 1-24.
501
502 29. Kump B. Contribution to the knowledge of the microfossils of the middle and upper
503 Eocene. Palaeontographica, A. 1953 ; 103 : 377-406.
504
505 30. Cookson IC. The cenozoic occurrence of Acacia in Australia-Australian Journal of
506 Botany.1954 ; 2(1) : 52-59.
507
508 31. Rull V. Paleofloristic and palaeovegetational changes across the Paleocene/Eocene
509 boundary in northern South America. Review of Palaeobotany and Palynology. 1999; 107:
510 83-95.
511
512 32. Atta-Peters D, Salami MB. Late Cretaceous to Early Tertiary pollen grains from offshore
513 Tano Basin, Southwestern Ghana. Revista Española de Micropaleontología. 2004 b ; 36 (3)
514 :451- 465.
515
516 33. Cecile OM. Paleocene-Eocene boundary in the Douala basin: biostratigraphy and
517 palaeoenvironment reconstruction test. PhD Thesis, Faculty of Science, Paleobiogeology,
518 Paleobotany, Paleopalynology, University of Liège. 2013; 221.
519
520 34. Pardo-Trujillo A, Jaramillo CA, Oboh-Ikuenobe FE. Paleogene Palynostratigraphy of the
521 Eastern middle Magdalena valley, Colombia. Palynology. 2003; 27: 155-178.
522
523 35. El-Beialy SY, Ali SA. Dinoflagellates from the Miocene Rudeis and Kareem Formations
524 borehole GS-78-1, Gulf of Suez, Egypt. Journal of African Earth Sciences. 2002; 35: 235-
525 245.
526
527 36. Oloto IN. Maestrichtian dinoflagellate cyst assemblage from the Nkporo Shale on the
528 Benin flank of the Niger Delta. Rev. Palaeobot. Palyn. 1989; 57: 173-186.
529
530 37. Masare E, Rauscher J, Jejax M, Schuler M, Ferre B. Cretaceous-Paleocene Palynology
531 from the Côte d'Ivoire-Ghana Transform margin, sites 956, 960, 961 and 962. Proceedings of
532 the Ocean Drilling Program, Scientific Results. 1998; 159:253-276.
533
534 38. Rull VA. Quantitative palynological record from the Early Miocene of western Venezuela,
535 with emphasis on mangroves. Palynology. 2001; 25: 109-126
536

537 39. Samant B, Phadtare NR. Stratigraphic palynoflora of the early Eocene Rajpardi
538 lignite, Gujarat and the lower age limit of the tareswar formation of south cambay basin, India
539 palaeontographica abt. 1997; 1(6): 1-108.
540

541 40. Bankole IS. Palynology and stratigraphy of three deep wells in the Neogene Agbada
542 Formation, Niger Delta, Nigeria. Implications for petroleum exploration and
543 paleoecology. PhD Thesis, Technical University of Berlin. 2010; 190.
544

545 41. Mahmoud MS, Schrank E, Late Cretaceous spores, pollen and dinoflagellates from two
546 boreholes (Nuqra-1 and 3) in the Aswan area, southeastern Egypt. Journal of
547 Paleobiology. 2007; 26 (2): 593-613.
548

549 42. Jörg P, Schmiedl G. Early Oligocene dinoflagellate cysts from the Upper Rhine
550 Graben (SWGermany): Paleoenviromnement and Paleoclimatic implications. Marine
551 Micropaleontology. 2002; 45: 1-24.
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585