

2

3 **Theoretical Investigation into the Change in**

4 **the Number of Water Molecules in Solvent**

5 **Inaccessible Region of an Enzyme and**

6 **Enzyme-Substrate Complex**

7 **ABSTRACT**

8 **Background:** There may be dry enzymes, but water remains indispensable for the catalytic action of enzymes. There is not as much interest in how the presence of a drug such as aspirin and a psychoactive compound such as ethanol may affect the water-mediated role of the enzyme.

Objectives: The objectives of this research are: 1) To assess the changes in the number of water molecules interacting with the enzyme-substrate complex and the solvent inaccessible region of a protein, 2) to determine the free energy difference due to preferential solvation and hydration, and 3) to re-examine theoretical issues in literature and relate them to the interpretation of the results.

Methods: A major theoretical research and minor experimentation using Bernfeld method.

Results and discussion: The presence of ethanol/aspirin alone yielded only dehydration of the osmolyte inaccessible region and the enzyme substrate complex (*ES*). There was positive free energy difference ($\Delta\Delta G$) if the equilibrium constant for hydration change ($K_{eq}(1)$) > the equilibrium constant for folding-unfolding transition ($K_{eq}(3)$); it is negative where $K_{eq}(3)$ > $K_{eq}(1)$. Analysis of various models made them valuable for the interpretation

of result for feature application.

Conclusion

The change in the number of water molecules in an osmolyte inaccessible region of the enzyme and those interacting with the *ES* may be either positive or negative due respectively to sucrose and ethanol/aspirin. The spontaneity of two processes, hydration and folding-unfolding transition, the free energy difference, differs. The model for water stripping, preferential interaction concept, and the KBI for osmolation and hydration can guide the interpretation of the effects of any cosolute.

9 *Keywords: Porcine pancreatic alpha amylase; change in Gibbs free energy; change in the*
10 *number of water molecules; enzyme-substrate complex; osmolyte-inaccessible region of*
11 *enzyme; cosolutes; Kirkwood-Buff Integrals(KBI).*

12 13 **1. INTRODUCTION**

14 There are quite a lot of controversies surrounding the hydration of biomolecules. The
15 hydration of biomolecules is not in doubt but the effect of such hydration on internal
16 dynamics of the biomolecules is commonly of general interest to scientists [1]. However, this
17 does not exclude intermolecular dynamics needed for contact with each other or with other
18 solution components otherwise the needed contact for whatever transformation may not
19 occur; hence the proposition that enzymes most diffuse towards the substrate to align itself
20 with it to achieve a catalytic orientation [2]. This is notwithstanding current trend in the
21 development of immobilised enzymes, from amylase family. It must however, be made clear
22 that it is very impossible to digest polysaccharide without hydration of both substrate and
23 enzyme. A lot of interest has been shown in immobilised enzymes [3] for different reasons.
24 In those studies concern has been shown for the need for hydration, its purpose and effect
25 on the kinetic and thermodynamic stability of the enzyme. But there **seems** not to be much
26 interest on how the presence of drug such as aspirin and psychoactive compound such as
27 ethanol may affect the role of all kinds of hydration of the enzyme.

28 Some enzymes are known to possess conserved water molecules as part of the
29 structure of the enzyme's active site suggesting they play an important function in the active
30 site stability, flexibility, ligand coordination and residue positioning, hence their guided
31 evolutionary conservation [4]. Nuclear Magnetic resonance spectroscopy (NMR) analysis of
32 the hydration process indicates that the onset of catalytic activity is a direct consequence of
33 an increase in enzyme's (lysozyme's) conformational flexibility; it has been suggested that
34 this increased flexibility may be due, in part, to the reduced interaction of charged and / or
35 polar amino acid residues within the enzyme molecule caused by water's ability to effect
36 dielectric screening [5]. Yet there is objection against total reliance on flexibility for function
37 considering that **an** enzyme activity can occur at very low hydration levels, coupled with
38 reduction in protein's flexibility; this according to [6] calls for a rethink regarding the dynamic
39 requirement for **an** enzyme activity and stability.

40 In this research, the changes in the number of water molecules interacting with the
41 enzyme via its enzyme-substrate complex due to the presence of the additives that appears
42 not to feature very prominently in literature have become the concern of this research. In this
43 regard the view by Damien is relevant. Citing other workers, **Laage et al** [7] posits that water
44 strongly influences the structure and function of biomolecules within it. According to them [7]
45 the most relevant interactions are hydrogen bonds, a mainly local type of weak bonding
46 among water molecules which also exist between water and the polar or ionic groups of the
47 biomolecule; this is apart from other long-range Coulomb forces between formally charged
48 groups of the biomolecule. **Other forces are hydrophobic forces; the latter is relevant for the**
49 **aggregation of hydrophobic moieties; it can also enhance protein folding. It is known**
50 **elsewhere [8] that hydrogen bonding occurs in binary mixtures of organic solutes such as**
51 **ethanol and sucrose in this research. There could be altered dielectric property of the**
52 **primary solvent, water that can influence changes in the conformational stability of the**
53 **enzyme.** It is obvious that the relevance of water is accomplished through various forms of

54 interaction that cannot preclude interaction energy and solution structure in the presence of
55 additives in particular.

56 It should be realised that the presence of cosolvent or cosolute can alter the effect of
57 aqueous solvent on the structure and function of the enzyme. The thermodynamic and
58 activation parameters in terms of energy associated with *ES* may not remain the same in the
59 presence of cosolvents, otherwise called osmolytes. The description of the interaction
60 requires mathematical models that will be briefly addressed in theoretical section while a
61 detailed qualitative aspect of theory is to be addressed in the discussion section as part of
62 interpretational goal. The objectives of this research are 1) To assess the changes in the
63 number of water molecules interacting with enzyme-substrate complex and solvent
64 inaccessible region of a protein, 2) determine the free energy difference due to preferential
65 solvation and hydration and 3) reexamine theoretical issues in literature and relate same to
66 the interpretation of results.

67 2. THEORY

68
69 To begin with there is need to state that the major motivation of this section is the
70 need to establish a justifiable theoretical background that can enhance the quality and
71 perhaps, the validity and serve as a basis for the generation and possibly the interpretation
72 of result. This section has two parts viz: The review of the derived equation related to
73 difference in interaction free energy and the changes in the number of water molecules
74 interacting with the enzyme substrate complex (*[ES]*); the second part is concerned with the
75 change in the number of water molecules in osmolyte-inaccessible regions. The equation [9]
76 adopted as in previous publication [10] in the quantitative determination of pair-wise solute-
77 solute interaction parameter is as follows:

78

$$\ln[k_{(m_3)}/k_{(m_3=0)}] = \frac{2[g_{cx} - g_c^\#]m_3}{RTm_0^2} - N\phi M_1 m_3 \quad (1)$$

79 where $k_{(m_3)}$ is the (pseudo –) first – order rate constant in a reaction mixture containing co
 80 – solute whose concentration is m_3 and $K_{(m_3=0)}$ is the rate constant in the absence of the
 81 co – solute; R and T are the molar gas constant and thermodynamic temperature; m_0 is the
 82 (hypothetical) ideal reference state and it is equal to 1 mol/kg; $g_{cx} - g_c^\#$ is the difference in
 83 interaction Gibbs free energies between the co–solute c and the reactants β (and by
 84 extension substrate and a biochemical catalyst) on one hand and the activated complex $\#$
 85 on the other hand; M_1 , ϕ , N and m_3 are the molar mass of water, practical osmotic
 86 coefficient for the aqueous solution, the number of water molecules, and the molarity of the
 87 added cosolute respectively [9]. The equation seems to represents another way of
 88 expressing preferential interaction, a thermodynamic phenomenon applicable to multi-
 89 component solution. In the original equation by Buurma *et al* [9],

$$90 \quad RT \ln (k_{(m_3)}/k_{(m_3=0)}) = \Delta G(c)m_3 - N\phi RT M_1 m_3 \quad (2)$$

91 Where, R is the universal gas constant.

92 Thus,

$$93 \quad G(c)m_3 = \frac{2[g_{cx} - g_c^\#]m_3}{m_0^2} \quad (3)$$

94 The most important function of the enzyme is the lowering of activation energy and
 95 free energy of activation. Enzyme – substrate cannot proceed to product without initial
 96 activation which however occurs at a lower energy cost. Previous research attempted to
 97 apply this concept of pair-wise solution component interaction to biological system such as
 98 enzyme catalysed reaction in the presence of cosolute [10]. Here a more straight forward
 99 approach is further adopted to achieve similar result. If assay is at very high enzyme
 100 concentration, and if the substrate is not soluble, and if the raw insoluble starch was the
 101 substrate as in this research, a situation that satisfies the condition for reverse quasi steady
 102 state approximation (rQSSA) [11], then the equilibrium dissociation constant of the substrate

103 from the complex given as $K_s = k_{-1}/k_1$ where k_{-1} and k_1 are the rate constant for the
 104 dissociation of enzyme-substrate complex (ES) and the 2nd order rate constant for the
 105 formation of the ES respectively, should be the case. But the concept is also applicable to a
 106 situation where the substrate concentration is very high such that $\frac{[E_0]}{K_M + [S_0]} \ll 1$ ($[E_0]$, K_M ,

107 and $[S_0]$) are the total concentration of enzyme, the Michaelis-Menten constant, and
 108 concentration of the substrate) as to satisfy the condition for standard QSSA (sQSSA) [11].

109 This takes the form $K_M = \frac{k_{-1} + k_2}{k_1}$ where k_2 is the rate constant for product formation and
 110 release. However, the key issue is that the rate constants for the dissociation of ES can be
 111 expressed respectively as

$$112 \quad k_{-1} = K_s k_1 \quad (4)$$

$$113 \quad k_{-1} = K_M k_1 - k_2 \quad (5)$$

114
 115
 116 In this research Eq. (4) unlike Eq. (5) does not present any issue because $\ln[k_{-1(m_3)}/$

$$117 \quad k_1 = - \frac{\partial \ln([S](t)/[S]_0)}{\partial t [E_0]} \quad (6)$$

118 Where the concentration of enzyme ($[E_0]$) assayed is held constant or fixed while the
 119 concentration of the substrate in time $t = 0$ is $[S_0]$, and $[S](t)$ is the concentration of the
 120 substrate in time, t .

$$121 \quad \ln(k_{s(m_3=0)}/(k_{s(m_3)})) = (\Delta G m_3 - \Delta G_{-1} m_3)/RT + M m_3 (\varphi_{-1} n_{-1} -$$

$$122 \quad n\varphi) \quad (7)$$

$$123 \quad = \frac{\Delta \Delta G m_3}{RT} + M m_3 \varphi \Delta n \quad (8)$$

124 It is not in doubt that Michaelis-Menten (MM) constant is a sum of equilibrium
 125 constants. This is to say that it is given as $K_M = \frac{k_{-1}}{k_1} + \frac{k_2}{k_1}$; this implies that

126



127 The variable, S_{FR} as explained elsewhere [12], is the fragment of the polysaccharide left
128 after a given catalytic cycle; no single polysaccharide is totally hydrolysed by an appropriate
129 hydrolase.

130 The change in the number of water molecules in osmolyte-inaccessible regions,
131 ΔN_w , is given by the slope of line relating $\ln(K_{eq})$ and the osmolyte concentration as follows
132 [13].

$$\ln K_{eq} = - \frac{\Delta N_w [\text{Osmolal}]}{55.56} \quad (9b)$$

134 Leading to Eq. (9b) from the perspective of osmolyte-inaccessible regions is simply
135 reaffirmation of the principle of preferential exclusion anchored on Kirkwood-Buff theory
136 [KBT] of solution structure that has been popularised in recent papers [14, 15]. The
137 theoretical interest arises from what appears to be a common ground for Eq. (2), Eq. (8), and
138 Eq. (9) in that the number of water molecules for different purposes can be calculated from
139 all equations, one from the slope (Eq. (9)) and the other from intercept of either Eq. (2) or
140 Eq. (8). Equation (8) which arises from theoretical exposition of Buurma *et al* [8] represented
141 the first time observation was made of the appearance of variable - concentration of cosolute
142 - in two places as an independent variable in an equation.

143 Soluble polar organic substances called osmolytes may be excluded from the
144 protein surface domain on account of their inability to penetrate protein's inner region. This
145 issue is important in the light of the fact that solvent accessibility change plays a critical role
146 in protein misfolding and aggregation, the culprit for several neurodegenerative diseases,
147 including amyotrophic lateral sclerosis (ALS) [16]. Furthermore, solvent accessibility may be
148 part of the structural environment of amino acids in the protein that might influence the
149 function-structural (mechanical) and catalytic in nature-of any of such amino acids [17].
150 Directly or indirectly, this important issues may have prompted research in this issue of

151 inaccessible core of the protein as exemplified in very recent research outcome which
152 showed that the solvent-inaccessible cores of the three classes of proteins are equally
153 densely packed [18]; this constitute steric hindrance to the penetration of relatively large
154 organic osmolytes. This may have promoted excess flexibility that caused increasing velocity
155 of hydrolysis with higher concentration of ethanol. One must not fail to point out that
156 osmophobic concept [19] has been advanced as basis for the action that compels a protein
157 to fold due to exclusion of such osmolyte which exist in nature from protein back bone.

158 On account of the issues raised in the text, there is need to recall that preferential
159 osmolation, either negative or positive, can yield (re) folding and unfolding as the case may
160 be leading to equilibrium state if a two-state model is assumed. Hence, the equilibrium
161 constants (K_{eq}) defined mathematically and given below are of paramount relevance to a
162 system in near dynamic equilibrium.

163
$$K_{eq(3)} = \frac{[U]}{[N]} \quad (10)$$

164 Equation (10) is adapted from the work by Pace [20] which the author restate as $K_{eq(3)} =$
165 $U/(1-U)$ where in this case U and $1-U$ denotes fraction of unfolded protein molecular
166 population and fraction of folded protein respectively. Equation (10) or its alternative is
167 expressible in two ways in accordance to whether or not the observed catalytic activity of the
168 enzyme in the presence of osmolyte is greater than the same activity in the absence of the
169 osmolyte. The equations which are applicable to the effects arising from the presence of an
170 osmolyte are to be stated in method subsection. But there is also preferential hydration and
171 dehydration due to preferential exclusion and binding of appropriate osmolyte respectively.
172 This creates directional aqueous molecular motion to and from the protein's surface domain
173 leading to an equilibrium system described by the second equation of equilibrium constant
174 given elsewhere [21] as

175
$$K_{eq(1)} = \exp\left(-\frac{\ln a_1 c_1}{c_3} \Delta\Gamma_{23}\right) \quad (11)$$

176 Where $\Delta\Gamma_{23}$ is the change in preferential interaction by either binding or relative exclusion of
177 an osmolyte; C_1 and C_3 are molar concentrations of water and osmolyte respectively; a_1 is
178 the activity of water in aqueous solution of osmolyte. Meanwhile the equation of preferential
179 interaction [15] is given as

$$180 \quad \Delta\Gamma_{23} = \frac{\ln K_{eq(3)}}{\ln a_3} \quad (12)$$

181 The emergence of Eq. (12), as in previous publication (15), is as a result of the proposition
182 that a parameter cannot be a devise-based measurable quantity (without definite or finite
183 magnitude) as well as a constant quantity. A measurable quantity is an extensive
184 thermodynamic parameter and, if a given ratio is always constant regardless of the
185 magnitudes of the compared parameters, it becomes an intensive thermodynamic quantity.
186 The report at the web site, en.Wikipedia.org (<https://www.en.Wikipedia.org>) shows that
187 Richard C. Tolman was the author who first introduced the concept of extensive and
188 intensive quantities.

189 **3. MATERIALS AND METHODS**

190 **3.1 Material**

191 As stated elsewhere[10], the materials are The chemicals used were: The chemicals
192 used were: Sucrose (St Lious France); raw (native) potato starch (Sigma Chemicals Co, USA);
193 ethanol, hydrochloric acid and sodium chloride (BDH Chemical Ltd, Poole England); 3,5-
194 dinitrosalicylic acid (DNA) (Lab Tech Chemicals, India); Tris (Kiran Light Laboratories, USA);
195 porcine pancreatic alpha amylase (EC 3.2.1.1) (Sigma, Adrich, USA); all other chemicals were of
196 analytical grade and solutions were made in distilled water. Aspirin was purchased from CP
197 Pharmaceuticals Ltd, Ash road North, Wrexham, LL 13 9UF, and U.K

198 **3.2 Equipment**

199 *pH* meter (tester) from Hanna Instruments, Mauritius; electronic weighing machine
200 from Wensar Weighing Scale Ltd, Chennai; Centrifuge, 300D model from China; 721/722
201 visible spectrophotometer from Spectrum Instruments Co Ltd, China.

202 3.3 Method

203 As stated elsewhere [21], 0.01 g of PPA was dissolved in 20 mL of distilled water to give
204 500 µg/L while potato starch was prepared by dissolving 1 g in tris-HCl buffer (aq.) buffer (90
205 mL), 5 mL, 6% (W/W), NaCl (aq.) and 5 mL distilled water to give 1 g/100 mL. Approximate
206 dilutions were carried out for the determination of K_M and V_{max} at 37°C and pH of 7.4 by
207 Lineweaver-Burk plot [22]. As in previous investigation [14] a method adopted for the
208 determination of velocity (v) where $C_3 \rightarrow 0$ is a re-modification of the same equation found in
209 literature [23]. It may appear theoretical but that is the essence of this research, a combination of
210 a major theory and minor experimentation. Increasing v with increasing C_3 , demands a plot of
211 $\log v$ versus C_3 which gives an intercept, being an extrapolated velocity ($v_{C_3 \rightarrow 0}$) of hydrolysis
212 as $C_3 \rightarrow 0$.

$$213 \quad \log v = \log v_{C_3 \rightarrow \infty} - \beta[C_3]. \quad (13)$$

214 Equation (13) is for the increasing v , while for the decreasing case, it is given as

$$215 \quad \log v = \log v_{C_3 \rightarrow \infty} - \beta/[C_3]. \quad (14)$$

216 However, in order to obviate the effect of outliers, linear regression line was allowed to link
217 the lowest point with the highest point for the determination of the minimum v as $C_3 \rightarrow 0$.
218 Assay for the generation of velocities of the hydrolysis of starch is according to Bernfeld
219 method [24].

220 The equation (Eq. (15)) below is adopted for the purpose of comparing the transition
221 state energies of two different equilibrium systems dehydration/hydration and
222 osmolation/exclusion equilibria due to the presence of osmolytes or cosolvents; it is
223 therefore, restated as

$$224 \quad \Delta\Delta G_{ES}^\# = -RT \ln \frac{K_{eq(1)}}{K_{eq(3)}} \quad (15)$$

225 The equilibrium constant $K_{eq(1)}$ is determined by substituting relevant parameters into Eq.
226 (11); $K_{eq(2)}$ is determined by exploring either Eq. (16) or Eq. (17) below.

227
$$K_{eq(3)} = \frac{V_N - V_{OBS}}{V_{OBS} - V_{MIN}} \quad (16)$$

228 Where $V_N > V_{OBS} > V_{MIN}$ and the subscripts, N, OBS, MIN are respectively, catalytic
229 activity of native enzyme, observed activity of treated enzyme and minimum activity of
230 treated enzyme.

231
$$K_{eq(3)} = \frac{V_N - V_{MIN}}{V_{MAX} - V_N} \quad (17)$$

232 The issues that led to the emergence of Eq. (16) and Eq. (17) were addressed in part
233 elsewhere [25]. Further details are currently in manuscript under preparation. Microsoft Excel
234 (2007) was used to plot the dependent variable versus independent variable.


235 3.4 STATISTICAL ANALYSIS

236 The velocities of hydrolysis were determined in triplicates. The mean values were
237 used to determine the equilibrium constant.


238 4. RESULTS AND DISCUSSION

239 Before analysis and discussion of results, there is a need to review Eq. (2) and Eq.
240 (8). A careful examination of the equations reveals that the slope and intercept may not be
241 positive or negative; the values depend on the magnitude of the ratio given as $K_{(m_3)}/$
242 $K_{(m_3=0)}$. If $K_{(m_3=0)} < K_{(m_3)}$, with increasing values of $K_{(m_3)}$, the value of the ratio should
243 be decreasing from smaller negative values to larger negative values, such that a plot
244 of $\ln(K_{(m_3)}/K_{(m_3=0)})$ versus m_3 should give a negative slope and definitely a negative
245 intercept. It is also probable that $\frac{K_{(m_3)}}{K_{(m_3=0)}} > 1$ such that any plot may give a positive slope or
246 correlation and either a positive intercept or intercept which is negative but much smaller in
247 magnitude. Therefore, characteristics such as the magnitudes and signs of the slope and

248 intercept of a straight line from the plot express the type of change in the number of water
249 molecules, which is either net hydration (positive) or net dehydration (negative). These
250 account for the shapes of various curves shown in Figure 1 through 6. To show the direction
251 of shift in the hydration process in the equilibrium, $E + S \rightleftharpoons ES$, a plot of Δn versus m_3
252 was carried out. The result (Fig. 1) shows that there was a decreasing trend in the change in
253 the number of water molecules interacting with the ES . The observed trend is due to the
254 effect of aspirin. The decreasing trend along the positive axis suggests that there was a
255 decrease in hydration due to the effect of aspirin alone (Fig. 1). It is a loss-dehydration- the
256 magnitude of which showed a decreasing trend, progressing towards net hydration (Fig. 2)
257 due to the effect of the second cosolute, sucrose. In this case, the variation of the change in
258 the number of water molecules with the molar concentrations of sucrose showed mixed
259 trend. With a lower concentration of the cosolute (1.55 mmol/L and 0.73 mmol/L) – aspirin –
260 there was an increasing trend unlike with higher concentration of the same cosolute, due
261 perhaps, to the effect of the 2nd cosolute (sucrose) in the reaction mixture (Fig. 3).
262




263 **Fig. 1. Variation of the change in the number of water molecules interacting with the**
264 **enzyme- substrate complex with different concentration of aspirin. n is the number of**
265 **water molecules.**
266
267



268
269
270
271
272
273
274
275

Fig. 2. Variation of the change in the number of water molecules interacting with enzyme substrate complex with different [Aspirin] at different fixed concentration of sucrose whose concentration ranges between 3.60-57.53 mmol/L. n and SUC denote the number of water molecules and sucrose.




276
277
278
279
280
281
282
283

Fig. 3. Variation of the change in the number of water molecules interacting with the enzyme-substrate complex with different concentration of sucrose (SUC) at different fixed concentration of aspirin. $n(1)$ and ASP denote the number of water molecules and aspirin respectively.


284 Like the trend observed with the effect of aspirin there is also a decreasing trend in
285 the positive values of Δn with increasing concentration of ethanol (Fig.4). Variation with
286 different concentrations of ethanol exhibited similar trend observed for the variation of Δn
287 with molar concentration of aspirin (Fig.5). With a mixture of ethanol and sucrose, there was,
288 as was the case with the effect of a mixture of aspirin and sucrose, a mixed trend in the
289 variation of Δn with molar concentrations of sucrose (Fig. 6). All these observation

289 notwithstanding, it is rather difficult to suggest why such observations cannot be mere
 290 coincidence taking into account the effect of high degree of improvisation in the conduct of
 291 the experiment. It is not an overemphasis to opine that ethanol is totally different from
 292 aspirin; while the former is essentially psychoactive, the latter is a well known non-steroidal
 293 anti-inflammatory drug [26, 27], and both have adverse effects on intestinal brush border
 294 membranes that could compromise the biological function of brush border membrane
 295 enzymes and transporters respectively.




296
 297
 298
 299
 300
 301
 302
 303

Fig. 4. Variation of the change in the number of water molecules interacting with enzyme substrate complex with different concentration of ethanol. $n(1)$ and ETH denote the number of water molecules and ethanol respectively.



304
 305

306 **Fig. 5. Variation of the change in the number of water molecules interacting with**
 307 **enzyme- substrate complex with different [Ethanol] at different fixed concentration of**
 308 **sucrose. SUC and $n(1)$ denote sucrose and number of water molecules respectively.**
 309



310
 311
 312 **Fig. 6. Variation of the change in the number of water molecules interacting with**
 313 **enzyme-substrate complex with different [Sucrose] at different fixed concentration of**
 314 **ethanol. ETH and $n(1)$ denote ethanol and number of water molecules respectively.**
 315

316
 317 **Table 1. Change in the number of water molecules in osmolyte-inaccessible regions**
 318 **due to ethanol-sucrose mixture**

319	<hr/>					
320	[Ethanol]/(mol/L)	1.25	3.23	5.28		
321	ΔN_W	-601.67	615.050	3809.19		
322	r^2	0.82	0.53	0.77		
323	<hr/>					
324	As function of ethanol concentration					
325	<hr/>					
326	[Sucrose]/(mmol/L)	0.00	3.57	7.14	14.29	28.57 57.14
327	ΔN_W	-34.56	-50.06	-69.26	-36.00	-9.06 -21.11
328	r^2	0.86	1 (2dpts)	0.95	0.94	0.95 0.60
329	<hr/>					

330 *Changes in the number of water molecules (ΔN_W) are calculated as the product of slope*
 331 *and 55.56; the slope may be obtained from the plot of $\ln K_{eq}$ versus [cosolute] at a fixed*
 332 *concentration of the 2nd cosolute; dpts mean data points.*
 333

334 The change in the number of water molecules (ΔN_W) on osmolyte inaccessible
 335 region as a function of sucrose concentration is similar to the exclusion of aqueous solvent
 336 or dehydration with lower concentration of ethanol unlike with higher concentration ethanol in
 337 which there was hydration (Table 1). But as function of the concentration of ethanol, there
 338 was irregular trend couple with a case of dehydration similar to result obtained in only
 339 ethanol treated enzyme (sucrose concentration = 0) (Table 1). This is not unexpected
 340 considering ethanol as a fluidising and water-stripping agent.

341 **Table 2. Change in the number of water molecules in osmolyte-inaccessible regions**
 342 **due to aspirin-sucrose mixture**

343						
344						
345	[Aspirin]/(mol/L)	0.76	3.05	6.10		
346	ΔN_W	2042.86	4380.01	6261.61		
347	r^2	0.98	0.97	1 (2dpts)		
348	As function of aspirin concentration					
349						
350	[Sucrose]/(mmol/L)	0.00	7.19	14.38	28.76	57.75
351	$\Delta N_W/\exp(3)$	-40.63	-80.84	-83.90	-52.85	-44.93
352						
353	r^2	0.87	1(2dpts)	1(2dpts)	0.99	0.98

354 *Changes in the number of water molecules (ΔN_W) are calculated as the product of slope*
 355 *and 55.56; the slope may be obtained from the plot of $\ln K_{eq}$ versus [cosolute] at a fixed*
 356 *concentration of the 2nd cosolute; dpts mean data points.*

357
 358 The change in the number of water molecules as a function of sucrose concentration
 359 showed increasing trend unlike such change as a function of aspirin concentration in which
 360 there was irregular trend and negative in sign as to imply dehydration (Table 2) similar to the
 361 result obtained due to the effect of aspirin alone. This may imply that aspirin like ethanol has
 362 water-stripping properties.

363 The difference in free energies between two thermodynamic processes
 364 dehydration/hydration and osmolation/exclusion arising from the effect of cosolutes and
 365 water are recorded in Tables (3a), (3b), (4a), and (4b). There is need to state that the data

366 generated is not an outcome of high precision measurement as a result of improvisation.
 367 This leaves room for further research using state-of-the-art facilities while the current data
 368 remains purely illustrative of the fact and principle enunciated in this research. Usually, a
 369 spontaneous process is one in which the free energy is relatively large and negative in sign.
 370 The effect of ethanol and aspirin separately alone, yielded a mixed result of negative and
 371 positive free energies as shown in Tables (3a) and (4a) respectively. The negative difference
 372 in free energy occurred with higher concentration of the cosolutes. With a mixture of ethanol
 373 and sucrose (Table 3b) and a mixture of aspirin and sucrose (Table 4b), the negative values
 374 occurred with higher concentration of ethanol and aspirin. What one can deduce is that
 375 positive $\Delta\Delta G$ occurs if $K_{eq}(1) > K_{eq}(3)$ and as such (de) hydration is more spontaneous.
 376 On the other hand if $K_{eq}(1) < K_{eq}(3)$, a negative $\Delta\Delta G$ may be given with the result that,
 377 osmolation/exclusion is more spontaneous.

378 **Table 3a. Difference in free energies between dehydration/hydration and**
 379 **osmolation/exclusion with only ethanol.**

380						
381						
382	[Ethanol]/mol/L	1.247	2.398	3.228	4.311	5.279
383						
384	$\Delta\Delta G$ /kJ/mol	18.680	0.960	-0.380	-0.360	-0.280

385 $\Delta\Delta G$ is the difference in free energy.

386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400

401 **Table 3b. Difference in free energies between (de)hydration and (de) osmolation with a**
 402 **mixture of ethanol and sucrose**

	[Sucrose]/mmol/L				
	3.57	7.14	14.29	28.57	57.14
[Ethanol]/mol/L	$\Delta\Delta G$ /kJ/mol				
1.247	3.53	4.22	2.43	2.42	1.81
3.228	-1.92	-0.49	0.12	0.28	0.64
5.279	-	-11.35	-4.84	-0.95	4.44

413 $\Delta\Delta G$ is the difference in free energy.

415 **Table 4a. Difference in free energies between dehydration/dehydration and**
 416 **osmolation/exclusion with only aspirin.**

[Aspirin]/mmol/L	0.76	1.53	3.05	4.58	6.10
$\Delta\Delta G$ /kJ/mol	2.63	2.49	1.84	- 2.85	- 9.55

422 $\Delta\Delta G$ is the difference in free energy.

424 **Table 4b. Difference in free energies between dehydration/hydration and**
 425 **osmolation/exclusion with a mixture of aspirin and sucrose**

	[Sucrose]/mmol/L			
	7.19	14.38	28.57	57.14
	$\Delta\Delta G$			
[Aspirin]/mmol/L	0.76	6.03	7.39	9.11
3.05	- 4.29	- 3.61	0.74	1.49
6.10	-	-	- 7.76	- 5.50

437 $\Delta\Delta G$ is the difference in free energy.

440 The results obtained so far are significant because of the biological role of water,
 441 and, there are a lot of theoretical basis for them. Such theoretical foundation or basis
 442 broadens the scope for further research. Beginning from what is known is the fact that
 443 proteins are strongly hydrated in aqueous medium. The density of water molecules close to

444 the protein surface due to effect of polar and non-polar groups is as high as 1.25 g/mL within
445 3-4.25 Å of protein surface, mainly as result of large number of water molecules that are
446 3.75 Å from non-polar atoms; within 2.5 Å of the protein surface there is a small increase in
447 density of water molecules due to electrostriction around the polar groups; but 3-4.5 Å from
448 the surface, there is a slight decrease in density [28]. Water molecules are clustered
449 perpendicular to the protein surface while in the parallel direction to the protein surface the
450 water molecules are more disperse [28]. This means that given suitable pH, an enzyme
451 exhibits a level of hydration needed for function. For a particular group, the fraction of time
452 when a water protein hydrogen bond is formed otherwise called hydrogen bond probability
453 (P_{hb}) is strongly dependent on protein accessible surface area (ASA). The lower the latter,
454 the higher the entropic barrier (cost) that should be paid to significantly reduce the flux of
455 water molecules on the protein surface hydration site where H-bond is expected [29]. In the
456 same vein, Ooi & Oobatake [30] also posited that each atomic group interacts with water in
457 proportion to its water-ASA. The effect of the presence of chaotropes is of major concern as
458 it has been observed that more polar organic solvents (tetrahydrofuran and acetonitrile)
459 replace mobile and weakly bound water molecules in the active site and leave primarily the
460 tightly bound water in that region [31].

461 In this research, aspirin and ethanol decreased the velocities of amylolysis of raw
462 potato starch. These velocities under the influence of aspirin and ethanol are respectively
463 21-74 U/mL and 38 – 61 U/mL, all being < the velocity of amylolysis (97 U/mL) by the
464 untreated enzyme. This means that the entropic cost of fixing water of hydration increased
465 as a consequence. This can be elucidated via the equation made popular by Petukhov [29].

466 The equation is

467
$$\Delta G_{hb} = -RT \ln \frac{P_{hb}}{1 - P_{hb}} \quad (18)$$

468 Equation (18) suggests that as $P_{hb} \rightarrow 1$ the value of the free energy of hydration tends
469 toward higher negative magnitude as an expression of feasibility or spontaneity of hydration.

470

$$P_{hb} = \frac{\exp\left(-\frac{\Delta G_{hb}}{RT}\right)}{1 + \exp\left(-\frac{\Delta G_{hb}}{RT}\right)} \quad (19)$$

471 Hence if $P_{hb} \rightarrow 1$ (or if it's equivalent 100 $P_{hb} \rightarrow 100\%$), the entropic cost should tend to
472 minimum. It is important to point out that the value of P_{hb} may be a function of the fraction of
473 water population that can form H-bond with 4 water molecules and 3 water molecules both of
474 which are a function of the prevailing temperature in a equation given according to Petukhov
475 [29] as

476

$$P_{hb} = 100(4X_1 + 3X_2)/4 \quad (20)$$

477 Where X_1 and X_2 are respectively the fraction of water that can H-bond with 4 and 3
478 molecules of water. As this research shows, the presence of sucrose seems to have
479 opposed the effect of aspirin and ethanol. As such it is expected that P_{hb} may have
480 increased as a result of the effect of sucrose.

481 The role of water, or the effect of hydration, has its theoretical foundation that
482 enhances the interpretation of results. It has been reported that “the hydration environment
483 of a protein significantly affects its dynamics. This is why changes in the number of water in
484 cosolvent inaccessible site of the protein have become very important because such can
485 affect enzyme function. A positive change indicates that there may have been hydration and
486 negative change means the opposite. Such changes may not have been possible if there
487 was no initial hydration and preferential interaction with molecules of water. Although the
488 method adopted by Mitchell and Litman [13] and Buurma [9] are different they have a
489 common ground for addressing the issue of hydration changes. This is the case because
490 osmolyte inaccessible region of the protein may accommodate the active site. The active site
491 is either located within the protein's inner part or at locations close to the surface domain of
492 the enzyme. Hence changes in the number of water molecules in an osmolyte inaccessible
493 region cannot totally exclude the active site. This is the case, if one recalls that ES complex

494 is the result of complex formation between substrate and active site of the enzyme.
495 Therefore, there could be changes in the number of water molecules interacting with *ES*.

496 In literature, following the application of osmotic stress, is the observation that
497 protein-DNA complex can be hydrated with measurable volume changes. The sign of the
498 changes of the number of water molecules interacting with the protein and *ES* as well as
499 osmolyte inaccessible region indicates the occurrence of either hydration or dehydration. As
500 in this research such change occurs when a cosolute is introduced into the medium. It has
501 been observed that the catalytic activity of lyophilised oxidative enzyme was lower when
502 directly suspended in organic solvents containing little water than when they are introduced
503 into the same largely nonaqueous media by first dissolving them in water and then diluting
504 with anhydrous solvents [32]. Despite the need for water for maximum catalytic activity of
505 enzymes, an obvious paradox exists to the effect that, some enzymes (subtilisin and alpha-
506 chymotrypsin) showed a 100 billion-fold enhancement in nonpolar solvent like octane with
507 just an amount of water much less than needed to form a monolayer [33]. This is attributed
508 to an increase in the kinetic barrier (activation energy) needed to be overcome in order to
509 transform from native to unfolded conformation [33]. This should not be surprising because
510 unlike polar solvents, e.g. ethanol and polar solute, e.g. aspirin in this research, that have
511 water-stripping power, octane does not being hydrophobic.

512 It is quite obvious that infinitesimal amount of water is needed to trigger catalytic
513 action as to imply that water may be described as a *prima facie* example of an inorganic
514 catalyst. Highly structured water molecules are needed around the protein surface as part of
515 efficient chemistry of the protein by which they promote the protein's three dimensional (3-D)
516 structures [34]. According to Csermely [35], water molecules within the region of perturbative
517 influence of the enzyme provides the environment by which fluctuating changes in hydrogen
518 bond could occur as a necessary requirement for protein flexibility, structural
519 rearrangements leading to conformational transitions needed for catalytic functions. This
520 very much in agreement with the observed hydration induced conformation and dynamic

521 changes which are completed just before the onset of enzyme biological function [36]. It
522 goes to confirm that an increased rigidity in the protein at low hydration can be reversed
523 when water is added to the dry enzyme leading to a “loosening up” or increase in flexibility
524 [36]. Protein flexibility means inter-domain and catalytic site mobility made possible waters of
525 hydration. The deduction one can make is that polar solvent like ethanol as in this research
526 displaces the weakly bound structural water molecules and preferential water of hydration
527 leading to alteration and distortion in the catalytic conformational transition needed for
528 function that culminate in lower velocity of amylolysis.


529 However, the hydration effects are strongly dependent on both temperature and
530 hydration. At cryogenic temperatures, hydration stiffens protein structure because of the
531 hydrogen-bond interaction, whereas at physiological temperatures, hydration softens the
532 structure through the activation of anharmonic motion”[1]. The hydration water dynamics and
533 their dynamical coupling with the protein are presumed to be essential for protein dynamics
534 and biological function [1]. The protein dynamics in question is actually intra-molecular
535 dynamics needed for conformational flexibility for function. According to Chaplin
536 (www1.lsbu.ac.uk), proteins are characterized by conformational flexibility, which entails a
537 wide range of hydration states, in a state of dynamic equilibrium, facilitated by the ease of
538 hydration. The ease of hydration is dependent on the activity of the surrounding water
539 molecules. The enzymatic function of the enzyme is dependent on the position of the
540 equilibrium, $ES \rightleftharpoons CS$ (where ES and CS mean the expanded state and compact state of
541 water respectively) around the protein; the es is also called the Ih-type with lower density-
542 the low density water (LDW) while CS is called II-type with higher density-the high density
543 water (HDW). The LDW and HDW are respectively more ordered and less ordered. Thus an
544 intermediate mixture of nonionic kosmotropes and nonionic chaotropes such as sucrose and
545 aspirin/ethanol respectively as in this research can enhance biological activity of the

546 enzyme: It is neither an excessive rigidity nor an over flexibility of the structure of the protein
547 that can enhance the function of an enzyme.


548 The effect of a mixture of protecting and destabilising cosolute as observed in this
549 research has its theoretical foundation. Beginning with aspirin and ethanol, the theory is
550 described as preferential interaction by osmolation or by binding and by exclusion. The
551 former, according to Timasheff [37], leads to water stripping according to the equation:



553 Where, L is the ligand otherwise called cosolute. Citing his previous paper, Timasheff [36]
554 posits that “the reference state is the protein dissolved in water, in which it is fully hydrated.
555 Therefore, in a binary solvent, the binding of the nonaqueous solvent component to any
556 locus must displace water, *i.e.*, binding is an exchange reaction” [37]. Unfortunately there is
557 no equation for exclusion of ligand as at this moment. Nonetheless, the following equation
558 may serve this purpose.



560 Equation (22) (which reflects only the expulsion of water) symbolically shows that when an
561 aqueous solution of a protein is introduced into a stabilising cosolute, the macromolecule will
562 not be totally free from the molecules of the cosolute. Hence β is the small portion of L
563 (stabilising osmolyte in this case) that binds while $L-\beta$ is the vast amount of L that is
564 excluded given that β is equal to the amount of water displaced. This could have given rise
565 to



567 However, one must not overlook the effect of osmotic stress that might be created when any
568 cosolute is excluded leaving the surrounding of the protein more concentrated as to create
569 concentration gradient. This can compel loosely bound water molecules to depart the protein

570 into the bulk; this may also be interpreted as a translational entropy gain of the aqueous
571 solvent that drives re-folding [38]. This is in agreement with the view that osmotic pressure
572 controls the activity of water in an aqueous compartment inaccessible to neutral solutes
573 (osmolytes). The osmotic stress created then induces the release of bound water from
574 macromolecules into bulk solvent. Macromolecular conformations are thus shifted toward the
575 state with the smallest volume, which is the state with the least amount of bound water [39,
576 40]. The folded state promoted by stabilisers such as sucrose in this research has smaller
577 volume. This is another evidence of the importance of hydration, be it water of hydration or
578 water of preferential hydration. Osmotic stress will always occur when there is the presence
579 of a stabiliser in particular.

580 Further evidence is the observation about individual, internal water molecules that
581 may be reactants in a catalyzed reaction and/or may be integral parts of a protein structure,
582 providing stereospecific interactions; the correlation between hydration and increased
583 activity means that it is likely that the observed hydration-induced dynamical changes may
584 facilitate activity [6]. However, the presence of amylolytic activities, in the absence of intra
585 molecular motion, indicates that the motions are not an absolute requirement; this seems to
586 imply that if enzyme activity can occur at very low hydration levels, and if at these levels
587 protein flexibility is reduced, then the dynamic requirement for enzyme activity and stability
588 may be questionable [6]. Here one must strongly disagree on account of this research
589 outcome and on the basis of common sense. The lower amylolytic activity of ethanol/aspirin
590 treated-enzyme suggest that the structural water within the protein core and active site in
591 particular may be weakly intact, but other catalytically supportive mobile water molecules
592 may have been stripped off. In other words the ubiquitous surface hydration shell has
593 influence on protein dynamics and function such that if adversely affected by the chaotropes
594 [6], a reduction in amylolytic activity becomes inevitable. This clearly explains the decrease
595 in the velocity of amylolysis for ethanol/aspirin treated enzyme. Scholars who are only
596 interested in balanced diet may consume the usual diet thrice without water for two days but

597 with implication of being inflicted with indigestion and constipation despite the fact that ab
598 initio, the gastrointestinal tract is not dry. No matter the length of time no reaction can be
599 noticed in a dry mixture of enzyme and its substrate.

600 This can be accounted for in terms of Kirkwood-Buff theory (KBT) of solution
601 structure which states that the average structure of all solutions [41] is given by radial
602 distribution function ($g_{\alpha\beta}(r)$) between two species, namely, α and β . The term, radial
603 distribution function, is a measure of the deviation from the random distribution of particles of
604 type β from a central particle of type α as a function of the distance (r) from the central
605 particle [41]. A positive or negative deviation of $g_{\alpha\beta}(r)$ (also known as pair correlation
606 function) from unity, at a certain distance corresponds to excess or deficit of β at the
607 indicated distance from the particle designated as α . The issue remains the combined effect
608 of aspirin/ethanol and sucrose.

609 According to Bolen and Baskakov [42], the osmophobic effect of osmolyte is a vital
610 property that is beneficial to life, being the capacity for an unfavourable interaction between
611 the osmolyte/cosolute and peptide backbone. In the same vein, Baskakov and Bolen [43]
612 opined that the osmophobic effect of stabilisers on the peptide back bone made the unfolded
613 state of protein in osmolyte solution very unfavourable relative to the folded state; therefore,
614 it was the strongly destabilising effect of stabilisers such as sucrose on the unfolded state as
615 in this research, that forces the enzyme to refold. From the perspective of thermodynamic
616 stability, Bolen and Baskakov [42], see solvophobic action which Schellman [44] and Rösger
617 *et al* [41] called excluded volume action, as a factor which raised the free energy of the
618 denatured state, shifting the equilibrium in favour of the native state. In this research sucrose
619 is a well known stabiliser which acts by preferential exclusion. On account of this sucrose
620 was able to enhance the amyolytic velocities of sucrose treated-enzyme in a reaction
621 mixture containing aspirin (3.052 mmol/L) and ethanol (3.228 mol/L): The velocities ranges
622 from 132-140 and 116-136 U/mL respectively. These values were higher than values

623 obtained for the untreated native enzyme (97 U/mL), only ethanol-treated (102 U/mL) and
624 only aspirin treated (69 U/mL) enzyme.

625 The phenomena of solvophobic and solvophilic effect are the root cause of the
626 change of biological function-either an increase or decrease in the velocity of catalytic action
627 for instance as noted in this research. Osmolytes (as cosolvents/cosolute) may be
628 solvophobic or solvophilic (preferential exclusion or osmolation *i.e.* preferential interaction by
629 binding) which causes respectively refolding and unfolding; this presupposes changes in the
630 volume or 3-D structure of the macromolecule. This needs interpretational analysis based on
631 what Rösgen *et al* called inverse KBT. It is usually a context between solvation and hydration
632 change expressed via the KB integrals (KBIs). From the point of view of preferential
633 hydration integral, the following equation is inevitable. The partial molar volume of the
634 protein is in contention. Thus, the change in G_{pw} due to folding to unfolding transition is
635 given as

$$636 \quad \Delta_N^D(G_{pw}) = -\Delta_N^D V_{prot} + \frac{m\phi_{os}}{RT} \quad (23)$$

637 Where $\Delta_N^D V_{prot}$ is the partial molar volume of the protein, ϕ_{os} is the volume fraction of the
638 osmolyte, G_{pw} is the KBI for hydration, m is the short form of m -value, the capacity of
639 osmolyte to cause conformational change and Δ_N^D means folded to unfolded transition. For
640 the ideal case as may be applicable to dilute solution of sucrose, positive m -value for the
641 protecting osmolyte, should be such that $\Delta_N^D(G_{pw})$ may be positive in sign as to imply an
642 increase in the number of water molecules around the protein. This view is premised on the
643 fact that the (re)folded state has smaller hydrodynamic radius than the unfolded which is
644 also more hydrated [40]. The implication of this premise is that $\Delta_N^D V_{prot}$ being small, implies
645 that it's negative magnitude ($-\Delta_N^D V_{prot}$) may be small. The outcome is that the right hand
646 side (RHS) may be large and positive. It must be made clear that $\Delta_N^D(G_{pw})$ needs to be

647 determined but it remains outside the scope of this research. This view explains the effect of
648 sucrose which promotes initial preferential hydration of the enzyme before other
649 physicochemical events such osmotic stress effect due to concentration gradient created by
650 the excluded osmolyte.

651 The equation for the osmolation case is given as

$$652 \quad \Delta_N^D(G_{po}) = -\Delta_N^D V_{prot} - \frac{m(1-\phi_{os})}{RT} \quad (24)$$

653 In this research, aspirin was noticed to have reduced the velocity of amylolysis as to imply
654 that the enzyme was destabilised. In this case, the m -value may be large and negative;
655 there may be a positive value of $\Delta_N^D(G_{po})$. This is to say the number of water molecules
656 around the protein decreases while there is a relative excess of the osmolyte around the
657 protein surface domain including some molecule that may have penetrated into the protein
658 3-D structure.

659 A very important deduction one can make is that hydration and dehydration are
660 merely precedent to the initial events, preferential exclusion and binding respectively.
661 Otherwise, upon unfolding due to osmolation, maximum hydration may occur as expected
662 for the unfolded protein [41]; this is clearly the case because if there is excess of the
663 osmolyte on the protein surface, osmotic gradient should be created towards the protein.
664 Diffusion of water towards the unfolded should occur, making available enough water
665 molecules for the hydration of exposed polar groups. This is without prejudice to the initial
666 displacement of weakly bound water by the binding of the osmolyte. On the other hand
667 translational entropy gain of departing water from hitherto hydrated protein due to excluded
668 osmolyte compels the protein to (re)fold. As presented in literature [41] the first order case
669 which seem to be applicable to highly concentrated osmolyte/cosolute, requires the
670 introduction of apparent hydrated molar volume of the former as follows: Equation (23), by
671 so doing, is transformed to

672
$$\Delta_N^D(G_{pw}) = -\Delta_N^D V_{prot} + \frac{m(1-C_3V_1)\phi_{os}}{RT} \quad (25)$$

673 Equation (25) enables the determination of the integral for hydration at none destabilising
 674 concentration of the stabilising osmolyte as long as $1 > C_3V_1$ and $V_1 \ll 1$. This equation is
 675 reserved for future investigation in which the concentration range of sucrose may be 0.25 -
 676 1.25 mol/L. For the purpose of discussion Eq. (25) reminds one of the high molar
 677 concentrations of ethanol explored in this research whose effect requires another equation
 678 slightly different from Eq. (25). The equation is

679
$$\Delta_N^D(G_{po}) = -\Delta_N^D V_{prot} - \frac{m(1-\phi_{os})(1-C_3V_1)}{RT} \quad (26)$$

680 Since ϕ_{os} is the volume fraction of cosolvent (or rather mole fraction which covers non-
 681 solvent and solvents, e.g. sucrose and ethanol respectively), and its value being < 1 means
 682 that $1 - \phi_{os}$ is always > 0 . The implication is that for the osmolation (positive preferential
 683 interaction parameter) case $1 - C_3V_1$ should also be > 0 . This explains the osmolation (and
 684 its effect) whereby $\Delta_N^D(G_{po})$ needs to be positive due to the binding of ethanol alone and
 685 only aspirin in separate assays. Osmolation leads to unfolding and consequently, a decrease
 686 in the amyolytic action of the enzyme as observed. The question that needs to be answered
 687 is, what means can be applied for the determination of V_1 ? The issue of interest is always
 688 the hydration changes linked either to the *ES* or cosolvent inaccessible region of the protein.
 689 This is despite objection against total reliance on flexibility for function considering that some
 690 enzyme activity can occur at very low hydration levels, coupled with a reduction in protein's
 691 flexibility. On the contrary, Poole [36] observed that hydration induced conformation and
 692 dynamic changes are completed just before the onset of enzyme activity which occurs
 693 before all polar groups are hydrated. There was confirmatory evidence via increased alpha -
 694 helicity that leads to increased rigidity in the protein at low hydration (dry); this led to the
 695 deduction that when water is added to the dry enzyme a "loosening up" or increase in

696 flexibility occurs around a threshold of hydration [36]. It appears therefore, that it is an
697 excessive flexibility that leads to total unfolding due to the action of destabilisers that
698 reduces the biological function of the protein as observed in this research.

699 In summary there may be changes in the number of water interacting with **ES**, be it
700 negative or positive. The change in the number of water molecules interacting with the **ES**
701 as a function of ethanol/aspirin concentration indicates dehydration more so with a lower
702 concentration of sucrose. Thus ethanol is destabilising. The change as a function of sucrose
703 concentration with different concentration of ethanol/aspirin shows mixed trend, increasing
704 hydration with lower fixed concentration of ethanol/aspirin and decreasing with higher fixed
705 concentration of ethanol/aspirin. Thus sucrose promotes hydration being a protecting
706 osmolyte.

707 Generally, the change in the number of water molecules (ΔN_W) in an osmolyte
708 inaccessible region of the enzyme as a function of sucrose concentration with different fixed
709 concentration of aspirin/ethanol is positive as to imply hydration. Perhaps, the increasing
710 solubility of raw starch in increasing concentration of ethanol may presumably account for
711 the negative ΔN_W with lower concentration of ethanol. The values of ΔN_W as a function of
712 aspirin/ethanol concentration with different fixed concentration of sucrose are negative as to
713 imply dehydration peculiar to osmolation by destabilising cosolute.

714 The spontaneity of the processes, folding to unfolding transition and accompanying
715 hydration changes, has been illustrated with the quantification of the free energy difference;
716 in line with the approach, the results shows that *ab initio* the equilibrium constant for
717 hydration change ($K_{eq}(1)$) may be $<$ or $>$ equilibrium constant ($K_{eq}(3)$) for folding to
718 unfolding transition. A positive free energy difference means that hydration change is more
719 spontaneous than folding transition which may be attributable to the effect of sucrose. The
720 converse is the case with ethanol/aspirin in which the free energy difference is negative
721 ($K_{eq}(3) > K_{eq}(1)$).

722 For the purpose of interpretation, theories in literature were adopted for the
723 elucidation of results. The model for water stripping effect of aspirin/ethanol, preferential
724 interaction concept and the KBT for KBI for osmolation and hydration guided the
725 interpretation of the root basis of the effects of the cosolutes.

726 5. CONCLUSION

727 The change in the number of water molecules in an osmolyte inaccessible region of
728 the enzyme and those interacting with the **ES** may be either positive or negative due
729 respectively to sucrose and ethanol/aspirin. The spontaneity of two processes, hydration and
730 folding-unfolding transition, the free energy difference, differs. The mathematical model for
731 water stripping, preferential interaction concept, and the KBI for osmolation and hydration
732 can guide the interpretation of the effects of any cosolute.

733

734 **COMPETING INTERESTS DISCLAIMER:**

735 **Authors have declared that no competing interests exist. The products used**
736 **for this research are commonly and predominantly use products in our area of**
737 **research and country. There is absolutely no conflict of interest between the**
738 **authors and producers of the products because we do not intend to use these**
739 **products as an avenue for any litigation but for the advancement of**
740 **knowledge. Also, the research was not funded by the producing company**
741 **rather it was funded by personal efforts of the authors.**

742 **COMPETING INTERESTS**

743 There is no competing interest.

744 **REFERENCES**

- 745 1. Nakagawa H, Kitao JA, Kataoka M. Hydration affects both harmonic and
746 anharmonic nature of protein dynamics. Biophys. J. 2008; 95(6): 2916–2923.
- 747 2. Bernazzani P. Structural changes associated with interactions between starch and
748 particles of TiO₂ ZnSe. J. Chem. Biochem. Mol. Biol. 2008; 2(1) 1 – 13.

- 749 3. Gangadharan D, Nampoothiri KM, Sivaramakrishnan S, Pandey A. Immobilized
750 bacterial α -amylase for effective hydrolysis of raw and soluble starch. *Food Res. Int.*
751 2009; 42: 436-442.
- 752 4. DR. Knight J, Hamelberg D, McCammon AJ, Kothary R. The role of conserved
753 water molecules in the catalytic domain of protein kinases. 2009; 76(3): 527-535.
- 754 5. Affleck R, Xu Z-F, Suzawa V, Focht K, Clark DS, Enzymatic catalysis and dynamics
755 in low-water environments. *Biochemistry.* 1992; 89: 1100 -1104.
- 756 6. Kurkal V, Daniel RM, Finney JL, Tehei M, Dunn RV, Smith JC. Enzyme Activity and
757 Flexibility at Very Low Hydration. *Biophys. J.* 2005; 89: 1282–1287.
- 758 7. Laage D, Elsaesser T, Hynes JT. Water dynamics in the hydration shells of
759 biomolecules. *Chem. Rev.* 2017; 117:10694-10725.
- 760 8. Maharolkar AP, Murugkar AG, Khirade PW, Mehrotra SC. Microwave dielectric
761 relaxation and polarization study of binary mixture of methylethylketone with
762 nitrobenzene. *Bull. Chem. Soc. Ethiop.* 2019; 33(2): 349-358.
- 763 9. Buurma, NJ, Pas Torello L, Blandermer JM, Engberts JBFN. Kinetic evidence for
764 hydrophobically stabilized encounter complexes formed by hydrophobic esters in
765 aqueous solutions containing monohydric alcohols. *J. Am. chem. Soc.* 2001; 123:
766 11848 – 11853.
- 767 10. Udema II, Onigbinde AO. Effect of interacting organic co-solutes with enzyme
768 substrate complex on the hydrolysis of raw soluble starch with α -amylase:
769 Theory and experimentation. *Adv. Res.* 2016; 7(1): 1-19.
- 770 11. Schnell S, Maini PK. Enzyme kinetics at high enzyme concentration. *Bull. Math. Biol.*
771 2000; 62: 483–499.
- 772 12. Udema II. Substrate mass conservation in enzyme catalyzed amylolytic activity. *Int.*
773 *J. Biochem. Res. Rev.* 2017; 18 (1): 1-10.

- 774 13. Mitchell DC, Litman DT. Effect of ethanol and osmotic stress on receptor
775 conformation reduced water activity amplifies the effect of ethanol on Metarhodopsin
776 II formation. *J. Biol. Chem.* 2000; 275 (8): 5355-5360.
- 777 14. Sirotkin VA, Kuchierskaya AA. Alpha-Chymotrypsin in water-ethanol mixtures: Effect
778 of preferential interactions. *Chem. Phys. Lett.* 2017; 689: 156-161.
- 779 15. Udema II, Onigbinde AO. Basic Kirkwood –Buff theory of solution structure and
780 appropriate application of Wyman linkage equation to biochemical phenomena.
781 *Asian J. Phys. Chem. Sci.* 2019; 7(1):1-14.
- 782 16. Sheng Y, Capri J, Waring A, Valentine JS, Whitelegge J. Exposure of solvent-
783 inaccessible regions in the amyloidogenic protein human SOD1 determined by
784 hydroxyl radical foot printing. *J. Am. Soc. Mass Spectrum.* 2019; 30 (2): 218-226.
- 785 17. Blundell TL, Gong S. Structural and functional restraints on the occurrence of single
786 amino acid variations in human proteins. *PLoS one.* 2010; 5(2): e9186, 1-12.
- 787 18. Gains JC, Acebes S, Virueta A, Butter M, Regan L, O' Hern CS. Comparing side
788 chain packing in soluble proteins, protein-protein interfaces, and transmembrane
789 proteins. *Proteins.* 2018; 85 (5):581-591.
- 790 19. Bolen DW, Baskakov IV. The osmophobic effect: Natural selection of a
791 thermodynamic force in protein folding. *J. Mol. Biol.* 2001; 310 (5): 955-963.
- 792 20. Pace CN. Measuring and increasing protein stability. *Trends Biotechnol.* 1990; 8: 93
793 – 98.
- 794 21. Udema II, Onigbinde AO. Activity coefficient of solution components and salts as
795 special osmolyte from Kirkwood-Buff theoretical perspective. *Asian Res. Biochem.*
796 2019; 4(3): 1-20.
- 797 22. Lineweaver H, Burk D. The determination of enzyme dissociation constants. *J. Am.*
798 *Chem. Soc.* 1934; 56: 658–666.

- 799 23. Kramer RM, Shende VR, Motl N, Pace N, Scholtz JM. Toward a molecular
800 understanding of protein solubility: Increased negative surface charge correlates
801 with increased solubility. *Biophys. J.* 2012; 102: 1907–1915.
- 802 24. Bernfeld P. Amylases, alpha and beta. *Methods. Enzymol.* 1955;1:149–152.
- 803 25. Udemia II. The effect of additives and temperature on the velocity of hydrolysis of raw
804 starch with human salivary α - amylase. *J. Biochem. Res. Rev.* 2016; 10(2): 1-17.
- 805 26. Sanyal NS, Kaushal N. Effect of two non-steroidal anti-inflammatory drugs, aspirin
806 and nimesulide on the G-glucose transport and disaccharide hydrolases in the
807 intestinal brush border membrane *Pharmacol Rep* 2005; 57: 833-838.
- 808 27. de Piña MZ, Saldaña-Balmori Y, Hernández-Tobias A, Piña E. Nonsteroidal
809 anti-inflammatory drugs lower ethanol-mediated liver increase in lipids and
810 thiobarbituric acid reactive substances. *Alcohol Clin Exp Res* 1994; 17(6):1228-
811 12232.
- 812 28. Levitt M, Sharon R. Accurate simulation of protein dynamics in solution. *Proc. Nat.*
813 *Acad. Sci. U.S.A.* 1988; 85: 7557-7561.
- 814 29. Petukhov M, Rychkov G, Firsov L., Serrano L. H-bonding in protein hydration
815 revisited. *Protein Sci.* 2004; 13 (8): 22120-2129.
- 816 30. Ooi T, Oobatake M. Prediction of the thermodynamics of protein unfolding: The
817 helix-coil transition of poly(L-alanine). *Biochemistry.* 1991; 88: 974-975.
- 818 31. Yang L, Dordick JS, Garde S. Hydration in nonaqueous media is consistent with
819 solvent dependence of its activity. *Biophys. J.* 2004; (87): 812-821.
- 820 32. Dai L, Klibanov AM. Striking activation of oxidative enzymes suspended in
821 nonaqueous media. *Proc. Nat. Acad. Sci. U.S.A.* 1999; 96: 9475 - 9478.
- 822 33. Zaks A, Klibanov AM. The effect of water on enzyme action in organic media. *J. Biol.*
823 *Chem.* 1988; 263 (17): 8017 - 8021.
- 824

- 825 34. Pal SKJ, Zewail AH. Biological water at the protein surface, dynamical solvation
826 probed directly with femtosecond resolution. Proc. Nat. Acad. Sci. U.S.A. U.S.A.
827 2001; 94 (4):1763-1768.
- 828 35. Csermely, P. (2001). Water and cellular folding process. Cell. Mol. Biol. 47 (5): 1-9.
- 829 36. Poole P. Hydration and enzymatic activity. J. Phys. Colloq. 1984; 45(c7): 249-253.
- 830 37. Timasheff SN. Protein solvent preferential interaction, protein hydration, and the
831 modulation of biochemical reactions by solvent components. Biochemistry. 2002; 99(15):
832 9721-9726.
- 833 38. Harano Y, Kinoshita M. Translational-entropy gain of the solvent upon protein
834 folding. Biophys. J. 2005; 89: 2701-2710.
- 835 39. Robinson CR, Sligar SG. Molecular recognition mediated by bound water: A
836 mechanism for star activity of the restriction endonuclease *EcoRI*. J.Mol. Biol. 1993;
837 234(2): 302-306.
- 838 40. Lynch TW, Sligar SG. Macromolecular hydration changes associated with *BamHI*
839 binding and catalysis. J. Biol. Chem. 2000; 275 (39): 30561-30565.
- 840 41. Rösgen J, Pettit MB, Bolen DW. Protein folding, stability, and solvation structure in
841 osmolyte solution. Biophys. J. 2005; 89: 2988–2997.
- 842 42. Bolen DW, Baskakov IV. The osmophobic effect: Natural selection of a
843 thermodynamic force in protein folding. J. Mol. Biol. 2001; 310 (5): 955-963.
- 844 43. Baskakov I, Bolen DW. Forcing thermodynamically unfolded proteins to fold
845 (communication). J. Biol. Chem. 1998; 273 (9): 1-5.
- 846 44. Schellman JA. Protein stability in mixed solvents: A balance of contact interaction
847 and excluded volume. Biophys J. 2003; 85:108 -105.

848

849

850

851

852

853

854

855

856

857

858

859

860