

**EFFECT OF ORGANIC MANURE AND MULCHING ON
THE GROWTH AND YIELD OF CARROT (*Daucus
carota* L.)**

ABSTRACT

An experiment on effect of organic manure & mulching on the growth and yield of carrot was conducted at the farm of Sher-e-Bangla Agricultural University, Dhaka during the period from November 2017 to February 2018. The research consisted in Randomized Complete Block Design (RCBD) with three replications under a factorial arrangement (of two factors), with Four levels of organic manure, as the factor A and four mulches such as factor B. Organic manure and mulching influenced significantly ($p=0.05$) all the studied variables. In case of organic manure the maximum plant height (44.55 cm) root length (14.73 cm), root diameter (4.11 cm), root weight (124.50 g), root yield (24.90 t/ha) and marketable root yield (23.85 ton/ha) were recorded from O₂ (Vermicompost) treatment. In case of mulches, maximum plant height (44.81 cm), root length (15.20 cm), root diameter (3.91 cm), root weight (117.85 g), root yield (23.57 t/ha) and marketable root yield (21.95 t/ha) found in M₂ (Black polythene) treatment. And the combined effect of the highest root yield (29.06 t/ha) was obtained from O₂M₂ (Vermicompost + black polythene) and lowest (13.20 t/ha) from O₀M₀ (control). So, it can be concluded that 10 ton/ha vermicompost with black polythene mulch was the best for carrot cultivation. The highest gross return (Tk. 563200), net return (Tk. 446355) and BCR (4.82) were obtained from the treatment combination from O₂M₂ where the lowest gross return (Tk. 207800), net return (Tk.112755) were obtained from O₀M₀ and lowest BCR (1.67) from O₁M₀.

Keywords: *Daucus carota*, growth, mulching, organic manure, yield

1. INTRODUCTION

Carrot (*Daucus carota* L.) is one of the most ancient vegetable. It is grown in spring, summer and autumn in temperate countries and during winter in tropical and subtropical countries [1] and extensively cultivated in North and South America, Europe, Asia, North Africa [2]. It belongs to Apiaceae family and said to be originated in Mediterranean region and its cultivation as a crop also began in that region [3]. It has high nutritive value and possible diversified use in making different palatable foods like soups, stews, curries, salad, pickles, halua and jam. It contains appreciable amount of carotene, which can contribute a lot to overcome blindness of children in Bangladesh. The carrot is one of the profitable crops in Bangladesh. The edible part of this crop is characterized by its high beta carotene content, a precursor of vitamin A [4] and acts as an excellent source of iron, calcium, phosphorus, vitamin B, sugar and folic acid. It has been reported that the entire production of carrot was 14075 metric tons under 1415 ha in

27 Bangladesh during 2009-2010 year [5]. An average yield of carrot is about 25 tons/ha in Bangladesh [6]
28 which are comparatively low from major carrot producer countries [7]. It grows successfully in Bangladesh
29 during Rabi season when temperature ranges from 11.17°C to 28.9°C [8] and mid November to early
30 December is the best time for its cultivation to get satisfactory yield [9]. The cultivation of carrot requires
31 an ample supply of plant nutrient. Use of organic manure is essential for its proper growth and
32 development. Organic matter content of Bangladesh soil is below 1% in about 60% cultivable land as
33 compared to an ideal minimum value of 5% organic matter. In the area of continuous cropping, organic
34 matter is supplied to soil through cowdung, compost, poultry manure etc. Several attempts have been
35 made to increase the yield potential of root crops, but farmers are concerned with the use of inorganic
36 fertilizers which results in decrease fertility of soil, soil health, contents of organic matter and decreases
37 the microbial activity of soil [10]. Vermicompost is the product of the composting process using various
38 species of worms, usually red wigglers, white worms, and other earthworms, to create a mixture of
39 decomposing vegetable or food waste, bedding materials. is regarded ecologically sound bio-fertilizer and
40 also cost-effective and eco-friendly [11]. Vermicompost is a potential source of readily available nutrients,
41 growth enhancing substances and a number of beneficial micro-organisms like N-fixing, P- solubilizing
42 and cellulose decomposing organisms [12,11]. It has been studied that the vermicompost effects on the
43 plant growth, yield and quality of crops considerably. As a result, the seed germination, vigour, flowering
44 and fruiting of plant, tuberization, development of root size colour shelf life and quality are apparently
45 improved [13,14]. On an average Vermicompost contains 1.6% N, 0.7% P, 0.8% K, 0.5% Ca and 0.2%
46 Mg. (Source: Panjab State Council for Science and Technology). In Bangladesh cowdung is used as an
47 organic manure. Due to inadequate knowledge about the merits of organic manure, the farmers
48 habituated extensive use of inorganic fertilizers. On an average, well rotten cowdung contains 0.5% N,
49 0.2% P₂O₅ and 0.5% K₂O [15]. In Bangladesh carrot is grown during winter season when the rainfall is
50 scanty. So irrigation is essential for cultivation. But it increases the cost of cultivation. Under such
51 condition mulching may be useful in reducing irrigation requirement. To serve this purpose water
52 hyacinth, wood ash and black polythene sheets may be used as mulching materials. Mulching is highly
53 effective in checking evaporation loss of soil moisture. Mulching protects the loss of soil moisture by soil
54 evaporation induced by wind and reduces the irrigation requirement [16]. It increases the efficiency of
55 applied N-fertilizer and irrigation [17]. Different mulches regulate soil moisture and temperature, suppress
56 weeds and improve germination and emergence [18]. In addition, mulches increase microbial activity in
57 the soil [19]. Higher yield and better quality, less infestation of insects diseases, earliness, prolong
58 growing season, higher nutritive value of the produced, improved storability are the advantages of
59 mulching [20]. In some extent, mulches reduce the invasion of insects and diseases [21].

61 2. MATERIAL AND METHODS

62 2.1 Experimental site

63 The experiment was conducted at the Horticulture Farm of the Sher-e-Bangla Agricultural University,
64 Dhaka during November, 2017 to February, 2018. Laboratory works were done at Horticulture Laboratory
65 in Sher-e-Bangla Agricultural University, Dhaka-1207. Location of the site is 23°74' N latitude and 90°35'
66 E longitude with an elevation of 8 meter from sea level in Agro-ecological zone of "Madhupur Tract" (AEZ-
67 28). The soil was sandy loam in texture having pH 5.46- 5.62 and EC 0.60 dS/m.

68 2.2 Experiment Frame Work

69 The research was consisted of two factors: Factor A: four level of organic manure as O₀ = No organic
70 manure, O₁= Cowdung (20 ton/ha), O₂ = Vermicompost (10 ton/ha), O₃ = Cowdung(10 ton/ ha) +
71 Vermicompost (5 ton/ha). Factor B: Different type of mulches M₀ = No mulch, M₁ = Water hyacinth M₂
72 =Black polythene M₃ = Wood ash. The two factor experiment was laid out in a Factorial Completely
73 Randomised Block Design (FCRBD) with three replications. The whole experimental area was 24.5m x
74 5.75m which was divided into three blocks. Each block was again divided into 16 plots and hence there
75 were 48 (16 x 3) unit plots. The treatments were assigned randomly in each block separately. The size of
76 unit plot was 1.25m x 1.0m. The distance between two adjacent blocks and plots were 1.0 m and 0.5 m
77 respectively.
78
79

80

81 **2.3 Manure and fertilizer application:**

82 The sources of applied N, P₂O₅, K₂O were as cowdung, vermicompost. The entire amounts of cowdung
83 and vermicompost were applied during the final land preparation as per treatments.

84

85 **2.4 Application of mulching:**

86 Before sowing of seeds mulching was done with water hyacinth, wood ash and black polythene as per
87 treatments. Fresh water hyacinth as chopped into small pieces (8-10cm) and dry then placed over the
88 plots with a thickness of 12cm approximately. Black polythene sheet with small holes at plant distance
89 was spread over the plots accordingly so that the seedlings could emerge easily through the holes.

90

91 **2.5 Economic analysis**

92 In computing economics, the varying levels of organic manure and different types of mulches were taken
93 into consideration apart from other costs common to all the treatments as per package of practices. The
94 benefit cost ratio (BCR) was calculated as follows:

95 $BCR = \text{Gross return per hectare (Tk.)} \div \text{Cost of production per hectare (Tk.)}$

96

97 **2.6 Statistical analysis**

98 The recorded data on different growth and yield parameters were calculated for statistical analysis.
99 Analyses of variances (ANOVA) for most of the characters under consideration were performed with the
100 help of MSTAT program. Treatment means were separated by Duncane's Multiple Range Test (DMRT) at
101 5% level of significance for interpretation of the results [29].

102


103 **3. RESULTS AND DISCUSSION**

104

105 **3.1 Plant height**

106 Plant height of carrot has measured at 30, 50, 70 days after sowing and at harvest. Figure 1, Showing
107 that the maximum plant height was found (44.55cm) from the O₂ (Vermicompost 10 ton/ha) treatment at
108 harvest, while the minimum height was observed (38.99cm) from O₀ (control) treatment. Rashid and
109 Shakur, [22] reported similar results in plant height. Among the mulches materials treatment M₂ (black
110 polythene) mulch produced the tallest plant (18.88 cm, 33.81 cm, 41.01 cm and 44.81 cm) at 30, 50, 70
111 DAS and at harvest respectively and followed by the M₁ (Water hyacinth) mulch at the same DAS. The
112 shortest plant was observed for the treatment M₀ (control) (Figure 2).


113 The combined effect of organic manure and mulching was found significantly influenced in terms of plant
114 height of carrot (Table 1). The maximum plant height (23.16 cm, 36.37 cm, 42.77 cm and 48.13 cm) was
115 recorded from O₂M₂ (Vermicompost, 10 ton/ha + black polythene mulch) treatment at 30, 50, 70 DAS and
116 at harvest respectively. On the other hand, the minimum plant height (12.00 cm, 17.22 cm, 30.93 cm and
117 35.43 cm) was found in plants of control plot O₀M₀ (No organic + no mulch) treatment at 30, 50, 70 DAS
118 and at harvest respectively. Rashid and Shakur, [22] reported that combined application of 6 ton/ha
119 vermicompost with water hyacinth mulch in carrot was the best for obtaining plant height, number of
120 leaves, root length, root fresh weight and highest marketable yield.



121
122
123
124
125
126

Figure 1: Effect of different organic manure on plant height at different days after sowing

[O₀ = No organic manure, O₁= Cowdung (20 ton/ha), O₂ = Vermicompost (10 ton/ha), O₃ = Cowdung (10 ton/ha) + Vermicompost (5 ton/ha)]



127
128
129
130
131
132

Figure 2: Effect of different mulches on plant height at different days after sowing

[M₀ = No mulch, M₁ = Water hyacinth, M₂ = Black polythene, M₃ = Wood ash]

3.2 Number of leaves per plant

The maximum number of leaves per plant (5.58, 10.48 and 11.72) was recorded from O₂ (Vermicompost, 10 ton/ha) treatment at 50, 70 DAS and at harvest respectively. The minimum leaf number (4.36, 8.25 and 10.24) was found from O₀ (control) treatment respectively for same DAS (Figure 3). Azarme et al.,[23] state that the addition of vermicompost at ratio of 15 ton/ha, significantly increased plant growth and yield compared to control.

133
134
135
136
137
138
139
140

Table 1: Combined effect of organic manure and mulching on plant height at different days of carrot


Treatments	Plant height (cm)			
	30 DAS	50 DAS	70 DAS	Harvest

O ₀ M ₀	12.00	17.22	30.93	35.43
O ₀ M ₁	13.86	24.47	37.47	40.07
O ₀ M ₂	14.36	32.20	39.33	42.33
O ₀ M ₃	12.40	21.90	31.07	38.13
O ₁ M ₀	13.47	20.30	31.67	35.73
O ₁ M ₁	16.73	26.90	38.93	43.00
O ₁ M ₂	17.57	32.40	40.67	43.67
O ₁ M ₃	15.73	24.80	35.53	39.87
O ₂ M ₀	17.87	26.83	37.13	43.53
O ₂ M ₁	18.53	29.50	40.73	46.07
O ₂ M ₂	23.16	36.37	42.77	48.13
O ₂ M ₃	17.87	28.67	38.1	40.47
O ₃ M ₀	15.20	25.10	35.67	40.87
O ₃ M ₁	16.87	27.00	39.47	44.80
O ₃ M ₂	20.47	34.26	41.27	45.13
O ₃ M ₃	15.87	26.83	36.73	40.40
CV (%)	8.41	9.67	9.25	10.98
LSD (0.05)	2.53	2.21	4.29	4.04

141 [O₀ = No organic manure, O₁ = Cowdung (20 ton/ha), O₂ = Vermicompost (10 ton/ha), O₃ = Cowdung (10 ton/ha) +
 142 Vermicompost (5 ton/ha), M₀ = No mulch, M₁ = Water hyacinth, M₂ = Black polythene, M₃ = Wood ash]
 143

144 At 50, 70 DAS and at harvest, the maximum number of leaves per plant (5.56, 10.55 and 12.08) was
 145 obtained from M₂ (Black polythene) treatment while the minimum number of leaves per plant (4.55, 8.15
 146 and 10.02) at the same DAS was found from M₀ (No mulch) treatment (Figure 4). Jaysawal et al., [24]
 147 reported that the treatment black polythene mulch was best among the various mulch treatments and
 148 recorded maximum plant height, number of leaves per plant, leaf fresh weight, leaf dry weight, root
 149 weight, root length, root diameter and total root yield of carrot.

150 The combined effect of different organic manure and mulch materials showed significant differences due
 151 to their application on number of leaves per plant of carrot at 50, 70 DAS and at harvest (Table-2). The
 152 maximum number of leaves per plant At 50, 70 DAS and at harvest was recorded 6.50, 11.77 and 13.00
 153 respectively from treatment combination O₂M₂ (Vermicompost, 10 ton/ha + black polythene) while the
 154 minimum number of leaves per plant at 50, 70 DAS and at harvest were 3.73, 7.06 and 9.16, respectively
 155 from O₀M₀ (No organic manure + no mulch).
 156



157
 158
 159 **Figure 3: Effect of different organic manure on no. of leaves per plant at different days after**
 160 **sowing**

161 [O₀ = No organic manure, O₁ = Cowdung (20 ton/ha), O₂ = Vermicompost (10 ton/ha), O₃ = Cowdung (10 ton/ha) +
 162 Vermicompost (5 ton/ha)]
 163




Figure 4: Effect of different mulches on no. of leaves per plant at different days after sowing
 [M₀ = No mulch, M₁ = Water hyacinth, M₂ = Black polythene, M₃ = Wood ash]

164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203

3.3 Root length (cm)

The longest root length 14.73 cm was recorded from O₂ (Vermicompost (10 ton/ha) treatment while the shortest root length 11.91 cm was observed from control plot (Table 3). This finding is an agreement with the result of Schuch et al., [25], they reported that the root length of carrot varied with different amount of manure application. Root length differed significantly due to the different mulch application. Maximum root length (15.21 cm) was recorded at treatment M₂ (Black polythene) and it was significantly different than other treatments. The minimum root length (11.96 cm) was found at M₀ (No mulch) (Table 5). This result is in accordance with the findings of Rahman et al., [26]. The combined effect of organic manures and mulches showed significant variation on root length of carrot (Table 7). The longest root 17.00 cm was observed from the treatment combination of O₂M₂ (Vermicompost, 10 ton/ha + black polythene). The shortest root length (10.50 cm) was recorded from control O₀M₀ (No organic manure + no mulch) treatment.

3.4 Root diameter (cm)

Diameter of root was significantly influenced by the application of different level of organic manure (Table 3). The maximum diameter of root (4.12 cm) was recorded from O₂ (Vermicompost, 10 ton/ha) treatment and it was significantly different than other organic manure treatments. On the other hand, the minimum root diameter (3.14 cm) was observed from O₀ treatment (control). Root diameter was also significantly varied due to the use of different mulching materials in carrot (Table 5). The highest root diameter (3.92 cm) was obtained at the mulches treatment M₂ (Black polythene). The lowest root diameter of root (3.15 cm) was obtained at the treatment of M₀ (No mulch). The maximum diameter of root (4.50 cm) was observed from the treatment combination of O₂M₂ (Vermicompost, 10 ton/ha + black polythene). The minimum diameter of root (2.43 cm) was recorded from control O₀M₀ (No organic manure + no mulch) treatment (Table 7).

3.5 Root weight per plant (g)

A significant variation was observed on root weight per plant due to use of different organic manures (Table 3). The maximum root weight per plant 124.50 g was recorded from O₂ (Vermicompost, 10 ton/ha) treatment. While the minimum root weight (47.58 g) was recorded from control plots. Lang [27] found that organic manures increased the yield of carrot (10%-20%) compared with control. Different mulches materials showed significant variation for root weight per plant of carrot (Table 5). The highest root weight per plant (117.85 g) was found from M₂ (Black polythene). The lowest root weight per plant (92.23 g) was recorded from control treatment. The maximum root weight per plant (82.53 g) was observed from treatment combination of O₂M₂ (Vermicompost, 10 ton/ha + black polythene) treatment. The minimum root weight per plant (35.73 g) was recorded from control plot O₀M₀ (No organic manure+ no mulch) (Table 7).

204 **Hasan et al.**, [28] reported that combined application of vermicompost (6 ton/ha) and water hyacinth
 205 mulching gave the highest marketable (27.89 ton/ha) and the gross yield (29.48 ton/ha) of carrot.

206 **3.6 Root yield per plot (kg)**

207 The maximum root weight per plot 3.11 kg was recorded from O₂ (Vermicompost, 10 ton/ha) treatment.
 208 While the minimum root weight (2.13kg) from control plots (Table 3). Different mulch materials showed
 209 significant variation for root weight per plot of carrot (Table 5). The highest root weight per plot (2.95 kg)
 210 was found from M₂ (Black polythene) treatment. The lowest root weight per plot (2.31kg) was recorded
 211 from control. The combined effect of organic manures and mulch materials was found significant variation
 212 was observed on root weight per plot (Table 7). The maximum root weight per plot (3.63 kg) was
 213 observed from treatment combination of O₂M₂ (Vermicompost, 10 ton/ha + black polythene). The
 214 minimum root weight per plot (1.65 kg) was recorded from control plot O₀M₀ (No organic manure + no
 215 mulch).
 216

217 **3.7 Yield per ha (ton)**

218 The root yield (24.90 ton/ha) recorded maximum from O₂ (Vermicompost, 10 ton/ha) treatment. The
 219 minimum root yield (17.00 ton/ha) was obtained from control O₀ (No organic manure) treatment (Table 4).
 220 The highest root yield (23.57 ton/ha) was recorded from M₂ (Black polythene) treatment, while the lowest
 221 (18.45 ton /ha) was obtained from the control (Table 6). The combined effect of organic manure and
 222 mulches was significantly varied on root yield (Table 8). However, the maximum root yield (29.07 ton/ha)
 223 was obtained from the treatment combination of O₂M₂ (Vermicompost, 10 ton/ha + black polythene) which
 224 was statistically significant different from the other treatments; whereas the minimum yield (13.20 ton/ha)
 225 was recorded from control plot O₀M₀ (No organic manure + no mulch).
 226

227 **3.8 Marketable yield (ton/ha)**

228 The marketable root yield (23.85 ton/ha) recorded maximum from O₂ (Vermicompost, 10 ton/ha)
 229 treatment. The minimum root yield (13.82 ton/ha) was obtained from control O₀ (No organic manure)
 230 treatment (Table 4). The highest marketable yield per ha (21.95 ton) was recorded from M₂ (Black
 231 polythene) treatment, while the lowest (15.63 ton/ha) was obtained from the control (Table 6).
 232 The combined effect of organic manures and mulches was significantly varied on Marketable yield per ha
 233 (Table 8). However, the maximum Marketable root yield per ha (28.16 ton) was obtained from the
 234 treatment combination of O₂M₂ (Vermicompost, 10 ton/ha + black polythene) which was statistically
 235 significant different from the other treatments; whereas the minimum yield (10.39 ton) was recorded from
 236 control plot O₀M₀ (No organic manure + no mulch).
 237

238 **3.9 Dry matter percentage of root (%)**

239 The maximum dry matter of root per plant (16.06N%) was recorded from O₂ (Vermicompost, 10 ton/ha)
 240 treatment while the minimum (12.16 %) from control plots (Table 4). Root dry weight was markedly
 241 increased and achieved maximum (15.46 %) values in treatment of M₂ (Black polythene) treatment. The
 242 lowest root dry weight (13.36 %) was observed in control (Table 6).
 243 A significant effect of organic manures and mulches combination on root dry weight was found (Table 8).
 244 The highest values of root dry weight (17.78 %) were recorded due to O₂M₂ (Vermicompost, 10 ton/ha +
 245 black polythene) treatment. On opposition to, the lowest ones (10.90 %) were proceeded from
 246 combination O₀M₀ (No organic manure + no mulch).
 247

248 **Table 2: Combined effect of organic manure and mulching on leaves number per plant at different**
 249 **days of carrot**

Treatments	Leaves number		
	50 days	70 days	Harvest
O ₀ M ₀	3.73	7.06	9.17
O ₀ M ₁	4.0	8.13	10.40
O ₀ M ₂	5.0	9.27	10.80
O ₀ M ₃	4.6	8.53	10.60
O ₁ M ₀	4.67	7.80	9.27

O ₁ M ₁	5.00	8.47	11.20
O ₁ M ₂	5.20	10.20	12.00
O ₁ M ₃	4.47	8.47	10.27
O ₂ M ₀	5.13	9.53	11.53
O ₂ M ₁	6.00	9.93	11.60
O ₂ M ₂	6.50	11.77	13.00
O ₂ M ₃	4.67	10.67	10.73
O ₃ M ₀	4.67	8.20	10.13
O ₃ M ₁	5.20	8.63	11.14
O ₃ M ₂	5.50	10.95	12.50
O ₃ M ₃	4.80	9.13	10.20
CV (%)	8.16	9.68	10.58
LSD (0.05)	0.83	0.85	0.38

250 [O₀ = No organic manure, O₁= Cowdung (20 ton/ha), O₂ = Vermicompost (10 ton/ha), O₃ = Cowdung (10 ton/ha) +
 251 Vermicompost (5 ton/ha), M₀ = No mulch, M₁ = Water hyacinth, M₂ =Black polythene, M₃ = Wood ash]

252 **Table 3: Effect of organic manure on yield parameters at harvest stage**

Organic Manure	Root length (cm)	Root diameter (cm)	Root weight per plant (g)	Root weight per plot (kg)
O ₀	11.91	3.14	85.83	2.13
O ₁	13.19	3.42	96.67	2.42
O ₂	14.74	4.12	124.50	3.11
O ₃	13.30	3.73	116.41	2.81
CV (%)	11.45	12.62	12.43	10.27
LSD (0.05)	0.86	0.29	5.39	0.22

253

254 **Table 4: Effect of organic manure on yield parameters at harvest stage**

Organic Manure	Yield (ton/ha)	Marketable yield (ton/ha)	Root dry matter (%)
O ₀	17.0	13.82	12.16
O ₁	19.33	16.92	12.86
O ₂	24.90	23.85	16.06
O ₃	22.29	20.39	13.97
CV (%)	11.56	10.37	11.42
LSD (0.05)	1.92	1.57	0.45

255

256 **Table 5: Effect of different mulching on yield parameters at harvest stage**

Mulching	Root length (cm)	Root diameter (cm)	Root weight Per plant (g)	Root weight Per plot (kg)
M ₀ (No mulch)	11.98	3.15	92.23	2.31
M ₁ (Water hyacinth)	12.98	3.75	109.39	2.71
M ₂ (Black polythene)	15.21	3.92	117.85	2.95
M ₃ (Wood ash)	13.57	3.59	103.93	2.59
CV (%)	11.45	12.62	12.43	10.27
LSD (0.05)	0.78	0.22	4.37	0.18

257

258 **Table 6: Effect of different mulching on yield parameters at harvest stage**

Organic Manure	Yield (ton/ha)	Marketable yield (ton/ha)	Root dry matter (%)
O ₀	18.45	15.63	12.62
O ₁	21.71	19.24	13.87
O ₂	23.57	21.95	15.46
O ₃	20.78	18.17	13.11
CV (%)	11.56	10.37	11.42

LSD (0.05)

1.13

1.31

0.41

259

260

Table 7: Combined effect of organic manure and mulching on yield parameters of Carrot

Treatments	Root length (cm)	Root diameter (cm)	Root weight Per plant (g)	Root weight Per plot (kg)
O ₀ M ₀	10.5	2.43	66	1.65
O ₀ M ₁	12.13	3.44	96.27	2.32
O ₀ M ₂	13.93	3.51	98.40	2.46
O ₀ M ₃	11.07	3.16	82.67	2.07
O ₁ M ₀	11.86	2.87	76.67	1.92
O ₁ M ₁	13.00	3.60	102.27	2.56
O ₁ M ₂	14.23	3.65	106.40	2.66
O ₁ M ₃	13.67	3.55	101.33	2.53
O ₂ M ₀	12.93	3.81	114	2.85
O ₂ M ₁	13.74	4.17	122.53	3.06
O ₂ M ₂	17.00	4.50	145.33	3.63
O ₂ M ₃	15.27	4.00	116.13	2.90
O ₃ M ₀	12.60	3.49	112.27	2.81
O ₃ M ₁	13.07	3.80	116.50	2.91
O ₃ M ₂	15.67	4.00	121.27	3.03
O ₃ M ₃	14.27	3.65	115.60	2.89
CV (%)	11.45	12.62	12.43	10.27
LSD (0.05)	1.72	0.20	15.79	0.48

261

262

263

Table 8: Combined effect of organic manure and mulching on yield parameters of Carrot

Treatments	Yield (ton/ha)	Marketable yield (ton/ha)	Root dry weight (%)
O ₀ M ₀	13.20	10.39	10.90
O ₀ M ₁	18.59	14.12	12.80
O ₀ M ₂	19.68	16.68	13.91
O ₀ M ₃	16.53	14.08	11.05
O ₁ M ₀	15.33	13.40	11.81
O ₁ M ₁	20.45	19.08	12.85
O ₁ M ₂	21.28	19.96	14.34
O ₁ M ₃	20.24	15.24	12.45
O ₂ M ₀	22.80	20.80	14.95
O ₂ M ₁	24.51	23.36	16.12
O ₂ M ₂	29.07	28.16	17.78
O ₂ M ₃	23.23	23.08	15.42
O ₃ M ₀	22.45	17.92	12.81
O ₃ M ₁	23.31	20.40	13.73
O ₃ M ₂	24.27	23.00	15.83
O ₃ M ₃	23.12	20.26	13.53
CV (%)	11.56	10.37	11.42
LSD (0.05)	3.85	3.18	0.71

264

265

266 **4. CONCLUSION**

267

268 **Based on the results of** the present study, it may be concluded that the efficient production of carrot is
269 increased by the judicious application of organic manure with black polythene. **The experimental result**
270 **revealed** that the organic manure vermicompost (10 ton/ha) gave the highest yield. Mulching materials
271 such as black polythene may be used for higher yield of carrot. Application of Vermicompost (10 ton/ha)
272 with black polythene mulch is one of the most effective management practices to improve soil
273 productivity. Thus, considering crop productivity, economic return and maintaining soil fertility, the
274 combined application of vermicompost (10 ton/ha) with black polythene mulch may be recommended to
275 farmers for **cost-effective** carrot production with affecting the soil health.

276

277 **COMPETING INTERESTS**

278

279 Authors have declared that no competing interests exist.

280

281

282 **REFERENCES**

283

- 284 1. Bose, T. K. and Som. M. G. 1990. Vegetable crops in India, Naya Prakash, Calcutta, India. 408-442.
- 285 2. Thomson H. C, Kelly W. C. 1957. Vegetable crop. (5th Edition) McGraw Hill book, New York, USA.
286 227-335.
- 287 3. Shinohara, S. E. 1984. Technology of Horticultural root crops. Pakistan J. Biol. Sci., 3(2): 13-16.
- 288 4. Chandha, K. L. 2003. Hand book of horticulture. Indian Council of Agricultural Research (ICAR), New
289 Delhi.
- 290 5. BBS, 2010. Yearbook of Agricultural Statistics of Bangladesh. Bangladesh Bureau of Statistics,
291 Ministry of Planning, Dhaka, Bangladesh. p. 124.
- 292 6. Rashidi, M. Abbassi, S. and Gholami, M. 2009. Interactive effects of plastic mulch and tillage method
293 on yield and yield components of tomato (*Lycopersicon esculentum*). American Eurasian J. Agric.
294 Env. Sci., 5: 420-427.
- 295 7. FAO, 2004. Production Year Book. Food and Agriculture Organization, Rome, Italy, 61(2): 99-111.
- 296 8. Alim, A. 1974. An Introduction to Bangladesh Agriculture. 1st Edn, Alim M, Dhaka. p. 9.
- 297 9. Rashid, M. M. 1993. Sabji Biggyan, 1st Edn. Golam Moyenuddin, Director, Text Book Division,
298 Bangla Academy, Dhaka. pp. 502-507.
- 299 10. Chen, Y. P., Liu, Q., Liu, Y. J., Jia, F. A. and He, X. H. 2014. Responses of soil microbial activity to
300 cadmium pollution and elevated CO₂. Sci. Rep., 4(1) 4287-4289.
- 301 11. Amooaghaie, R. and Golmohammadi, S. 2017. Effect of Vermicompost on Growth, Essential Oil, and
302 Health of *Thymus Vulgaris*. Compost sci. utilization 25(3):1-12
- 303 12. Archana, A. B. and Anubha, K. 2011. Standardization of Herbal Drugs: An Overview IRJP. 2 (12), 56-
304 60.
- 305 13. Ansari, A. A. 2008. Effect of vermicompost and vermiwash on the productivity of spinach (*Spinacia*
306 *oleracea*), onion (*Allium cepa*) and potato (*Solanum tuberosum*). World J. Agric. Sci. 4(5): 554-557.
- 307 14. Chanda, G. C., Bhunia, G. and Chakraborty, S. K. 2011. The effect of vermicompost and other
308 fertilizers on the cultivation of tomato plants. J Hortic. Forest., 3(2): 42-45.
- 309 15. Yawalkar, K. S., Agrawal, J. P. and Bokde, S. 1984. Manures and Fertilizers. Agric. Horticultural
310 Publishing House, Nagpur-440010, India. pp. 29-85.
- 311 16. Roy, A. K., Mushi, A. A. and Khan, A. H. 1990. Effect of different mulches on the growth of potato.
Bangladesh J. Bot., 19(1): 41-46.

- 312 17. Rhee, K. M., Yoon, J. H. and Park, J. K. 1990. Effect of polythene film mulching on the changes of
313 soil chemical properties during the sesame cropping season, Res. Rep. Rur. Adm., soil Fert. Abstr.
314 54: 116-27.
- 315 18. Frazier, W. A. 1957. Plastic mulches for horticultural crop. Bull. Agril Exptl. Stat., p. 562.
- 316 19. Aldefer, R. B. 1946. Seasonal variability in aggregation of hagers town silt loam. Soil sci., 62: 151-
317 168.
- 318 20. Ahmed, K. U. 1999. Influence of different mulches on growth and yield of sweet potato. MS Thesis,
319 Department of Crop Botany, Bangladesh Agricultural University, Mymensingh.
- 320 21. Brown, J. E., Stevens, C., Vsborn, M. C. and Bryce, H. M. 1989. Black plastic mulch and spun
321 bonded polyster row cover as method of southern blight control in bell peper. Plant Dis., 73(11): 931-
322 932.
- 323 22. Rashid, M. M. and Shakur, M. A. 1986. Effect of date of planting and duration of growing period on
324 the yield of carrot. Bangladesh Hort., 14(2): 28-32.
- 325 23. Azarmi, R., Ziveh, P. S. and Satari, M. R. 2008. Effect of Vermicompost on Growth, Yield and
326 Nutrition Status of Tomato (*Lycopersicum esculentum*). Pakistan J. Bio. Sci. 11: 1797-1802.
- 327 24. Jaysawal, N., Singh, Dr. G., Kanojia, Dr. A. and Debbarma, B. 2018. Effect of different mulches on
328 growth and yield of carrot (*Daucus carota* L.). Int. J. chem. stud., 6(4): 381-384.
- 329 25. Schuch, S. M. L., Soares M. H. G. and Schuck, E. 1999. Evaluation of carrot cultivars using two
330 sources for organic manures, in Porto Alegre County, RS, Brazil. Pesquisa-Agropecuaria-Gaucha, 5
331 (2): 193-200.
- 332 26. Rahman, M. A., Islam, T., Mamun, M. A. A., Rahman, M. S. and Ashraf, Ms. S. 2018. Yield and
333 Quality Performance of Carrot under Different Organic and Inorganic Nutrient Sources with Mulching
334 Options. Asian J. Agric. Hort. Res. 1(4): 1-8.
- 335 27. Lang, H. 1984. Use of carrot cultivars and organic manures in early carrot growing for reliability, yield
336 and quality. Kar to effelbau. 35(2): 65-69.
- 337 28. Hasan, M. M., Ali, M. A., Rubel, M. M. K., Shah, M., Alzahrani, Y. and Hakeem, K. R. 2018.
338 Influences of Vermicompost and Organic Mulching on Growth, Yield and Profitability of Carrot
339 (*Daucus carota* L.). J. Agri. Bio., 12(4): 34-39.
- 340
- 341 29. Mazed, H. K., Pulok, M. A. I., Chowdhury, M. S. N., & Ferdous, J. (2015). Effect of Different Types of
342 Organic Manure and Mulching on the Growth and Yield of Carrot (*Daucus Carota* L.). *International*
343 *Journal of Scientific and Research Publications*, 309
344
345
- 346
347
348