

Effect of Nitrogen and Phosphorus on the Growth and Seed Yield of Spinach

ABSTRACT

The experiment was conducted at the farm of Sher-e-Bangla Agricultural University, Dhaka, Bangladesh during Robi (November 2017 to March 2018) season to find out the growth, yield and economic benefit of spinach seed as influenced by nitrogen and phosphorus. The research involved two factors. Factor A: Nitrogen management 4 levels; $N_0 = 0$ kg/ha, $N_1 = 27.6$ kg/ha, $N_2 = 55.2$ kg/ha, $N_3 = 82.8$ kg/ha, and factor B: Phosphorus management 4 levels; $P_0 = 0$ kg/ha; $P_1 = 15.84$ kg/ha, $P_2 = 31.68$ kg/ha, $P_3 = 47.52$ kg/ha. There were 16 treatment combinations in the experiment and laid out in Randomized Complete Block Design (RCBD) with three replications. Quality tests of seeds were done based on the germination test (%), seed vigor test (Electrical conductivity). In case of nitrogen, the highest seed yield (1.10 t ha^{-1}), germination percentage (87.33 %) and lowest value in EC test (13.87 dS/cm) were obtained from N_2 , while the lowest seed yield (0.81 t ha^{-1}), germination percentage (79.33 %) and highest value in EC test (11.16 dS/cm) from N_0 . For similar levels of phosphorus, the highest seed yield (1.05 t ha^{-1}), germination percentage (87.08 %) and lowest value in EC test (13.35 dS/cm) were recorded from P_2 , whereas the lowest seed yield (0.84 t ha^{-1}), germination percentage (79.91 %) and highest value in EC test (11.79 dS/cm) from P_0 . Due to mutual effect, the highest seed yield (1.30 t ha^{-1}), germination percentage (91.33 %) and lowest value in EC test (14.83 dS/cm) were noted from N_2P_2 , whereas the lowest seed yield (0.69 t ha^{-1}), germination percentage (72.66 %) and highest value in EC test (10.74 dS/cm) from N_0P_0 . From the economic point of view, the highest Benefit-Cost Ratio (BCR) (1.59) was found in the treatment of N_2P_2 and the lowest BCR (1.01) was found in the treatment of N_0P_0 . It is apparent that the treatment combination N_2P_2 gave the best performance for the seed yield and economic benefit of spinach.

Keywords: Growth, Nitrogen, Phosphorus, Seed Yield

1. INTRODUCTION

Spinach (*Spinacia oleracea*) is a leafy green cool-season vegetable that is known for its nutritive value and is considered one of the most popular vegetables in Bangladesh. It is believed to have originated from Persia. Its leaves are a common edible vegetable. By weight, spinach consists of 91.4% water, 3.6% carbons, and 2.9% protein [1]. There are 23 calories in 100 grams of spinach and further benefits of spinach leaves consumption can be of more value e.g vital mineral elements in spinach compared to other vegetables e.g iron which is very good for women. The seed is produced for commercial consumption and for seed companies that supply seed throughout the country. In Bangladesh spinach occupies 22000 acres [2] with an annual production of 66000 tons. In comparison to other countries, this yields much lower. So, to use of quality seeds of high yield varieties are the foremost important technique for maximizing yield per unit area. Quality seed can increase vegetable production by up to 25-50% [1]. Farmers save seeds annually of about 50 tons and used

every year, which in most cases are of inferior in quality [3]. Appropriate fertilizer application to the plants greatly affects their growth and production. Nitrogen strongly stimulates growth, expansion of the crop canopy and interception of solar radiation [4]. Increasing the levels of nitrogen during the vegetative stage can strengthen and allow a plant to grow more rapidly and produce large amounts of succulent, green foliage, which in turn can generate higher yields [5]. Similarly, phosphorus (P) is an essential nutrient act as catalysts in the conversion of numerous key biochemical reactions in plants. Phosphorus stimulates root development, improves flower formation, seed production and improves crop quality and increases resistance to plant diseases [6]. Leafy vegetables, particularly, spinach is highly responsive to fertilization [7] and oxalates are the main indexes of the quality due to a very efficient uptake system and inefficient reductive systems [8]. The fertilizer requirements on sandy and sandy loams are 85 to 120 kg N, 75 to 85 kg P₂O₅, and 85 to 150 kg K₂O. On heavier clay soils, 75 kg ha⁻¹ of each nutrient should be adequate. If the fertilizer is banded at seeding it should be placed along each side of the rows 2 to 3 inches below the level of the seed and 6 inches to the side of the row; fertilizer should never come in contact with the seed and two or three splits of 85 to 120 kg ha⁻¹ N would be adequate as side-dressing [9]. An adequate supply of fertilizers can promote plant growth and increase crop production, but excessive and inappropriate use of chemical fertilizers causes accumulation of compounds in the edible products which can be detrimental to human health in addition cause environmental pollution and economic losses [10]. [11] found that N increased the spinach yield and enhanced the accumulation of N and P in leaves. [12] reported that application of 40 kg N + 15.0 kg P₂O₅ increased plant fresh yield by 27.2 and 42.3% and 16.3 and 10.4% in seed yield in relation to the control in the first and second seasons, respectively. [13] achieved the highest yield with 120 kg N ha⁻¹ Farmers in Bangladesh generally do not use any improved or special techniques for quality seed production. There for to achieve higher seeds yield with good quality fertilizer management is an important practice which cannot be emphasized.

2. MATERIAL AND METHODS

2.1. Experimental Site

The experiment was conducted at the Agronomy Research Farm of Sher-e-Bangla Agricultural University (SAU), Dhaka, Bangladesh from November 2017 to March 2018. Experimental site situated an elevation of 8 meters above the sea level in the Agro-ecological zone of "Madhupur Tract" (AEZ-28). The soil was sandy loam and medium high land in texture having pH 5.46- 5.62.

2.2 Experiment Frame Work

The research was consisted of two factors: Factor A: Nitrogen management 4 levels; N₀= 0 kg/ha N₁= 27.6 kg/ha, N₂=55.2 kg/ha, N₃= 82.8 kg/ha, and factor B: Phosphorus management 4 levels; P₀= 0 kg/ha; P₁ = 15.84 kg/ha, P₂= 31.68 kg/ha, P₃= 47.52 kg/ha. The two factors experiment was laid out following the Randomized Complete Block Design (RCBD) with three replications. The experiment was divided into three equal blocks where each block was divided into 16 plots. Then 16 treatment combinations were allotted randomly in each block. The size of each unit plot was 1.5 m × 1 m. The distance maintained between two blocks and two plots were 0.75 m and 0.5 m, respectively. Row to row distance was 30 cm and plant to plant distance was 20 cm.

2.3. Application of manure and fertilizers

About 5 t ha⁻¹ well-decomposed cow dung was applied as the control (N₀P₀) treatment) plot and incorporated adequately to the soil during final land preparation whereas other plots were applied with inorganic fertilizer as per treatment. What about N₀ and P₀? Doses of inorganic fertilizers (Urea and Triple superphosphate) were applied in the experimental plot according to the treatments (Table 1). The whole amount of TSP and half the amount of urea

and MoP (180 kg/ha) were also applied as basal dose before sowing of seed in the main field. 1st top dressing of urea was applied when seedlings established in the main field about 10 days after seed sowing. 2nd top dressing of urea and the rest amount of MoP was applied about 25 days after 1st top dressing. Then the rest amount of urea was applied as 3rd installment after flowering. Each top dressing was followed by manual irrigation.

Table 1. Doses of nutrients application in the main field as per treatment

Treatments	Available nutrients (kg ha ⁻¹)	Fertilizers (kg ha ⁻¹)	Doses (g plot ⁻¹)	Treatments	Available nutrients (kg ha ⁻¹)	Fertilizers (kg ha ⁻¹)	Doses (g plot ⁻¹)
	N	Urea	Urea		P	TSP	TSP
N ₀	0	0	0	P ₀	0	0	0
N ₁	27.6	60	9	P ₁	15.84	75	11.25
N ₂	55.2	120	18	P ₂	31.68	150	22.5
N ₃	82.8	180	27	P ₃	47.52	225	33.75

Here, N₀= 0 kg/ha; N₁=27.6 kg/ha; N₂= 55.2 kg/ha; N₃= 82.8 kg/ha; P₀= 0 kg/ha; P₁= 15.84 kg/ha; P₂= 31.68 kg/ha; P₃=47.52 kg/ha.

2.4. Economic analysis

The cost of production was analyzed in order to find out the most economic treatment of nitrogen and phosphorus for quality seed production of spinach. All the non-material and material input costs and interests in running capital were considered for computing the cost of production. The benefit-cost ratio (BCR) was calculated by the following formula:

Benefit-cost ratio (BCR) = Gross return (tk/ha) ÷ Total cost of production (tk/ha)

2.5. Statistical analysis

The data obtained for different characters were statistically analyzed to observe the significant difference among the treatment by using the STATISTIX-10 computer package program. The mean values of all the characters were calculated and analysis of variance was performed. The significance of the difference among the treatments means was estimated by the Least Significant Different Test (LSD) at 5% level of probability.

3. RESULTS AND DISCUSSION

3.1 Plant height (cm)

A significant variation was observed on the plant height of spinach due to the application of different levels of nitrogen (Table 2.). The highest plant height 13.94cm and 47.69cm were obtained before flowering and at the time of harvest respectively from N₃ while the lowest plant height 11.01cm and 34.88cm were obtained before flowering and at the time of harvest respectively from N₀. P₃ showed the highest plant height 13.64cm and 46.68cm before flowering and at the time of harvest respectively and the lowest plant height 11.60 cm and 35.18 cm at before flowering and at the time of harvest respectively in P₀ (Table 3).

The combined effect of different levels of nitrogen and phosphorus application showed a significant effect on the plant height of spinach (Table 4). The highest plant height 15.21cm and 55.25cm before flowering and at the time of harvest, respectively was observed in N₃P₃ while the lowest plant height 10.10cm and 27.7cm before flowering and at the time of harvest, respectively in N₀P₀.

3.2. Number of leaves per plant

121 The effect of nitrogen on the number of leaves per plant of spinach was significant (Table 2).
 122 The highest number of leaves (8.46) was produced from N_2 and the lowest number of leaves
 123 (6.42) was observed in N_0 . P_2 showed the maximum leaves per plant (8.30) and the
 124 minimum leaves per plant (6.36) was observed in P_0 (Table 3).

125 The combined effect of different levels of nitrogen and phosphorus showed a significant
 126 effect on the number of leaves per plant of spinach (Table 4). The maximum leaves per plant
 127 (10.35) were observed in N_2P_2 and the lowest leaves per plant (5.01) were recorded with
 128 N_0P_0 , [14] also found a similar result.
 129

130 **Table 2.** Effect of nitrogen on plant height (before flowering and at the time of harvest)
 131 leaves per plant of spinach

Treatments	Plant height before flowering (cm)	Plant height at the time of harvest (cm)	Leaves per plant
N_0	11.01	34.88	6.42
N_1	12.34	40.35	7.24
N_2	13.07	43.75	8.46
N_3	13.94	47.69	7.84
LSD	0.87	3.12	0.54
CV %	8.34	9.00	8.70

132 Here, $N_0 = 0$ kg/ha; $N_1 = 27.6$ kg/ha; $N_2 = 55.2$ kg/ha; $N_3 = 82.8$ kg/ha

133 **Table 3.** Effect of phosphorus on plant height (before and at the time of harvest), Leaves per
 134 plant of spinach

Treatments	Plant height before flowering	Plant height at the time of harvest	Leaves per plant
P_0	11.60	35.18	6.36
P_1	12.16	40.37	7.24
P_2	12.96	44.44	8.30
P_3	13.64	46.68	8.06
LSD	0.87	3.11	0.543
CV %	8.34	9.00	8.70

135 Here, $P_0 = 0$ kg/ha; $P_1 = 15.84$ kg/ha; $P_2 = 31.68$ kg/ha; $P_3 = 47.52$ kg/ha.

136 **Table 4.** Combined effect of nitrogen and phosphorus on the plant height (before and at the
 137 time of harvesting), leaves per plant of spinach

Treatments	Plant height before flowering (cm)	Plant height at the time of harvest (cm)	Leaves per plant
N_0P_0	10.10	27.7	5.01
N_0P_1	10.63	34.34	6.25
N_0P_2	11.25	37.31	6.66
N_0P_3	12.07	40.18	7.76
N_1P_0	11.49	34.65	6.60
N_1P_1	12.15	41.03	7.33
N_1P_2	12.60	42.44	7.46
N_1P_3	13.12	43.29	7.58
N_2P_0	12.14	37.88	6.91
N_2P_1	12.54	41.80	7.81
N_2P_2	13.42	47.31	10.35
N_2P_3	14.17	48.01	8.76

$N_3 P_0$	12.65	40.51	6.93
$N_3 P_1$	13.35	44.30	7.56
$N_3 P_2$	14.56	50.71	8.73
$N_3 P_3$	15.21	55.25	8.13
LSD	1.75	6.25	1.08
CV %	8.34	9.00	8.70

Here, $N_0 = 0$ kg/ha; $N_1 = 27.6$ kg/ha; $N_2 = 55.2$ kg/ha; $N_3 = 82.8$ kg/ha.
 $P_0 = 0$ kg/ha; $P_1 = 15.84$ kg/ha; $P_2 = 31.68$ kg/ha; $P_3 = 47.52$ kg/ha.

3.3. Number of inflorescence per plant

Statistically significant differences were found on the number of inflorescence per plant of spinach due to the application of nitrogen (Table 5). The highest number of inflorescence per plant (8.03) was recorded from N_2 whereas, the lowest number (5.17) was observed from N_0 . The highest number of inflorescence per plant (7.80) was recorded from P_2 and the lowest (4.94) was found from P_0 (Table 6).

The combined effect of nitrogen and phosphorus showed significant variation in the number of inflorescences per plant (Table 7). The highest number of inflorescence per plant (10.53) was recorded from $N_2 P_2$ and the lowest number of inflorescence per plant (3.90) from $N_0 P_0$.

3.4. Length of inflorescence

A significant variation was observed on the length of an inflorescence of spinach when different levels of nitrogen were applied (Table 5). The highest length of inflorescence (29.54cm) was recorded in N_2 and the lowest length of inflorescence (19.09 cm) from N_0 . The highest length of inflorescence (28.97 cm) was recorded in P_3 and the lowest length of inflorescence (20.58 cm) was recorded in P_0 (Table 6).

The combined effect of different levels of nitrogen and phosphorus showed a significant effect on the length of inflorescence (Table 7). The highest length of inflorescence (35.04 cm) was observed in $N_2 P_2$ and the lowest length of inflorescence (15.19 cm) was recorded in $N_0 P_0$.

3.5. Number of seeds per inflorescence

Statistically significant differences were found on the number of seeds per inflorescence of spinach due to the application of different nitrogen levels (Table 5). The maximum seeds per inflorescence (58.68) was recorded from N_2 (55.2 kg/ha) and the minimum (38.15) was observed from N_0 . The highest seeds per inflorescence (57.19) were recorded from P_2 and the lowest number of seeds per inflorescence (39.13) in P_0 (Table 6).

The number of seeds per inflorescence was significantly influenced by the combined application of nitrogen and phosphorus (Table 7). The maximum number of seeds per inflorescence (70.41) was recorded from $N_2 P_2$ and the lowest number of seeds per inflorescence (35.33) in $N_0 P_0$.

Table 5. Effect of nitrogen on number of inflorescence per plant, inflorescence length and seeds per inflorescence of spinach

Treatments	Number of inflorescence per plant	Inflorescence length (cm)	Seeds per inflorescence
N_0	5.17	19.99	38.15
N_1	6.30	23.56	47.37
N_2	8.03	29.54	58.86

	N₃	7.13	29.01	56.28
	LSD	0.47	1.857	3.40
	CV %	8.48	8.73	8.12
175	Here,	N ₀ = 0 kg/ha;	N ₁ =27.6 kg/ha;	N ₂ = 55.2 kg/ha;
176				N ₃ = 82.8 kg/ha.

177 **Table 6.** Effect of phosphorus on number of inflorescence per plant, inflorescence length
178 and seeds per inflorescence of spinach

Treatments	Number of inflorescence per plant	Inflorescence length (cm)	Seeds per inflorescence
P ₀	4.94	20.58	39.13
P ₁	6.26	24.12	48.19
P ₂	7.80	28.43	57.19
P ₃	7.64	28.97	56.15
LSD	0.48	1.85	3.39
CV %	8.48	8.73	8.12
Here, P ₀ = 0 kg/ha; P ₁ = 15.84 kg/ha; P ₂ = 31.68 kg/ha; P ₃ = 47.52 kg/ha.			

180 **Table 7.** Combined effect of nitrogen and phosphorus on number of inflorescence per plant,
181 inflorescence length and seeds per inflorescence of spinach

Treatments	Number of inflorescence per plant	Inflorescence length (cm)	Seeds per inflorescence
N ₀ P ₀	3.9	15.19	28.03
N ₀ P ₁	5.03	18.95	35.33
N ₀ P ₂	5.70	21.86	43.82
N ₀ P ₃	6.06	23.96	45.43
N ₁ P ₀	5.08	19.91	36.27
N ₁ P ₁	6.23	22.91	47.79
N ₁ P ₂	6.75	24.72	51.74
N ₁ P ₃	7.15	26.69	53.70
N ₂ P ₀	5.23	22.84	45.25
N ₂ P ₁	7.05	26.15	54.29
N ₂ P ₂	10.53	35.04	70.41
N ₂ P ₃	9.31	34.15	65.48
N ₃ P ₀	5.55	24.38	47.00
N ₃ P ₁	6.73	28.48	55.34
N ₃ P ₂	8.21	32.10	62.80
N ₃ P ₃	8.03	31.10	60.00
LSD	0.94	3.71	6.79
CV (%)	8.48	8.73	8.12

182

183

Here,

N₀= 0 kg/ha;
P₀= 0 kg/ha;

N₁=27.6 kg/ha;
P₁= 15.84 kg/ha;

N₂= 55.2 kg/ha;
P₂= 31.68 kg/ha;

N₃= 82.8 kg/ha.
P₃= 47.52 kg/ha.

185 3.6. Seed yield per hectare (ton)

186 Statistically significant differences were found for seed yield per hectare of spinach due to
187 different nitrogen levels (Table 8). The maximum seed yield (1.10 t ha⁻¹) was recorded from
188 N₂ and the lowest (0.81 t ha⁻¹) was recorded from N₀. The maximum seed yield (1.05 t ha⁻¹)
189 was recorded from P₂ and the lowest (0.84 t ha⁻¹) was recorded from P₀ (Table 9.). The
190 combined effect of nitrogen and phosphorus showed significant variation in the seed yield of
191 spinach (Table 10). The highest seed yield (1.30 t/ha) was recorded from the combination of
192 N₂P₂ and the lowest (0.69 t ha⁻¹) was recorded from N₀P₀ treatment combination.

193
194

3.7. 1000 seed weight (g)

195 Statistically significant differences were found for 1000 seed weight of spinach due to the
196 nitrogen level (Table 8). The maximum 1000 seed weight (10.19 g) was recorded from N₂
197 and the lowest (9.24 g) was recorded from N₀. The maximum 1000 seed weight (10.20 g)
198 was recorded from P₂ and the lowest (9.23 g) was recorded from P₀ (Table 9). The combined
199 effect of nitrogen and phosphorus showed significant variation in 1000 seed weight of
200 spinach (Table 10). The highest 1000 seed weight (11.06 g) was recorded from the
201 combination of N₂P₂ and the lowest (8.77 g) from N₀P₀.

3.8. Germination percentage

203 A significant difference was found on germination % due to fertilizer level (Table 8). The
204 maximum germination percentage (87.33 %) was recorded from N₂ and the minimum (79.33
205 %) was found in N₀. [15] reported that seed yield and its quality of spinach obtained that
206 increasing the rate of nitrogen levels up to the levels at 60 kg N produced higher seed yield
207 with the best quality, germination percentage and germination rate. The maximum
208 germination percentage (87.08 %) was recorded from P₃ and the minimum (79.91 %) was
209 found from P₀ (Table 9).

210 The combined effect of nitrogen and phosphorus was significant on the germination
211 percentage of spinach (Table 10). The highest germination percentage (91.33 %) was
212 recorded from the combination of N₂P₂ and the lowest germination percentage (72.66 %)
213 was recorded in N₀P₀ (control). [14] experimented on the farm of the Department of
214 Horticulture, BSMRAU, Salna, Gazipur on 6 spinach genotypes to observe their seed
215 production potentiality and to evaluate the quality of produced seed. They reported that the
216 quality test of seed was done based on germination test (%), seed vigor test (Electrical
217 conductivity), moisture test (%) and thousand seed weight (g) of seeds.

3.9. Electrical conductivity test

219 A significant difference was found in electrical conductivity test value due to the application
220 of different levels of nitrogen (Table 8). The highest EC test value (13.87 dS/cm) was
221 recorded from treatment N₀ whereas, the minimum (11.16 dS/cm) was found in N₂. The
222 maximum EC test value (13.35 dS/cm) was recorded from treatment P₀ and the lowest EC
223 test value (11.79 dS/cm) was recorded in P₂ (Table 9). The combined effect of nitrogen and
224 phosphorus was significant on the EC test value of spinach (Table 10). The highest EC test
225 value (14.83 dS/cm) was recorded from the combination (N₀P₀) and the lowest EC test value
226 (10.20 dS/cm) was recorded with N₂P₂ treatment combination. Why highest in N₂P₂ and not
227 N₃P₃
228

229 **Table 8. Effect of nitrogen on seed yield per hectare 1000 seed weight, germination**
230 **percentage and electrical conductivity test**

Treatments	Seed yield per hectare (t)	1000 seed weight (g)	Germination (%)	Electrical conductivity test (ds/cm)
N ₀	0.81	9.24	79.33	13.87
N ₁	0.91	9.58	83.41	12.78
N ₂	1.10	10.19	87.33	11.16
N ₃	1.01	10.14	87.25	11.73
LSD	0.06	0.75	4.95	0.85

CV (%) 8.28 9.19 7.05 8.25

Table 9. Effect of phosphorus on seed yield per hectare, 1000 seed weight, germination percentage and electrical conductivity test

Treatments	Seed yield per hectare (t)	1000 seed weight (g)	Germination (%)	Electrical conductivity test (ds/cm)
P ₀	0.84 c	9.23 b	79.91	13.35
P ₁	0.93 b	9.56 ab	83.75	12.57
P ₂	1.05 a	10.20 a	86.58	11.79
P ₃	1.01 a	10.18 a	87.08	11.81
LSD	0.067	0.75	4.95	0.85
CV (%)	8.28	9.19	7.05	8.25

Here, P₀= 0 kg/ha; P₁= 15.84 kg/ha; P₂= 31.68 kg/ha; P₃= 47.52 kg/ha.

Table 10. The combined effect of nitrogen and phosphorus seed yield per hectare, 1000 seed weight, germination percentage and electrical conductivity test

Treatments	Seed yield per hectare (t)	1000 seed weight (g)	Germination (%)	Electrical conductivity test (ds/cm)
N ₀ P ₀	0.69	8.77	72.66	14.83
N ₀ P ₁	0.83	9.22	80.00	13.97
N ₀ P ₂	0.85	9.43	81.00	13.37
N ₀ P ₃	0.88	9.56	83.66	13.30
N ₁ P ₀	0.83	9.15	79.00	13.55
N ₁ P ₁	0.91	9.47	84.00	12.85
N ₁ P ₂	0.95	9.76	84.66	12.58
N ₁ P ₃	0.96	9.95	86.00	12.14
N ₂ P ₀	0.93	9.26	83.33	12.59
N ₂ P ₁	0.99	9.74	84.66	11.46
N ₂ P ₂	1.30	11.06	91.33	10.20
N ₂ P ₃	1.16	10.70	90.00	10.42
N ₃ P ₀	0.92	9.73	84.66	12.45
N ₃ P ₁	0.98	9.80	86.33	12.01
N ₃ P ₂	1.09	10.53	89.33	11.03
N ₃ P ₃	1.04	10.50	88.66	11.41
LSD	0.13	1.50	9.91	1.70
CV (%)	8.28	9.19	7.05	8.25

Here, N₀= 0 kg/ha; N₁=27.6 kg/ha; N₂= 55.2 kg/ha; N₃= 82.8 kg/ha.
P₀= 0 kg/ha; P₁= 15.84 kg/ha; P₂= 31.68 kg/ha; P₃= 47.52 kg/ha.

3.10. Benefit cost ratio (BCR)

The combination of nitrogen and phosphorus showed different benefit cost ratio. The highest benefit cost ratio (1.59) which is economically more benefitted was obtained from N₂P₂ and the lowest benefit cost ratio (1.01) which is economically less benefitted was obtained from the treatment combination of N₀P₀(Table.11).

Table 11. Cost and return analysis of spinach seed considering nitrogen and phosphorus

Treatments	Seed yield (t/ha)	Gross return (Tk/ha)	Total cost of production(Tk)	Net return (Tk/ha)	Benefit cost ratio (BCR)
N ₀ P ₀	0.69	117000	116656.8	13156	1.01
N ₀ P ₁	0.83	124500	118342	6158	1.05
N ₀ P ₂	0.85	127500	120027.3	7472.7	1.06
N ₀ P ₃	0.88	132000	121712.5	10287.5	1.08
N ₁ P ₀	0.83	124500	117870.1	6629.9	1.05
N ₁ P ₁	0.91	136500	119555.4	16944.6	1.14
N ₁ P ₂	0.95	142500	121240.6	21259.4	1.17
N ₁ P ₃	0.96	144000	122925.9	21074.1	1.17
N ₂ P ₀	0.93	133500	119083.5	14416.5	1.12
N ₂ P ₁	0.99	148500	120768.8	27731.2	1.22
N ₂ P ₂	1.30	195000	122454	72546	1.59
N ₂ P ₃	1.16	174000	124139.3	49860.7	1.40
N ₃ P ₀	0.92	138000	120296.9	17703.1	1.14
N ₃ P ₁	0.98	147000	121982.1	25017.9	1.20
N ₃ P ₂	1.09	163500	123667.4	39832.6	1.32
N ₃ P ₃	1.04	156000	125352.6	30647.4	1.24

248 Here,

- 249 ➤ Sale of marketable seed @ Tk.150/kg
250 ➤ Gross return = Marketable yield x Tk./kg
251 ➤ Net income = Gross return-Total cost of production
252 ➤ BCR = Gross return ÷ cost of production

253

254 4. CONCLUSION

255

256 Both seed yield and economic benefit of crop are important for the seed production.
257 Application of nitrogen (55.2 kg/ha) represents higher seed yield in spinach than without no
258 nitrogen and (82.8 kg/ha) nitrogen. According to the results of the present experiment it may
259 be concluded that efficient production of spinach seed is increased by the application of
260 phosphorus (31.68 kg/ha). The combined effect of nitrogen and phosphorus had positive
261 effect on morphological characters, yield contributing characters but seed yield and seed
262 quality of spinach increased by the combined application of (N 55.2 kg/ha+ P 31.68 kg/ha).
263 On the basis of benefit cost ratio, it may be suggested that application of N 55.2 kg/ha with P
264 31.68 kg/ha combination seemed to be more suitable for getting higher seed yield in
265 spinach.

266

267

268 COMPETING INTERESTS

269

270 The authors have declared that no competing interests exist.

271

272 REFERENCES

273

- 274 1. George, R.A.T. (1985). Vegetable seed production. New York, America. p-318.

- 275 2. BBS. (2016). Year book of Agricultural statistics of Bangladesh.2016 28th ed.
 276 Bangladesh Bureau of statistics. Statistics Division, Ministry of planning. Govt of the
 277 people Republic of Bangladesh.p-204.
- 278 3. Anonymous. (1995). Requirement and supply of seeds. Report of a Task Force,
 279 Ministry of Agriculture, Government of the Peoples Republic of Bangladesh, Dhaka.
 280 p-284.
- 281 4. Milford, G.F.J., Armstrong, M.J., Jarvis, P.J., Houghton, B.J., Bellett, D.M., Jones, J.
 282 and Leigh, R.A. (2000). Effects of potassium fertilizer on the yield, quality and
 283 potassium offtake of sugar beet crops grown on soils of different potassium status.
 284 *J. Agric. Sci.* **135**: 1-10.
- 285 5. Eckert, D. (2010). Efficient Fertilizer Use of Nitrogen. John Willy and Sons, New
 286 York, America. pp. 1-19.
- 287 6. Griffith, B. (2010) Efficient Fertilizer Use - Phosphorus. McMillan Co. Amsterdam,
 288 Netherland. pp. 1-7.
- 289 7. Cantliffe, D. (1992). Nitrate accumulation in vegetable crops as affected by
 290 photoperiod and light duration (beets, radish, spinach, beans). *J. Am. Soc. Hort.*
 291 *Sci.*, **97**: 414-418.
- 292 8. Jaworska, G. (2005) Content of nitrates, nitrites, and oxalates in New Zealand
 293 spinach. *Food Chem.* **89**: 235-242.
- 294 9. Canali, S., Montemurro, F., Tittarelli, F., and Masetti, O. (2008).Effect of nitrogen
 295 fertilization reduction on yield, quality and N utilization of processing spinach. *J.*
 296 *Food. Agric. Environ.* **6**(3&4): 242-247.
- 297 10. Wang, Z. H., Zong, Z.Q., Li, S. X., and Chen, B. M. (2002). Nitrate accumulation in
 298 vegetables and its residual in vegetable fields. *Environ. Sci.* **23**(11): 79-83.
- 299 11. Fadaly, K.A., and Mishriky, J.F. (2000). Effect of nitrogen sources and levels on
 300 growth, yield and mineral composition of spinach. Bulletin of Faculty of Agriculture,
 301 University of Cairo. **41**(3): 973-988.
- 302 12. Assiouty, M.M., Sedera, S.A. and Giza, D. (2005). Effect of bio and chemical
 303 fertilizers on seed production and quality of spinach. *Int. J. Agril.Biol.* **7**(6): 947-952.
- 304 13. Boroujerdnia, M. and Ansari, N.A. (2007).Effect of Different Levels of Nitrogen
 305 Fertilizer and Cultivars on Growth, Yield and Yield Components of Romaine Lettuce
 306 (*Lactuca sativa* L.). *Middle Eastern and Russian J. Plant. Sci. Biotech.* **1**(2): 47-53.
- 307 14. Ambia, K., Hossain, M., Haque, M., Mosiur, A., Halder, C. and Krisna, S. (2016).
 308 Evaluation Seed Yield and Quality of Different Genotypes of Spinach (*Beta*
 309 *vulgaris*). Evaluation of seed yield of different genotypes of spinach. *Sci. Agri.* **16** (2):
 310 61-66.
- 311 15. Zaki, M.S. and Halawa, S.S. (2016). Effect of nitrogen fertilizer and growth
 312 stimulants on flowering behavior, seed yield and its quality of spinach. *Annal. Agric.*
 313 *Sci.* **54**(2): 345-354.