

Effect of Nitrogen and Phosphorus on the Growth and Seed Yield of Spinach

ABSTRACT

The experiment was conducted at the farm of Sher-e-Bangla Agricultural University, Dhaka, Bangladesh during Rabi (November 2017 to March 2018) season to find out the growth, yield and economic benefit of spinach seed as influenced by nitrogen and phosphorus. The research involved two factors. Factor A: Four levels of nitrogen; $N_0 = 0$ kg/ha, $N_1 = 27.6$ kg/ha, $N_2 = 55.2$ kg/ha, $N_3 = 82.8$ kg/ha, and factor B: Four levels of phosphorus; $P_0 = 0$ kg/ha, $P_1 = 15.84$ kg/ha, $P_2 = 31.68$ kg/ha, $P_3 = 47.52$ kg/ha. There were 16 treatment combinations in the experiment and laid out in Randomized Complete Block Design (RCBD) with three replications. Quality tests of seeds were done based on the germination test (%), seed vigor test (Electrical conductivity). In case of nitrogen, the highest seed yield (1.10 t ha^{-1}), germination percentage (87.33 %) and lowest value in EC test (11.87 dS/cm) were obtained from N_2 , while the lowest seed yield (0.81 t ha^{-1}), germination percentage (79.33 %) and highest value in EC test (13.87 dS/cm) from N_0 . For phosphorus levels, the highest seed yield (1.05 t ha^{-1}), germination percentage (86.58 %) and lowest value in EC test (11.79 dS/cm) were recorded from P_2 , whereas the lowest seed yield (0.84 t ha^{-1}), germination percentage (79.91 %) and highest value in EC test (13.35 dS/cm) from P_0 . Due to mutual effect, the highest seed yield (1.30 t ha^{-1}), germination percentage (91.33 %) and lowest value in EC test (10.2 dS/cm) were noted from N_2P_2 , whereas the lowest seed yield (0.69 t ha^{-1}), germination percentage (72.66 %) and highest value in EC test (14.83 dS/cm) from N_0P_0 . From the economic point of view, the highest Benefit-Cost Ratio (BCR) (1.59) was found in the treatment of N_2P_2 and the lowest BCR (1.01) was found in the treatment of N_0P_0 . It is apparent that the treatment combination N_2P_2 gave the best performance for the seed yield and economic benefit of spinach.

Keywords: Growth, Nitrogen, Phosphorus, Seed Yield

1. INTRODUCTION

Spinach (*Spinacia oleracea*) is a leafy green cool-season vegetable that is known for its nutritive value and is considered one of the most popular vegetables in Bangladesh. It is believed to have originated from Persia. Its leaves are a common edible vegetable. By weight, spinach consists of 91.4% water, 3.6% carbons, and 2.9% protein [1]. There are 23 calories in 100 grams of spinach and further benefits of spinach leaves consumption can be of more value e.g vital mineral elements in spinach compared to other vegetables e.g iron which is very good for women. The seed is produced for commercial consumption and for seed companies that supply seed throughout the country. In Bangladesh spinach occupies 22000 acres [2] with an annual production of 66000 tons. In comparison to other countries, this yields much lower. So, to use of quality seeds of high yield varieties are the foremost important technique for maximizing yield per unit area. Quality seed can increase vegetable production by up to 25-50% [1]. Farmers save seeds annually of about 50 tons and used

every year, which in most cases are of inferior in quality [3]. Appropriate fertilizer application to the plants greatly affects their growth and production. Nitrogen strongly stimulates growth, expansion of the crop canopy and interception of solar radiation [4]. Increasing the levels of nitrogen during the vegetative stage can strengthen and allow a plant to grow more rapidly and produce large amounts of succulent, green foliage, which in turn can generate higher yields [5]. Similarly, phosphorus (P) is an essential nutrient act as catalysts in the conversion of numerous key biochemical reactions in plants. Phosphorus stimulates root development, improves flower formation, seed production and improves crop quality and increases resistance to plant diseases [6]. Leafy vegetables, particularly, spinach is highly responsive to fertilization [7] and oxalates are the main indexes of the quality due to a very efficient uptake system and inefficient reductive systems [8]. The fertilizer requirements on sandy and sandy loams are 85 to 120 kg N, 75 to 85 kg P₂O₅, and 85 to 150 kg K₂O. On heavier clay soils, 75 kg ha⁻¹ of each nutrient should be adequate. If the fertilizer is banded at seeding it should be placed along each side of the rows 2 to 3 inches below the level of the seed and 6 inches to the side of the row; fertilizer should never come in contact with the seed and two or three splits of 85 to 120 kg ha⁻¹ N would be adequate as side-dressing [9]. An adequate supply of fertilizers can promote plant growth and increase crop production, but excessive and inappropriate use of chemical fertilizers causes accumulation of compounds in the edible products which can be detrimental to human health in addition cause environmental pollution and economic losses [10]. [11] found that N increased the spinach yield and enhanced the accumulation of N and P in leaves. [12] reported that application of 40 kg N + 15.0 kg P₂O₅ increased plant fresh yield by 27.2 and 42.3% and 16.3 and 10.4% in seed yield in relation to the control in the first and second seasons, respectively. [13] achieved the highest yield with 120 kg N ha⁻¹ Farmers in Bangladesh generally do not use any improved or special techniques for quality seed production. There for to achieve higher seeds yield with good quality fertilizer management is an important practice which cannot be emphasized.

2. MATERIAL AND METHODS

2.1. Experimental Site

The experiment was conducted at the Agronomy Research Farm of Sher-e-Bangla Agricultural University (SAU), Dhaka, Bangladesh from November 2017 to March 2018. Experimental site situated an elevation of 8 meters above the sea level in the Agro-ecological zone of "Madhupur Tract" (AEZ-28). The soil was sandy loam and medium high land in texture having pH 5.46- 5.62.

2.2 Experiment Frame Work

The research was consisted of two factors: Factor A: Four levels of nitrogen; N₀= 0 kg/ha N₁= 27.6 kg/ha, N₂=55.2 kg/ha, N₃= 82.8 kg/ha, and factor B: Four levels of phosphorus; P₀= 0 kg/ha; P₁ = 15.84 kg/ha, P₂= 31.68 kg/ha, P₃= 47.52 kg/ha. The two factors experiment was laid out following the Randomized Complete Block Design (RCBD) with three replications. The experiment was divided into three equal blocks where each block was divided into 16 plots. Then 16 treatment combinations were allotted randomly in each block. The size of each unit plot was 1.5 m × 1 m. The distance maintained between two blocks and two plots were 0.75 m and 0.5 m, respectively. Row to row distance was 30 cm and plant to plant distance was 20 cm.

2.3. Application of manure and fertilizers

About 5 t ha⁻¹ well-decomposed cow dung was applied as the control (N₀P₀) treatment) plot and incorporated adequately to the soil during final land preparation whereas other plots were applied with inorganic fertilizer as per treatment. Doses of inorganic fertilizers (Urea and Triple superphosphate) were applied in the experimental plot according to the treatments (Table 1). The whole amount of TSP and half the amount of urea and MoP (180

kg/ha) were also applied as basal dose before sowing of seed in the main field. 1st top dressing of urea was applied when seedlings established in the main field about 10 days after seed sowing. 2nd top dressing of urea and the rest amount of MoP was applied about 25 days after 1st top dressing. Then the rest amount of urea was applied as 3rd installment after flowering. Each top dressing was followed by manual irrigation.

Table 1. Doses of nutrients application in the main field as per treatment

Treatments	N rates (kg ha ⁻¹)	Fertilizers (kg ha ⁻¹)	Doses (g plot ⁻¹)	Treatments	P rates (kg ha ⁻¹)	Fertilizers (kg ha ⁻¹)	Doses (g plot ⁻¹)
	N	Urea	Urea		P	TSP	TSP
N ₀	0	0	0	P ₀	0	0	0
N ₁	27.6	60	9	P ₁	15.84	75	11.25
N ₂	55.2	120	18	P ₂	31.68	150	22.5
N ₃	82.8	180	27	P ₃	47.52	225	33.75

2.4. Economic analysis

The cost of production was analyzed in order to find out the most economic treatment of nitrogen and phosphorus for quality seed production of spinach. All the non-material and material input costs and interests in running capital were considered for computing the cost of production. The benefit-cost ratio (BCR) was calculated by the following formula:

Benefit-cost ratio (BCR) = Gross return (tk/ha) ÷ Total cost of production (tk/ha)

2.5. Statistical analysis

The data obtained for different characters were statistically analyzed to observe the significant difference among the treatment by using the STATISTIX-10 computer package program. The mean values of all the characters were calculated and analysis of variance was performed. The significance of the difference among the treatments means was estimated by the Least Significant Different Test (LSD) at 5% level of probability.

3. RESULTS AND DISCUSSION

3.1 Plant height (cm)

A significant variation was observed on the plant height of spinach due to the application of different levels of nitrogen (Table 2.). The highest plant height 13.94cm and 47.69cm were obtained before flowering and at the time of harvest respectively from N₃ while the lowest plant height 11.01cm and 34.88cm were obtained before flowering and at the time of harvest respectively from N₀. P₃ showed the highest plant height 13.64cm and 46.68cm before flowering and at the time of harvest respectively and the lowest plant height 11.60 cm and 35.18 cm at before flowering and at the time of harvest respectively in P₀ (Table 3).

The combined effect of different levels of nitrogen and phosphorus application showed a significant effect on the plant height of spinach (Table 4). The highest plant height 15.21cm and 55.25cm before flowering and at the time of harvest, respectively was observed in N₃P₃ while the lowest plant height 10.10cm and 27.7cm before flowering and at the time of harvest, respectively in N₀P₀.

3.2. Number of leaves per plant

The effect of nitrogen on the number of leaves per plant of spinach was significant (Table 2). The highest number of leaves (8.46) was produced from N₂ and the lowest number of leaves

121 (6.42) was observed in N_0 . P_2 showed the maximum leaves per plant (8.30) and the
 122 minimum leaves per plant (6.36) was observed in P_0 (Table 3).

123 The combined effect of different levels of nitrogen and phosphorus showed a significant
 124 effect on the number of leaves per plant of spinach (Table 4). The maximum leaves per plant
 125 (10.35) were observed in N_2P_2 and the lowest leaves per plant (5.01) were recorded with
 126 N_0P_0 , [14] also found a similar result.
 127

128 **Table 2.** Effect of nitrogen on plant height (before flowering and at the time of harvest)
 129 leaves per plant of spinach

Treatments	Plant height before flowering (cm)	Plant height at the time of harvest (cm)	Leaves per plant
N_0	11.01	34.88	6.42
N_1	12.34	40.35	7.24
N_2	13.07	43.75	8.46
N_3	13.94	47.69	7.84
LSD	0.87	3.12	0.54
CV %	8.34	9.00	8.70

130 Here, $N_0=0$ kg/ha; $N_1=27.6$ kg/ha; $N_2=55.2$ kg/ha; $N_3=82.8$ kg/ha

131 **Table 3.** Effect of phosphorus on plant height (before and at the time of harvest), Leaves per
 132 plant of spinach

Treatments	Plant height before flowering	Plant height at the time of harvest	Leaves per plant
P_0	11.60	35.18	6.36
P_1	12.16	40.37	7.24
P_2	12.96	44.44	8.30
P_3	13.64	46.68	8.06
LSD	0.87	3.11	0.543
CV %	8.34	9.00	8.70

133 Here, $P_0=0$ kg/ha; $P_1=15.84$ kg/ha; $P_2=31.68$ kg/ha; $P_3=47.52$ kg/ha.

134 **Table 4.** Combined effect of nitrogen and phosphorus on the plant height (before and at the
 135 time of harvesting), leaves per plant of spinach

Treatments	Plant height before flowering (cm)	Plant height at the time of harvest (cm)	Leaves per plant
N_0P_0	10.10	27.7	5.01
N_0P_1	10.63	34.34	6.25
N_0P_2	11.25	37.31	6.66
N_0P_3	12.07	40.18	7.76
N_1P_0	11.49	34.65	6.60
N_1P_1	12.15	41.03	7.33
N_1P_2	12.60	42.44	7.46
N_1P_3	13.12	43.29	7.58
N_2P_0	12.14	37.88	6.91
N_2P_1	12.54	21.80	7.81
N_2P_2	13.42	47.31	10.35
N_2P_3	14.17	48.01	8.76
N_3P_0	12.65	40.51	6.93
N_3P_1	13.35	44.30	7.56

$N_3 P_2$	14.56	50.71	8.73
$N_3 P_3$	15.21	55.25	8.13
LSD	1.75	6.25	1.08
CV %	8.34	9.00	8.70

Here, N_0 = 0 kg/ha; N_1 =27.6 kg/ha; N_2 = 55.2 kg/ha; N_3 = 82.8 kg/ha.
 P_0 = 0 kg/ha; P_1 = 15.84 kg/ha; P_2 = 31.68 kg/ha; P_3 = 47.52 kg/ha.

3.3. Number of inflorescence per plant

Statistically significant differences were found on the number of inflorescence per plant of spinach due to the application of nitrogen (Table 5). The highest number of inflorescence per plant (8.03) was recorded from N_2 whereas, the lowest number (5.17) was observed from N_0 . The highest number of inflorescence per plant (7.80) was recorded from P_2 and the lowest (4.94) was found from P_0 (Table 6).

The combined effect of nitrogen and phosphorus showed significant variation in the number of inflorescences per plant (Table 7). The highest number of inflorescence per plant (10.53) was recorded from $N_2 P_2$ and the lowest number of inflorescence per plant (3.90) from $N_0 P_0$.

3.4. Length of inflorescence

A significant variation was observed on the length of an inflorescence of spinach when different levels of nitrogen were applied (Table 5). The highest length of inflorescence (29.54cm) was recorded in N_2 and the lowest length of inflorescence (19.09 cm) from N_0 . The highest length of inflorescence (28.97 cm) was recorded in P_3 and the lowest length of inflorescence (20.58 cm) was recorded in P_0 (Table 6).

The combined effect of different levels of nitrogen and phosphorus showed a significant effect on the length of inflorescence (Table 7). The highest length of inflorescence (35.04 cm) was observed in $N_2 P_2$ and the lowest length of inflorescence (15.19 cm) was recorded in $N_0 P_0$.

3.5. Number of seeds per inflorescence

Statistically significant differences were found on the number of seeds per inflorescence of spinach due to the application of different nitrogen levels (Table 5). The maximum seeds per inflorescence (58.68) was recorded from N_2 (55.2 kg/ha) and the minimum (38.15) was observed from N_0 . The highest seeds per inflorescence (57.19) were recorded from P_2 and the lowest number of seeds per inflorescence (39.13) in P_0 (Table 6).

The number of seeds per inflorescence was significantly influenced by the combined application of nitrogen and phosphorus (Table 7). The maximum number of seeds per inflorescence (70.41) was recorded from $N_2 P_2$ and the lowest number of seeds per inflorescence (35.33) in $N_0 P_0$.

Table 5. Effect of nitrogen on number of inflorescence per plant, inflorescence length and seeds per inflorescence of spinach

Treatments	Number of inflorescence per plant	Inflorescence length (cm)	Seeds per inflorescence
N_0	5.17	19.99	38.15
N_1	6.30	23.56	47.37
N_2	8.03	29.54	58.86
N_3	7.13	29.01	56.28
LSD	0.47	1.857	3.40

	CV %	8.48	8.73	8.12
173	Here,	N ₀ = 0 kg/ha;	N ₁ =27.6 kg/ha;	N ₂ = 55.2 kg/ha; N ₃ = 82.8 kg/ha.
174				

175 **Table 6.** Effect of phosphorus on number of inflorescence per plant, inflorescence length
176 and seeds per inflorescence of spinach

Treatments	Number of inflorescence per plant	Inflorescence length (cm)	Seeds per inflorescence
P ₀	4.94	20.58	39.13
P ₁	6.26	24.12	48.19
P ₂	7.80	28.43	57.19
P ₃	7.64	28.97	56.15
LSD	0.48	1.85	3.39
CV %	8.48	8.73	8.12

177 Here, P₀= 0 kg/ha; P₁= 15.84 kg/ha; P₂= 31.68 kg/ha; P₃= 47.52 kg/ha.

178 **Table 7.** Combined effect of nitrogen and phosphorus on number of inflorescence per plant,
179 inflorescence length and seeds per inflorescence of spinach

Treatments	Number of inflorescence per plant	Inflorescence length (cm)	Seeds per inflorescence
N ₀ P ₀	3.9	15.19	28.03
N ₀ P ₁	5.03	18.95	35.33
N ₀ P ₂	5.70	21.86	43.82
N ₀ P ₃	6.06	23.96	45.43
N ₁ P ₀	5.08	19.91	36.27
N ₁ P ₁	6.23	22.91	47.79
N ₁ P ₂	6.75	24.72	51.74
N ₁ P ₃	7.15	26.69	53.70
N ₂ P ₀	5.23	22.84	45.25
N ₂ P ₁	7.05	26.15	54.29
N ₂ P ₂	10.53	35.04	70.41
N ₂ P ₃	9.31	34.15	65.48
N ₃ P ₀	5.55	24.38	47.00
N ₃ P ₁	6.73	28.48	55.34
N ₃ P ₂	8.21	32.10	62.80
N ₃ P ₃	8.03	31.10	60.00
LSD	0.94	3.71	6.79
CV (%)	8.48	8.73	8.12

180	Here,	N ₀ = 0 kg/ha;	N ₁ =27.6 kg/ha;	N ₂ = 55.2 kg/ha;	N ₃ = 82.8 kg/ha.
181		P ₀ = 0 kg/ha;	P ₁ = 15.84 kg/ha;	P ₂ = 31.68 kg/ha;	P ₃ = 47.52 kg/ha.
182					

183 3.6. Seed yield per hectare (ton)

184 Statistically significant differences were found for seed yield per hectare of spinach due to
185 different nitrogen levels (Table 8). The maximum seed yield (1.10 t ha⁻¹) was recorded from
186 N₂ and the lowest (0.81 t ha⁻¹) was recorded from N₀. The maximum seed yield (1.05 t ha⁻¹)
187 was recorded from P₂ and the lowest (0.84 t ha⁻¹) was recorded from P₀ (Table 9.). The
188 combined effect of nitrogen and phosphorus showed significant variation in the seed yield of
189 spinach (Table 10). The highest seed yield (1.30 t/ha) was recorded from the combination of
190 N₂P₂ and the lowest (0.69 t ha⁻¹) was recorded from N₀P₀ treatment combination.

192 3.7. 1000 seed weight (g)

Statistically significant differences were found for 1000 seed weight of spinach due to the nitrogen level (Table 8). The maximum 1000 seed weight (10.19 g) was recorded from N₂ and the lowest (9.24 g) was recorded from N₀. The maximum 1000 seed weight (10.20 g) was recorded from P₂ and the lowest (9.23 g) was recorded from P₀ (Table 9). The combined effect of nitrogen and phosphorus showed significant variation in 1000 seed weight of spinach (Table 10). The highest 1000 seed weight (11.06 g) was recorded from the combination of N₂P₂ and the lowest (8.77 g) from N₀P₀.

3.8. Germination percentage

A significant difference was found on germination % due to fertilizer level (Table 8). The maximum germination percentage (87.33 %) was recorded from N₂ and the minimum (79.33 %) was found in N₀. [15] reported that seed yield and its quality of spinach obtained that increasing the rate of nitrogen levels up to the levels at 60 kg N produced higher seed yield with the best quality, germination percentage and germination rate. The maximum germination percentage (87.08 %) was recorded from P₃ and the minimum (79.91 %) was found from P₀ (Table 9).

The combined effect of nitrogen and phosphorus was significant on the germination percentage of spinach (Table 10). The highest germination percentage (91.33 %) was recorded from the combination of N₂P₂ and the lowest germination percentage (72.66 %) was recorded in N₀P₀ (control). [14] experimented on the farm of the Department of Horticulture, BSMRAU, Salna, Gazipur on 6 spinach genotypes to observe their seed production potentiality and to evaluate the quality of produced seed. They reported that the quality test of seed was done based on germination test (%), seed vigor test (Electrical conductivity), moisture test (%) and thousand seed weight (g) of seeds.

3.9. Electrical conductivity test

A significant difference was found in electrical conductivity test value due to the application of different levels of nitrogen (Table 8). The highest EC test value (13.87 dS/cm) was recorded from treatment N₀ whereas, the minimum (11.16 dS/cm) was found in N₂. The maximum EC test value (13.35 dS/cm) was recorded from treatment P₀ and the lowest EC test value (11.79 dS/cm) was recorded in P₂ (Table 9). The combined effect of nitrogen and phosphorus was significant on the EC test value of spinach (Table 10). The highest EC test value (14.83 dS/cm) was recorded from the combination (N₀P₀) and the lowest EC test value (10.20 dS/cm) was recorded with N₂P₂ treatment combination. Why highest in N₂P₂ and not N₃P₃

Table 8. Effect of nitrogen on seed yield per hectare 1000 seed weight, germination percentage and electrical conductivity test

Treatments	Seed yield per hectare (t)	1000 seed weight (g)	Germination (%)	Electrical conductivity test (ds/cm)
N ₀	0.81	9.24	79.33	13.87
N ₁	0.91	9.58	83.41	12.78
N ₂	1.10	10.19	87.33	11.16
N ₃	1.01	10.14	87.25	11.73
LSD	0.06	0.75	4.95	0.85
CV (%)	8.28	9.19	7.05	8.25

230 **Table 9.** Effect of phosphorus on seed yield per hectare, 1000 seed weight, germination
 231 percentage and electrical conductivity test

Treatments	Seed yield per hectare (t)	1000 seed weight (g)	Germination (%)	Electrical conductivity test (ds/cm)
P ₀	0.84 c	9.23 b	79.91	13.35
P ₁	0.93 b	9.56 ab	83.75	12.57
P ₂	1.05 a	10.20 a	86.58	11.79
P ₃	1.01 a	10.18 a	87.08	11.81
LSD	0.067	0.75	4.95	0.85
CV (%)	8.28	9.19	7.05	8.25

232 Here, P₀ = 0 kg/ha; P₁ = 15.84 kg/ha; P₂ = 31.68 kg/ha; P₃ = 47.52 kg/ha.

233 **Table 10.** The combined effect of nitrogen and phosphorus seed yield per hectare, 1000
 234 seed weight, germination percentage and electrical conductivity test

Treatments	Seed yield per hectare (t)	1000 seed weight (g)	Germination (%)	Electrical conductivity test (ds/cm)
N ₀ P ₀	0.69	8.77	72.66	14.83
N ₀ P ₁	0.83	9.22	80.00	13.97
N ₀ P ₂	0.85	9.43	81.00	13.37
N ₀ P ₃	0.88	9.56	83.66	13.30
N ₁ P ₀	0.83	9.15	79.00	13.55
N ₁ P ₁	0.91	9.47	84.00	12.85
N ₁ P ₂	0.95	9.76	84.66	12.58
N ₁ P ₃	0.96	9.95	86.00	12.14
N ₂ P ₀	0.93	9.26	83.33	12.59
N ₂ P ₁	0.99	9.74	84.66	11.46
N ₂ P ₂	1.30	11.06	91.33	10.20
N ₂ P ₃	1.16	10.70	90.00	10.42
N ₃ P ₀	0.92	9.73	84.66	12.45
N ₃ P ₁	0.98	9.80	86.33	12.01
N ₃ P ₂	1.09	10.53	89.33	11.03
N ₃ P ₃	1.04	10.50	88.66	11.41
LSD	0.13	1.50	9.91	1.70
CV (%)	8.28	9.19	7.05	8.25

235 Here, N₀ = 0 kg/ha; N₁ = 27.6 kg/ha; N₂ = 55.2 kg/ha; N₃ = 82.8 kg/ha.
 236 P₀ = 0 kg/ha; P₁ = 15.84 kg/ha; P₂ = 31.68 kg/ha; P₃ = 47.52 kg/ha.

237 **3.10. Benefit cost ratio (BCR)**

239
 240 The combination of nitrogen and phosphorus showed different benefit cost ratio. The highest
 241 benefit cost ratio (1.59) which is economically more benefitted was obtained from N₂P₂ and
 242 the lowest benefit cost ratio (1.01) which is economically less benefitted was obtained from
 243 the treatment combination of N₀P₀ (Table.11).

244 **Table 11.** Cost and return analysis of spinach seed considering nitrogen and phosphorus
 245

Treatments	Seed yield (t/ha)	Gross return (Tk/ha)	Total cost of production(Tk)	Net return (Tk/ha)	Benefit cost ratio (BCR)
N ₀ P ₀	0.69	117000	116656.8	13156	1.01

$N_0 P_1$	0.83	124500	118342	6158	1.05
$N_0 P_2$	0.85	127500	120027.3	7472.7	1.06
$N_0 P_3$	0.88	132000	121712.5	10287.5	1.08
$N_1 P_0$	0.83	124500	117870.1	6629.9	1.05
$N_1 P_1$	0.91	136500	119555.4	16944.6	1.14
$N_1 P_2$	0.95	142500	121240.6	21259.4	1.17
$N_1 P_3$	0.96	144000	122925.9	21074.1	1.17
$N_2 P_0$	0.93	133500	119083.5	14416.5	1.12
$N_2 P_1$	0.99	148500	120768.8	27731.2	1.22
$N_2 P_2$	1.30	195000	122454	72546	1.59
$N_2 P_3$	1.16	174000	124139.3	49860.7	1.40
$N_3 P_0$	0.92	138000	120296.9	17703.1	1.14
$N_3 P_1$	0.98	147000	121982.1	25017.9	1.20
$N_3 P_2$	1.09	163500	123667.4	39832.6	1.32
$N_3 P_3$	1.04	156000	125352.6	30647.4	1.24

Here,

- Sale of marketable seed @ Tk.150/kg
- Gross return = Marketable yield x Tk./kg
- Net income = Gross return-Total cost of production
- BCR = Gross return ÷ cost of production

4. CONCLUSION

Both seed yield and economic benefit of crop are important for the seed production. Application of nitrogen (55.2 kg/ha) represents higher seed yield in spinach than without no nitrogen and (82.8 kg/ha) nitrogen. According to the results of the present experiment it may be concluded that efficient production of spinach seed is increased by the application of phosphorus (31.68 kg/ha). The combined effect of nitrogen and phosphorus had positive effect on morphological characters, yield contributing characters but seed yield and seed quality of spinach increased by the combined application of (N 55.2 kg/ha+ P 31.68 kg/ha). On the basis of benefit cost ratio, it may be suggested that application of N 55.2 kg/ha with P 31.68 kg/ha combination seemed to be more suitable for the highest values of gross return, net return as well as highest beneficial cost ratio of spinach..

COMPETING INTERESTS

The authors have declared that no competing interests exist.

REFERENCES

1. George, R.A.T. (1985). Vegetable seed production. New York, America. p-318.
2. BBS. (2016). Year book of Agricultural statistics of Bangladesh.2016 28th ed. Bangladesh Bureau of statistics. Statistics Division, Ministry of planning. Govt of the people Republic of Bangladesh.p-204.
3. Anonymous. (1995). Requirement and supply of seeds. Report of a Task Force, Ministry of Agriculture, Government of the Peoples Republic of Bangladesh, Dhaka. p-284.
4. Milford, G.F.J., Armstrong, M.J., Jarvis, P.J., Houghton, B.J., Bellett, D.M., Jones, J. and Leigh, R.A. (2000). Effects of potassium fertilizer on the yield, quality and

- 281 potassium offtake of sugar beet crops grown on soils of different potassium status.
282 *J. Agric. Sci.* **135**: 1-10.
- 283 5. Eckert, D. (2010). Efficient Fertilizer Use of Nitrogen. John Willy and Sons, New
284 York, America. pp. 1-19.
- 285 6. Griffith, B. (2010) Efficient Fertilizer Use - Phosphorus. McMillan Co. Amsterdam,
286 Netherland. pp. 1-7.
- 287 7. Cantliffe, D. (1992). Nitrate accumulation in vegetable crops as affected by
288 photoperiod and light duration (beets, radish, spinach, beans). *J. Am. Soc. Hort.*
289 *Sci.*, **97**: 414-418.
- 290 8. Jaworska, G. (2005) Content of nitrates, nitrites, and oxalates in New Zealand
291 spinach. *Food Chem.* **89**: 235-242.
- 292 9. Canali, S., Montemurro, F., Tittarelli, F., and Masetti, O. (2008).Effect of nitrogen
293 fertilization reduction on yield, quality and N utilization of processing spinach. *J.*
294 *Food. Agric. Environ.* **6**(3&4): 242-247.
- 295 10. Wang, Z. H., Zong, Z.Q., Li, S. X., and Chen, B. M. (2002). Nitrate accumulation in
296 vegetables and its residual in vegetable fields. *Environ. Sci.* **23**(11): 79-83.
- 297 11. Fadaly, K.A., and Mishriky, J.F. (2000). Effect of nitrogen sources and levels on
298 growth, yield and mineral composition of spinach. Bulletin of Faculty of Agriculture,
299 University of Cairo. **41**(3): 973-988.
- 300 12. Assiouty, M.M., Sedera, S.A. and Giza, D. (2005). Effect of bio and chemical
301 fertilizers on seed production and quality of spinach. *Int. J. Agril.Biol.* **7**(6): 947–952.
- 302 13. Boroujerdnia, M. and Ansari, N.A. (2007).Effect of Different Levels of Nitrogen
303 Fertilizer and Cultivars on Growth, Yield and Yield Components of Romaine Lettuce
304 (*Lactuca sativa* L.). *Middle Eastern and Russian J. Plant. Sci. Biotech.* **1**(2): 47-53.
- 305 14. Ambia, K., Hossain, M., Haque, M., Mosiur, A., Halder, C. and Krisna, S. (2016).
306 Evaluation Seed Yield and Quality of Different Genotypes of Spinach (*Beta*
307 *vulgaris*). *Sci. Agri.* **16** (2): 61-66.
- 308 15. Zaki, M.S. and Halawa, S.S. (2016). Effect of nitrogen fertilizer and growth
309 stimulants on flowering behavior, seed yield and its quality of spinach. *Annal. Agric.*
310 *Sci.* **54**(2): 345-354.