

**Growth Performance of *Crescentia cujete* (Robx) Seedlings as Influenced by
Different Watering Regimes**

ABSTRACT

This study investigated the effects of different watering regimes on the growth of *Crescentia cujete*. The seedlings were subjected to six different watering regimes which include: W_1 – control (watering everyday), W_2 – watering once every two days, W_3 – watering once every three days, W_4 – watering once every four days, W_5 – watering once every five days and W_6 – watering once every six days; each treatment replicated six times. The experiment was arranged in a Completely Randomized Design (CRD). The study was carried out at the Central Nursery of Forestry Research Institute of Nigeria, Ibadan. Growth parameters assessed include: plant height (cm), collar diameter (mm), leaf production and the biomass accumulation which was sectioned into leaf, stem and the root. The data collected were subjected to one-way Analysis of Variance (ANOVA) and means separation was done using Duncan Multiple Range Test (DMRT) at 5% level of probability. The results showed that there were significant differences among the treatments in plant height and number of leaves produced while there was no significant difference in the collar diameter. However, seedlings watered once every 5 days (W_5) performed best in height (20.48 cm) and leaf production (18.42). It was revealed that there were significant differences in the leaf and stem dry weights among the treatments with W_5 having the highest weight while there were no significant differences in the root dry weight. *Crescentia cujete* optimized water shortage for its growth.

Keywords: Crescentia cujete, watering regime, growth, dry weight

1. INTRODUCTION

Crescentia cujete commonly referred to as Calabash tree is a forest tree that belongs to the family of Bignoniaceae which originated from tropical America and later introduced to Africa [1]. According to [2], it grows to about 6 to 9 m in height with a wide crown and long branches covered with clusters of tripinnate leaves and gourd-like fruit. The large fruit has a thin hard shell and whitish pulp with dark-brown seeds that are thin and flat. The spoon-shaped leaves are 5 to 18 cm long and 2 to 5 cm broad and they are arranged in clusters along the stout twigs. It is frequently cultivated as ornamental tree.

Nearly all parts of the tree are utilized for diverse articles and equipment, the wood is used as tools handles, ribs in boat building and gourds for containers and musical instruments [3]. They are valued in the manufacture of handicrafts and musical instruments. The wood has specific gravity of 0.6 to 0.8; it is strong, flexible, moderately hard and heavy. Hence, it can be used for firewood and construction in rural

36 areas and in the manufacture of handles for agricultural implements. The fruit pulp is laxative, emollient,
37 expectorant and fever medicine. Fresh seeds are ground and mixed with water to make a refreshing drink
38 which has sweet and pleasant taste [4].

39 Phenols and tannins are present in the fruit sample of *C. cujete*. Phenol and phenolic compounds have
40 been extensively used in disinfections and remain the standard with which other bactericides are
41 compared in official test [5]; [3]. This underscores why *C. cujete* is used as disinfectants and bactericides
42 in emollient healing and in the treatment of burns [1]. Flavonoids found in *C. cujete* can act as anti-
43 oxidants and protects the cells of the body form radical damage; free radicals are believed to damage cell
44 and inflict various health-related problems [6].

45 Water, which is in a continual state of flux, is a very essential factor in the growth and development of
46 plants [7]. Various vital processes in plants such as cell division, cell elongation, stem as well as leaf
47 enlargement and chlorophyll formation depends on plant water availability [8]; [9]. There is usually
48 structural deformity in plants frequently leading to death when there is insufficient water below critical
49 level [10]. The reduction in relative water contents affects physiological processes and hence plant growth
50 (Awodola, 1984 unpublished MSc dissertation); [11].

51 Water is required by plants for the manufacture of carbohydrates and as a means for transportation of
52 essential nutrients and minerals and cooling of plant leaves. About ninety-five percent of water in plants is
53 used up in cooling via evapotranspiration while the remainder is utilized in physiological processes such
54 as photosynthesis and respiration [12] and evapotranspiration rates increases with higher temperature
55 [13]; [14]. However, water availability is seasonal coupled with climate change effects which manifests in
56 longer dry spell or inundation giving way to waterlogging.

57 The water requirements of tree species differ due to their varying morphological and/or genetic makeup;
58 this explains the disparate adaptive tendencies of different plants to water availability extremes such as
59 water deficit and waterlogging [15]. Season-dependent water availability in nurseries spawns the need to
60 study the responses of different tree seedlings in the nursery to different watering regimes in order to
61 achieve optimum growth and engender effective water management.

62 The information about the response of *C. cujete* to different watering regime is sparse despite its diverse
63 medicinal and domestic potentials. Hence, the objective of this study is to investigate the effects of
64 different watering regimes on the early growth of *C. cujete* with a view to contributing to its conservation
65 and efficient water management in the nursery.

66 **2. MATERIALS AND METHODS**

67 The experiment was carried out at the central nursery of Forestry Research Institute of Nigeria, Ibadan
68 located on latitude 7°23`N and longitude 3°58`E. The climate is characterized by wet (April to October)
69 and dry seasons (November to March). The annual rainfall ranging between 1300 - 1500 mm and annual
70 mean relative humidity of 80-85% (Forestry Research Institute of Nigeria Annual Meteorological Report,
71 2015).

72 The experiment was carried out between January and March, 2017. The seeds of *C. cujete* were
73 obtained from a mother tree at the herbal garden of Forestry Research Institute of Nigeria, Ibadan, Oyo
74 State, Nigeria. The top soil (0-20 cm) was collected from the floor of a *Gmelina arborea* plantation. The
75 seedlings were raised in a germination tray for two weeks, with sterilized river sand used as the growing
76 media. The two weeks old seedlings were transplanted into polythene pots filled with 2 kg of top soil and
77 watered to pot capacity with respect to the treatments. The growth parameters such as plant height (cm),
78 collar diameter (mm) and leaf production were assessed weekly while the biomass estimation was carried
79 out at the end of the growth assessment.

80 The height of the seedlings was measured with the aid of a measuring tape graduated in centimeters
81 (cm) from the soil surface while the collar diameter was determined with the aid of a Vernier caliper.

82 Biomass estimation was carried out by selecting three (3) seedlings from each treatment. The seedlings
83 were sectioned into leaves, stem and root and oven-dried at 70°C for 24 hours. The dry weight of the
84 leaves, stem and root of the seedlings were weighed and recorded. The mean of the dry weights of the
85 three seedlings selected in each treatment were calculated [16].

86 There were six treatments and each treatment was replicated six times making a total of thirty-six
87 experimental units in all.

88 The treatments are:

89 W_1 – control (watering everyday)

90 W_2 – watering once every two days

91 W_3 – watering once every three days

92 W_4 – watering once every four days

93 W_5 – watering once every five days

94 W_6 – watering once every six days

95 The experimental design used was Completely Randomized Design (CRD). The data collected were
96 subjected to a one-way Analysis of Variance (ANOVA) at 5% level of probability while the means found to
97 be significant were separated using Duncan Multiple Range Test (DMRT) at 5% level of probability [17].

98 3. RESULTS AND DISCUSSION

99 **Table 1: Particle size distribution and chemical properties (0-20 cm)**

Soil Properties	Values
pH (H ₂ O)	5.71
Organic carbon (g/kg)	9.6
Total Nitrogen (g/kg)	0.72

P (mg/kg)	7
K (cmol/kg)	0.6
Ca (cmol/kg)	7.0
Mg (cmol/kg)	5.79
Mn (mg/kg)	23.6
Fe (mg/kg)	6.0
Cu (mg/kg)	12
Zn (mg/kg)	3.08
Sand (%)	84.5
Clay (%)	9.9
Silt (%)	5.6
Textural Class	Loamy sand

100

101 According to Federal Fertilizer Department of the Federal Ministry of Agriculture and Rural Development
 102 (2011), the soil was moderately acidic; the nitrogen and organic carbon contents were low while there
 103 was moderate amount of phosphorus and potassium (Table 1).

104 **Table 2: Weekly Mean Temperatures and Relative Humidity in the Study Area**

Weeks	Maximum temperature (°C)	Minimum temperature (°C)	Mean relative humidity (%)
1	35.6	21.5	50.2
2	34.3	23.1	56.9
3	34.4	24	69.5
4	35.3	23.5	61
5	31.7	23.9	78
6	32.8	24.3	68.9
7	35	22.2	38.4
8	36.1	24.4	59.8

9	35.4	24.3	64.2
10	35.1	24.2	70.1
11	34.6	24	68.2
12	31.8	23.8	74.9

105 **Source: Forestry Research Institute of Nigeria Meteorology Section**

106 The maximum temperature ranged between 31.7 °C to 36.1°C, the peak was recorded on the eighth week
 107 while the minimum was recorded on the fifth week. The minimum temperature ranged between 21.5 °C to
 108 24.4 °C and the mean relative humidity ranged between 50.2% to 78% (Table 2).

109 **Table 3: Analysis of Variance of the Growth Parameters**

Parameters	Source of variation	df	Sum of squares	Mean square	F	Significance
Height	Watering regime	5	25.327	5.065	4.871	0.002*
	Error	30	31.194	1.040		
	Total	35	56.522			
Collar Diameter	Watering regime	5	0.341	0.068	0.622	0.684 ^{ns}
	Error	30	3.292	0.110		
	Total	35	3.633			
Leaf Production	Watering regime	5	59.206	11.841	3.142	0.021*
	Error	30	113.065	3.769		
	Total	35	172.271			

110 *ns – not significant; * - significant (p≤0.05)*

111
 112 **Table 4: Mean Values of the Growth Parameters of *Crescentia cujete* Subjected to Selected**
 113 **Watering Regimes**

Treatments	Height (cm)	Leaves	Collar diameter (mm)
W ₁	18.01 ^a	15.76 ^a	4.82

W ₂	18.86 ^{ab}	14.56 ^a	4.79
W ₃	19.99 ^{bc}	16.82 ^{ab}	4.65
W ₄	20.01 ^{bc}	14.96 ^a	4.97
W ₅	20.48 ^c	18.42 ^b	4.88
W ₆	19.85 ^{bc}	16.57 ^{ab}	4.81

114 **Means followed by the same superscripts in the same column are not significantly different (p=**
 115 **.05)**

116 3.1 Mean Height, Collar Diameter and leaf production

117 The effect of watering regime was significant on height of seedlings of *Crescentia cujete* at 5% level of
 118 probability (Table 3). The seedlings watered once in five days (W₅) significantly recorded the highest
 119 mean value of 20.48 cm, which was not significantly different from the seedlings watered once in four
 120 days (W₄) with mean value 20.01 cm, W₆ (19.85 cm) and W₃ (19.99 cm) while seedlings watered once
 121 **everyday** (W₁) had the least mean value of 18.01 cm which was not significantly different from W₂ (18.86
 122 cm) (Table 4).

123 There was no significant difference in the collar diameter of the seedlings of *Crescentia cujete* at 5% level
 124 of probability. However, seedlings watered once in four days (W₄) had the highest mean value of 4.97 mm
 125 closely followed by seedlings watered once in five days (W₅) with mean value of 4.88 mm while the
 126 seedlings watered once in three days (W₃) had the least mean value of 4.65 mm (Table 4).

127 In terms of leaf production, there was significant difference among the treatments used at 5% level of
 128 probability. The seedlings watered once in five days (W₅) produced the highest number of leaves with
 129 mean value of 18.42 followed by seedlings watered once in three days (W₃) with mean value of 16.82 and
 130 the seedlings watered once in two days (W₂) produced the least number of leaves with mean value of
 131 14.56 (Table 4).

132 **Table 5: Mean Values of the Biomass of Sectioned Parts of *Crescentia cujete***

Treatments	Leaf Dry Weight (g)	Stem Dry weight (g)	Root Dry weight (g)
W ₁	2.52 ^{ab}	1.75 ^b	2.87
W ₂	1.93 ^a	1.18 ^a	2.30

W ₃	2.25 ^{ab}	1.20 ^a	2.37
W ₄	1.82 ^a	1.18 ^a	2.32
W ₅	2.92 ^b	1.80 ^b	3.08
W ₆	2.29 ^{ab}	1.30 ^{ab}	2.18

133 **Means followed by the same superscripts in the same column are not significantly different (p**
 134 **=.05)**

135 **3.2 Biomass accumulation**

136 **3.2.1 Leaf Biomass Accumulation**

137 The result showed that the selected watering regimes had significant effect on the leaf biomass
 138 accumulation of *Crescentia cujete* with W₅ recording the highest mean value of 2.29 g and W₄ recording
 139 the least mean value which was not significantly different from mean value recorded by W₂ (1.93 g) (Table
 140 5).

141 **3.2.2 Stem biomass accumulation**

142 The result showed that the stem biomass accumulation of *Crescentia cujete* was significantly affected by
 143 the selected watering regimes. W₅ had the highest mean value of 1.80 g which is not significantly different
 144 from W₁ and W₆ with mean values of 1.75 g and 1.30 g respectively while W₂ and W₄ recorded the least
 145 mean value of 1.18 g (Table 5).

146 **3.2.3 Root biomass accumulation**

147 The effect of watering regime on root biomass accumulation of *Crescentia cujete* is not significantly
 148 different among the treatments. This implies that irrespective of the selected watering regimes, the root
 149 dry weight of *Crescentia cujete* may not differ (Table 5).

150 **4. DISCUSSION**

151 The resultant effect of different watering regimes on the height of *Crescentia cujete* agrees with the
 152 findings of [11] who reported that *Parkia biglobosa* subjected to watering once in five days had the best
 153 performance in growth. [18] recommended watering once in two to three days for the optimum growth for
 154 *Acacia senegal*. [9] reported that watering twice a week is suitable to achieve the optimum growth of
 155 *Acacia senegal* in the nursery.

156 The seedlings watered once in five days (W₅) produced the highest number of leaves which was not
 157 significantly different from the seedlings watered once in three days (W₃) and the seedlings watered once
 158 in six days (W₆). [11] observed similar results in studies of different watering regimes on the growth of
 159 *Parkia biglobosa* seedlings where seedlings watered once in three days or once in five days yielded the

160 highest number of leaves. [19] reported that plants optimize the morphology, physiology and metabolism
161 of their organs and cells in order to maximize productivity under the moisture stress conditions.
162 [11] also observed that *Parkia biglobosa* seedlings watered once in three days gave the highest dry plant
163 weight. [20] reported that reduced soil moisture may cause a reduction in root growth and limit nutrient
164 uptake by roots; this is contrary to the result from this study. It has been widely reported that extensive
165 root growth is an adaptive feature under drought stressed conditions [21]. Shoot dry weight increases with
166 increase in soil moisture but under water stress conditions, root growth exhibits better growth than the
167 shoot. This causes the increase in root to shoot ratio [22].

168

169 **5. CONCLUSION**

170 In this study, the effect of selected watering regimes on *Crescentia cujete* was examined. It is evident that
171 *C. cujete* can optimize water deficit for its growth. For effective water management coupled with optimum
172 growth during the early growth of *C. cujete* seedlings, water can be applied once in four to five days.

173 **REFERENCES**

- 174 1. Michael A. Trees, Shrubs and Liannas of West Africa Dry Zones. Grad Margae Publishers
175 GMBH, MNHN. 2004; 191.
- 176 2. Gilman EF and Watson DG. Fact Sheet Florida Cooperative Extension Service. University of
177 Florida; 1993
- 178 3. Ejelonu BC, Lasisi AA, Olaremu AG and Ejelonu OC. The Chemical Constituents of Calabash
179 (*Crescentia cujete*). *African Journal of Biotechnology* 2011;10(84):19631-19636.
- 180 4. UNCTAD. Market Brief in the European Union for Selected Natural Ingredients Derived from
181 Native Species - *Crescentia cujete* (Calabash, Totumo). *United Nations Conference on Trade and*
182 *Development/Biotrade Facilitation Programme*. 2005
- 183 5. Carter SJ. Tutorial pharmacy. 6th Edn. Pitman Medical Publishing Co. Ltd. Tunbridge Well Kent,
184 England. 1979:29-30.
- 185 6. Phaniendra A, Jestadi DB and Periyasamy L. Free Radicals: Properties, Sources, Targets, and
186 their Implications in Various Diseases. *Indian Journal of Clinical Biochemistry*. 2015;30(1):11-26.
- 187 7. Ördög V. The Agricultural Engineering *MSc curriculum development TÁMOP- 4.1.2-08 / 1 / A-*
188 *2009-0010 project*. 2011
- 189 8. Price DT, Black TA and Kelliher FM. Effects of Sala Understorey Removal on Photosynthesis
190 Rate and Stomata Conductance of Young Douglas-Fir Trees. *Canadian Journal of Forest*
191 *Resource*. 1986;16: 90-97pp.

- 192 9. Ogun MB, Adeduntan SA and Suberu SA. Influence of Watering Regime and Mycorrhizae
193 Inoculations on the Physiology and Early Growth of *Acacia senegal* (L.) Wild. *African Journal of*
194 *Plant Science* 2010;4(7):210-216.
- 195 10. Levy Y and Krikum J. Effects of Irrigation, Water and Salinity and Root-Stock on the Vertical
196 Distribution of Vesicular Arbuscular Mycorrhiza on Citrus Roots. *New Phytology*. 1983; 15:397-
197 403.
- 198 11. Sale FA. Evaluation of Watering Regimes and Different Pot Sizes in the Growth of *Parkia*
199 *biglobosa* (Jacq) Benth Seedlings under Nursery Condition, *European Scientific Journal*
200 2015;11(12):313-325
- 201 12. Pallardy, SG. Physiology of Woody Plants. 3rd ed. *Academic Press*. 2008;95. ISBN 978-0-12-
202 088765-1
- 203 13. Chaouche K, Neppel L, Dieulin C, Pujol N, Ladouche B, Martin E, *et al*. Analyses of Precipitation,
204 Temperature and Evapotranspiration in a French Mediterranean Region in the Context of Climate
205 Change. *Comptes Rendus Geoscience* 2010; 342(3):234-243.
- 206 14. Synder RL, Moratiel R, Zhenwei S, Swelam A, Jomaa I and Shapland T Evapotranspiration
207 Response to Climate Change. 28th *International Horticultural Congress on Science and*
208 *Horticulture for People: International Symposium on CLIMWATER 2010: Horticultural Use of*
209 *Water in a Changing Climate*. Eds: J.E. Fernandez and M. I. Ferrereira. *Acts Hort*. 922
- 210 15. Chavarria G and Pessoa dos Santos H. Plant Water Relations: Absorption, Transport and Control
211 Mechanisms, *Advances in Selected Plant Physiology Aspects*, Dr. Giuseppe Montanaro (Ed.),
212 2012 ISBN:978-953-51-0557-2, InTech, Available from:
213 [http://www.intechopen.com/books/advances-in-selected-plantphysiology-aspects/plant-water-](http://www.intechopen.com/books/advances-in-selected-plantphysiology-aspects/plant-water-relations-absorption-transport-and-control-mechanisms)
214 [relations-absorption-transport-and-control-mechanisms](http://www.intechopen.com/books/advances-in-selected-plantphysiology-aspects/plant-water-relations-absorption-transport-and-control-mechanisms)
- 215 16. Shipley B and Vu T. Dry Matter Content as a Measure of Dry Matter Concentration in Plants and
216 their Parts. *New Phytologist* 2002; 153:359-364
- 217 17. Steel RG and Torrie JH. *Biostatistics: Principles and Procedures*. McGraw Hill, Mexico. 1998
- 218 18. Isah AD, Bello AG, Maishanu HM and Abdullahi S. Effect of Watering Regime on the Early
219 Growth of *Acacia senegal* (linn) willd. Provenances. *International Journal of Plant, Animal and*
220 *Environmental Sciences* 2013; 3:52-56
- 221 19. Lisar SYS, Motafakkerzad R, Hossain MM and Rahman IMM. *Water Stress in Plants: Causes,*
222 *Effects and Responses*. Prof. Ismail Md. Mofizur Rahman (Ed.). 2012. ISBN: 978-953-307-963-9.

- 223 20. Arndt SK, Clifford SC, Wanek W, Jones HG and Popp M Physiological and Morphological
224 Adaptation of the Fruit Tree *Ziziphus rotundifolia* in Response to Progressive Drought Stress.
225 *Tree Physiology*. 2001; 21:705-715.
- 226 21. Guoxiong C, Krugman T, Fahima T, Korol AB and Nevo E. Comparative Study on Morphological
227 and Physiological Traits Related to Drought Resistance Between Xeric and Mesic *Hordeum*
228 *spontaneum* lines in Israel. *Barley Genetics Newsletter*. 2002; 32:22-33.
- 229 22. Chiatante D, Di-Iorio A, Sciandra S, Stefania G and Mazzoleni S. Effect of drought and fire on root
230 development in *Quercus pubescens* Willd. and *Fraxinus ornus* L. seedlings. *J. Environ. Exp. Bot.*
231 2006; 56:190-197.
232