

### **Response of Heat Tolerant Variety (Kufri Surya) of Potato (*Solanum tuberosum*) Under Different Levels of Nitrogen**

#### **Abstract**

Keeping in view, the deficiency of detailed information on adoption of heat tolerant potato (*Solanum tuberosum*) variety 'Kufri Surya' in Terai Agro-Climatic situation of West Bengal, the field experiment was conducted at the Instructional farm of Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar, West Bengal during the rabi season of 2016 to study the effect on heat tolerant variety (Kufri Surya) with different nitrogen levels. Experiment was laid out in a Split-plot design taking two varieties 'Kufri Jyoti' and 'Kufri Surya' as main plot with six different levels of nitrogen of 0 kg N ha<sup>-1</sup>, 50 kg N ha<sup>-1</sup>, 100 kg N ha<sup>-1</sup>, 150 kg N ha<sup>-1</sup>, 200 kg N ha<sup>-1</sup> and 250 kg N ha<sup>-1</sup> as subplot. Results of the experiment showed that the higher values of the growth attributes like dry matter accumulation, leaf area index, in all the sampling dates of experimentation was recorded with 100 kg N ha<sup>-1</sup>. Owing to the higher leaf area index and dry matter accumulation in shoot, tuber yield was recorded highest from the treatment having 100 kg N ha<sup>-1</sup> (28.46 t ha<sup>-1</sup>).

**Key words:** Potato, Nitrogen, Kufri Surya, Kufri Jyoti, Heat Tolerant Variety.

## 23 Introduction

24 Potato (*Solanum tuberosum* L.) is the third most important food crop in the  
25 world after rice and wheat in terms of human consumption. India ranks as the world's  
26 2<sup>nd</sup> largest potato producing nation after China. Production in India is about 48.52  
27 million tonnes (Government of India, 2018) of which 26% are produced by West  
28 Bengal itself. Potato is a cool season long day crop. High temperatures and long days  
29 favour assimilate partitioning to the above ground vegetative parts, as a result, above  
30 ground bio-mass and plant height is increased and tuber yield is reduced (Wolf *et al.*,  
31 1990). Potato gives good yield at day temperature of 30-35 °C. But if night temperature  
32 go beyond 22 °C, there will be little tuberization even when day temperature is 25-27  
33 °C. Due to intense climate change the favourable temperatures for its growth is  
34 increased at its later stages hampering the tuberization. On this context, whether the  
35 heat tolerant variety 'Kufri Surya' could perform better than check variety 'Kufri Jyoti'  
36 was brought into notice from the experiment. Nitrogen is beneficial for the tuber  
37 quality, dry matter production, size of tubers etc. More application of nitrogen  
38 fertilizers can increase size of tubers and hence the yield but there is a particular dose  
39 limit up to which it will show positive results; beyond that limit the application of  
40 nitrogen fertilizer will not increase the yield but rather it would be harmful because of  
41 deposition of nitrogen in tubers in the form of nitrates (Mohammad and  
42 Mohammadreza, 2012) which is not at all favourable for human consumption and  
43 moreover excessive application can cause environmental pollution. The use of low N  
44 results in reduction in yield of potato. Judicious use of balanced dose of fertilizers is  
45 very critical for higher tuber yield. Keeping the above facts in view, a field experiment  
46 was undertaken to study the effect of different levels of nitrogen on growth and yield of  
47 potato cultivars 'Kufri Surya' and check variety of 'Kufri Jyoti'.  
48

## 49 Materials and Methods

50 A field experiment was conducted to study the effect of different doses of  
51 nitrogen on two different varieties of potato that is, 'Kufri Jyoti' and heat tolerant  
52 variety 'Kufri Surya' at Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Bihar,  
53 West Bengal during *rabi* season of 2016. The farm is situated at 26°12'78"N latitude  
54 and 89°24'55" E longitude at an elevation of 43 meters above mean sea level. The  
55 climatic zone where the farm is situated is in *Terai* zone which is subtropical in nature  
56 having its prominent characteristics of very high rainfall, high humidity and a  
57 prolonged winter season. The average rainfall of this zone varies from 2000-3000 mm.  
58 The soil of the experimental field was sandy loam in texture, a true representative of the  
59 *terai* region of West Bengal with a pH of 5.6. The experiment was carried out in split  
60 plot design with two varieties of potato 'Kufri Surya' and 'Kufri Jyoti' as main plots  
61 and six nitrogen levels as subplots i.e, 0 kg N ha<sup>-1</sup>, 50 kg N ha<sup>-1</sup>, 100 kg N ha<sup>-1</sup>, 150 kg  
62 N ha<sup>-1</sup>, 200 kg N ha<sup>-1</sup> and 250 kg N ha<sup>-1</sup>. The experiment had three replications with a plot  
63 size of 5m x 3.45 m and a spacing of 45 x 15 cm. The crop was planted on 26<sup>th</sup>  
64 November of 2016 and harvested on 4<sup>th</sup> March of 2017.

65 Healthy cut tubers were selected each having two-three eyes weighing 25-40  
66 g. The seed tubers were treated to protect them from an attack of fungal diseases when  
67 planted in the field. So, before planting, the seed tubers were dipped in solution of  
68 Acetochlor @ 2.5 g lit.<sup>-1</sup>+ streptomycin (Plantamycin) @ 2.5 g lit.<sup>-1</sup>of water for 15  
69 minutes and then they were dried in shade to protect it from direct sunlight prior to  
70 planting. Farmyard manure was applied on the field @ 5t ha<sup>-1</sup> at the time of final land  
71 preparation. The different doses of nitrogen were 0, 50, 100, 150, 200 and 250 kg N ha<sup>-1</sup>  
72 + 100 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>+100 kg K<sub>2</sub>O ha<sup>-1</sup>were given respective plots. Out of these doses  
73 1/3rd of nitrogen and full dose of P<sub>2</sub>O<sub>5</sub> and full dose of K<sub>2</sub>O were applied as basal at the  
74 time of planting of tubers. The rest half of the 2/3rd nitrogen was given in two equal  
75 splits, one as first top dressing at 21 DAP ( days after planting) and the second split of  
76 Nitrogen was applied at second top dressing at 41 DAP. The fertilizers were applied by  
77 broadcasting method in the form of Urea, Single Super Phosphate (SSP) and Muriate of  
78 Potash (MOP) as the sources of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O respectively. Two irrigations were  
79 given to the crop. First irrigation was given at 22 DAP after first top dressing and  
80 earthing up. Second irrigation was given at 44 DAP after second top dressing. Before  
81 10 days of harvesting of the crop dehaulming was done.

82 The growth attributes like number of haulms per plant, leaf area index (LAI),  
83 crop growth rate(CGR),net assimilation rate( NAR) were recorded at 20,40, 60 and 80  
84 DAP( days after planting). The photosynthetic and transpiration parameters like net  
85 photosynthesis rate, leaf stomatal conductance, transpiration rate were recorded at 20,  
86 40, 60DAP. Since at 80 DAP plants starts showing senescence no parameters were  
87 recorded. Though the photosynthetic and transpiration parameters were recorded by the  
88 instrument CI -340 Handheld Photosynthesis system.

89 The CI-340 Handheld Photosynthesis System is a portable, light-weight single-  
90 handed tool that measures photosynthesis, respiration, transpiration, stomatal  
91 conductance, PAR and internal CO<sub>2</sub>. It was designed for field use. It has some optional  
92 accessory modules which allow researchers to control CO<sub>2</sub>, H<sub>2</sub>O, temperature, light  
93 intensity, and measure chlorophyll fluorescence, while the ten different customized  
94 chambers can accommodate any leaf size, including conifer needles and cacti. Direct  
95 chamber connection to the CO<sub>2</sub>/H<sub>2</sub>O gas analyzer reduces measurement delay and  
96 enables rapid measurement of gas exchange with minimal delay. For measurement of  
97 photosynthetic activity, simply a potato leaf was secured within the CI-340 chamber  
98 and thereby “ambient” or “closed” system measurement was chosen. In moments, leaf  
99 function data was recorded and stored in the instrument, as the CI-340 analyzed  
100 photosynthesis rate , transpiration rate, and stomatal conductance rate. The data was  
101 downloaded from computer afterwards from the data memory chip present in the  
102 instrument.

103 **Theory of how the CI-340 Handheld Photosystem works is as follows : The rate at**  
104 **which photosynthesis occurs is determined by measuring the rate at which a**  
105 **known leaf area assimilates the CO<sub>2</sub> concentration in a given time. It is known that**  
106 **transpiration is the primary determinant of leaf energy balance and plant water**  
107 **status. The rate of transpiration is determined by the accumulation of water vapor**  
108 **flux per one-sided leaf area in a given time. Stomatal conductance is the water loss**

109 of a leaf's conductance considered in parallel or series. It can be obtained by  
 110 measuring the transpiration and leaf surface temperature (°C). The required  
 111 parameters can be obtained by applying following calculations (Handheld  
 112 Photosynthesis System CI-340 from CID Bio-Science, 2019).

113 The Net Photosynthesis Rate (Pn) (micro mol m<sup>-2</sup> s<sup>-1</sup>) is calculated by the  
 114 following formula:

$$115 \quad P_n = -W \times (C_o - C_i) = -2005.39 \times \frac{V \times P}{T_a \times A} \times (C_o - C_i)$$

116 Where, C<sub>o</sub>(C<sub>i</sub>): outlet (inlet) CO<sub>2</sub> concentration (ppm or micro mol/ mol)  
 117 W = Mass flow rate per leaf area, V = Leaf chamber volume, T<sub>a</sub> = air temperature (K), P  
 118 = Atmospheric pressure (bar) ,A = Leaf area (cm<sup>2</sup>).

119 The Transpiration Rate (E) (milimol m<sup>-2</sup> s<sup>-1</sup>) was calculated by:

$$120 \quad E = \frac{e_o - e_i}{P - e_o} \times W \times 10^3$$

$$121 \quad e_o = hr_o \times es100^{-1}$$

$$122 \quad e_i = hr_i \times es100^{-1}$$

123 Where, e<sub>o</sub>(e<sub>i</sub>): Outlet (inlet) water vapour (bar) ,W = Mass flow rate per leaf area ,T<sub>a</sub> =  
 124 air temperature (K) ,P = Atmospheric pressure (bar) ,hr<sub>o</sub> (hr<sub>i</sub>) = Outlet (inlet) relative  
 125 humidity (%).

$$126 \quad \text{Leaf Stomatal Conductance} = \frac{W}{e_o - e_i} \times \frac{p - e_o}{p} \times 1000$$

$$127 \quad \text{Leaf Stomatal Conductance} = \frac{e_{\text{leaf}} - e_o}{e_o - e_i} \times \frac{p - e_o}{p}$$

128 Where, e<sub>leaf</sub> = saturated water vapour at leaf temperature (bar) W = Mass flow  
 129 rate per leaf area, P = Atmospheric pressure (bar) ,e<sub>o</sub> (e<sub>i</sub>) : outlet (inlet) water vapour  
 130 (bar) ,R<sub>b</sub> = leaf boundary layer resistance (m<sup>2</sup>s mol<sup>-1</sup>) 0.3m<sup>2</sup>s mol<sup>-1</sup> is used. Observation  
 131 were done at 20,40,60 DAP.

132 The data collected from the field and laboratory experiments were subjected to  
 133 statistical analysis with appropriate design and treatment variations were tested for  
 134 significance by F-test (Cochran and Cox, 1955). The standard error of mean and critical  
 135 difference is indicated in the tables. For determination of critical difference at 5% level  
 136 of significance Fisher and Yates (1963) table was consulted. The statistical analysis was  
 137 evaluated by SPSS software.

138

## 150 Results and Discussion

### 151 *Effect on growth attributes and yield:*

152 Among the two varieties 'Kufri Surya' and 'Kufri Jyoti' both of them have  
153 given statistically similar results for most of the growth attributes at 20 and 80 DAP.  
154 This was because at 20 DAP the plants are yet to be developed because of its early  
155 stages of growth and at 80 DAP the plants started showing symptoms of **senescence** in  
156 stems and leaves. But quite significant differences were observed when the observations  
157 were taken at 40 and 60 DAP because it was the peak period of vegetative growth of the  
158 potato crop. In case of number of haulms per plant at any stages of the crop there were  
159 no significant differences among the varieties. At 40 and 60 DAP it was observed that  
160 'Kufri Jyoti' had performed 29 % better than 'Kufri Surya' at 40DAP and 24 % better  
161 at 60DAP. This can be pertained to 'KufriJyoti' having higher leaf area than 'Kufri  
162 Surya' whose leaves are narrower in shape resulting in lesser leaf area. Dry matter  
163 accumulation at 40 DAP was 19% more in 'KufriJyoti' because of more leaf area index,  
164 for which photosynthesis was more resulting in better accumulation of photosynthates  
165 and at 60 DAP Kufri Surya ( $303.13 \text{ g m}^{-2}$ ) performed better than KufriJyoti ( $279.53 \text{ g}$   
166  $\text{m}^{-2}$ ). This was due to mild attack of *Phoma spp.* on KufriJyoti at 60DAP for which  
167 growth of the plant was hampered. In case of crop growth rate Kufri Jyoti performed  
168 better at 20- 40 DAP ( $7.315 \text{ g m}^{-2} \text{ day}^{-1}$ ) and at 40-60 DAP Kufri Surya ( $8.569 \text{ g m}^{-2}$   
169  $\text{day}^{-1}$ ) gave maximum crop growth rate. Crop growth rate was hampered for Kufri Jyoti  
170 due to the same reason for which the dry matter accumulation was less at 60DAP. Net  
171 assimilation rate which is the amount of dry matter produced in gram per unit area of  
172 leaf per day was found significant at 40-60 DAP for both the varieties due to its peak  
173 period of growth in which Kufri Surya has performed 65% better than Kufri Jyoti at 40-  
174 60 DAP as it was resistant to pathogen attack. Though Kufri Jyoti ( $25.80 \text{ t ha}^{-1}$ ) was  
175 mildly affected by *Phoma spp.* at 60DAP, timely control measures had helped  
176 immensely to revert back its negative effects on yield and hence had shown better yield  
177 compared to Kufri Surya ( $19.76 \text{ t ha}^{-1}$ ). The reasons might be due to bigger size and  
178 weight of tubers per plant in case of Kufri Jyoti (Neshev et al.,2014).

179 Nitrogen is a very essential nutrient for growth of plants because its an  
180 important constituent of key photosynthetic enzyme RuBP carboxygenase / oxygenase.  
181 Total sugar accumulation in leaves and tubers are positively influenced by nitrogen  
182 application. Total sugar increased with the rate of N-fertilizer application. The higher  
183 sugar content was due to higher photosynthetic rate, which is enhanced due to  
184 enzymatic activity. Increase in nitrogen levels increases the carbohydrate production by  
185 more number of chlorophylls. But there is a limit of nitrogen application beyond which  
186 if nitrogen fertilizers are added the plants won't show a positive result. It was observed  
187 that number of haulms increased linearly with increase in dose of nitrogen since  
188 nitrogen has a positive role in increase in vegetative growth of plant ( Sriom et al .,  
189 2017). There was significant difference for most of the growth attributes in all the stages  
190 among the various nitrogen levels except 20 DAP because of early stages of growth. So,  
191 in 40 DAP maximum number of haulms were observed in the treatment of  $200 \text{ kg N ha}^{-1}$

192 <sup>1</sup> (3.35) and in 60 DAP for 250 kg N ha<sup>-1</sup> (5.75). For dry matter accumulation 100 kg N  
193 ha<sup>-1</sup> was found optimum for the maximum dry matter production at 40 DAP(171.46 g  
194 m<sup>-2</sup>) and 60 DAP (322.25 g m<sup>-2</sup>). These results were in accord with the findings of  
195 **Sharma and Sharma** (1991) and Kundu et al.,(2019).This might be assigned to LAI at  
196 40 DAP and 60 DAP having the highest value for 100 kg N ha<sup>-1</sup> as it was optimum  
197 amount nitrogen required for enlargement of leaves resulting in production of more  
198 photosynthates . Crop growth rate among the various nitrogen levels for both 20-40  
199 DAP and 40-60 DAP were statistically at par with each other. There was no significant  
200 difference among the various nitrogen levels in 20-40 DAP except 40-60 DAP.  
201 Maximum net assimilation rate (g m<sup>-2</sup> day<sup>-1</sup>) was observed at 200 kg N ha<sup>-1</sup> in 40-60  
202 DAP(1.087). Highest yield was obtained at 100 kg N ha<sup>-1</sup> due to better tuber  
203 development at the optimum level( Patel and Patel, 2001).

204

UNDER PEER REVIEW

**Table 1.**Effect of Variety and Nitrogen levels on number of haulms per plant, LAI and dry matter accumulation of plant.

TREATMENTS	Number of haulms per plant			Leaf Area Index(LAI)				Dry matter accumulation(g m <sup>-2</sup> )			
	20 DAP	40 DAP	60 DAP	20 DAP	40 DAP	60 DAP	80 DAP	20 DAP	40 DAP	60 DAP	80 DAP
V <sub>1</sub>	2.50	2.87	4.47	0.15	3.98	4.46	2.20	8.26	150.97	279.53	274.55
V <sub>2</sub>	2.38	2.93	5.00	0.14	3.08	3.60	2.10	8.68	131.75	303.13	272.66
<b>SEm(±)</b>	0.01	0.03	0.59	0.05	0.12	0.02	0.09	0.24	2.83	1.18	.94
<b>CD(0.05)</b>	NS	NS	NS	NS	0.72	0.10	NS	NS	17.24	7.18	NS
<b>Nitrogen levels</b>											
N <sub>0</sub>	2.15	2.25	3.40	0.11	3.04	3.39	1.78	7.21	112.78	218.01	218.33
N <sub>1</sub>	2.40	2.65	4.30	0.12	3.50	3.77	1.93	8.25	136.15	290.72	261.76
N <sub>2</sub>	2.70	2.90	5.05	0.13	4.40	4.70	2.45	9.53	171.46	322.25	327.76
N <sub>3</sub>	2.30	3.05	5.00	0.19	4.01	4.18	2.69	8.54	154.28	321.60	306.71
N <sub>4</sub>	2.75	3.35	4.90	0.17	3.10	4.53	1.97	9.15	128.46	297.72	263.63
N <sub>5</sub>	2.35	3.20	5.75	0.16	3.14	3.60	2.11	8.15	145.03	297.66	263.44
<b>SEm(±)</b>	0.17	0.13	0.20	0.01	0.14	0.10	0.06	0.41	3.88	7.34	5.75
<b>CD(0.05)</b>	NS	0.39	0.58	NS	0.41	0.30	NS	NS	11.46	21.65	16.96

V1-Kufri Jyoti, V2-Kufri Surya.N0-0 kg ha<sup>-1</sup>, N1-50 kg ha<sup>-1</sup>, N2-100 kg ha<sup>-1</sup>, N3- 150 kg ha<sup>-1</sup>,N4-200 kg ha<sup>-1</sup>, N5-250 kg ha<sup>-1</sup>.

209 **Table 2.**Effect of Variety and Nitrogen levels on Crop Growth Rate, Net Assimilation Rate and Yield of crop.

TREATMENTS	Crop growth rate ( g m <sup>-2</sup> )		Net assimilation rate( g m <sup>-2</sup> day <sup>-1</sup> )		Yield(t ha <sup>-1</sup> )
	20- 40 DAP	40-60 DAP	20-40 DAP	40-60 DAP	
V <sub>1</sub>	7.135	6.428	2.734	0.679	25.80
V <sub>2</sub>	6.153	8.569	2.825	1.120	19.76
<b>SEm(±)</b>	0.130	0.083	0.057	0.045	0.19
<b>CD(0.05)</b>	0.789	0.503	NS	0.142	1.17
<b>Nitrogen levels</b>					
N <sub>0</sub>	5.278	5.262	2.458	0.641	9.28
N <sub>1</sub>	6.395	7.729	2.832	0.789	20.45
N <sub>2</sub>	8.116	6.313	2.861	0.985	28.46
N <sub>3</sub>	7.287	8.366	2.954	0.905	26.78
N <sub>4</sub>	5.947	9.690	2.555	1.087	26.08
N <sub>5</sub>	6.844	7.632	3.016	0.991	25.63
<b>SEm(±)</b>	7.135	6.428	0.082	0.061	0.45
<b>CD(0.05)</b>	NS	NS	NS	0.179	1.33

210 V1-Kufri Jyoti, V2-Kufri Surya.N0-0 kg ha<sup>-1</sup>, N1-50 kg ha<sup>-1</sup>, N2-100 kg ha<sup>-1</sup>, N3- 150 kg ha<sup>-1</sup>,N4-200 kg ha<sup>-1</sup>, N5-250 kg ha<sup>-1</sup>.



211 ***Effect on photosynthetic parameters:***

212 There were significant differences among the varieties, maximum net  
213 photosynthesis rate was observed in 20 DAP (7.7 micro mol m<sup>-2</sup> s<sup>-1</sup>), 40 DAP  
214 (14.77 micro mol m<sup>-2</sup> s<sup>-1</sup>) in KufriJyoti and in 60 DAP in Kufri Surya (10.88 micro mol  
215 m<sup>-2</sup> s<sup>-1</sup>) which might be due to the possible reason of attack of pathogen on KufriJyoti  
216 hampering its healthy leaf growth and hence the photosynthetic activity. Both the  
217 varieties were statistically at par with each other with respect to transpiration  
218 rate. Maximum stomatal conductance rate was in Kufri Surya (269.42 millimol m<sup>-2</sup> s<sup>-1</sup>)  
219 at 20 DAP, in KufriJyoti (368.55 millimol m<sup>-2</sup> s<sup>-1</sup>) at 40 DAP which might be due to  
220 faster development of leaves of Kufri Surya at 20 DAP and better development of  
221 leaves and number of stomata in KufriJyoti at 40DAP.

222 Significant differences were observed between different nitrogen levels for  
223 various photosynthetic parameters. All the photosynthetic characters have been  
224 recorded maximum at 150 kg N ha<sup>-1</sup>. Stomatal conductance rate is the rate at which  
225 carbon dioxide is uptaken and water vapour is released through stomata. Nitrogen plays  
226 an important role in stomatal conductance by cell expansion and altering the cation and  
227 anion concentration of cytoplasmic environment which can actually change the stomatal  
228 conductance rate (Nasab et al., 2014). More nitrogen application also increases the leaf  
229 growth and hence the number of stomata increasing the stomatal conductance rate.  
230 Highest stomatal observations were found at 150 kg nitrogen ha<sup>-1</sup>, for all the stages with  
231 a maximum of 395.33 millimol m<sup>-2</sup> s<sup>-1</sup> at 60 DAP because this was the optimum dose  
232 above which no such effect was seen. Since stomatal conductance rate is closely related  
233 with transpiration rate, maximum transpiration rate similarly observed at 150 kg N ha<sup>-1</sup>  
234 at all stages of growth. Net photosynthetic rate may be assigned to the possible reasons  
235 of larger number of chlorophyll and stomata due to optimum doses of nitrogen.

236 From the above experiment it can be concluded that Kufri Surya didn't perform  
237 better than check variety KufriJyoti because the high temperature at which Kufri Surya  
238 might have shown better performance with respect to yield than KufriJyoti due to its  
239 heat tolerant characteristics which was not obtained.

240

241

242

243

244

245

246

247

248

249 **Table 3.**Effect of Variety and Nitrogen levels on photosynthetic parameters.

TREATMENTS	Stomatal conductance rate (millimol m <sup>-2</sup> s <sup>-1</sup> )			Transpiration rate (millimol m <sup>-2</sup> s <sup>-1</sup> )			Net Photosynthesis rate (micro mol m <sup>-2</sup> s <sup>-1</sup> )		
	20 DAP	40 DAP	60 DAP	20 DAP	40 DAP	60 DAP	20 DAP	40 DAP	60 DAP
V <sub>1</sub>	235.43	368.55	355.63	1.06	2.80	2.85	7.77	14.17	9.12
V <sub>2</sub>	269.42	343.33	365.55	0.99	3.19	2.95	5.66	12.64	10.88
<b>SEm(±)</b>	1.44	2.79	0.58	0.07	0.05	0.05	0.04	0.02	0.22
<b>CD(0.05)</b>	8.76	16.98	NS	NS	0.30	NS	0.27	0.10	1.42
N <sub>0</sub>	220.60	315.27	355.89	0.72	3.18	2.04	4.29	11.94	7.97
N <sub>1</sub>	229.09	333.97	353.43	1.01	2.44	2.54	5.45	13.14	8.77
N <sub>2</sub>	244.39	365.07	387.76	1.04	2.94	3.04	7.46	13.60	9.34
N <sub>3</sub>	292.51	389.69	395.33	1.33	3.68	3.54	10.15	15.05	12.52
N <sub>4</sub>	258.91	371.73	349.98	1.02	2.85	3.13	7.16	13.07	11.49
N <sub>5</sub>	269.06	359.92	321.16	1.04	2.87	3.09	5.77	13.65	10.14
<b>SEm(±)</b>	2.49	3.96	10.02	0.09	0.10	0.09	0.21	0.16	0.15
<b>CD(0.05)</b>	7.35	11.68	29.57	0.26	0.29	0.26	0.62	0.47	0.44

269

V1-Kufri Jyoti, V2-Kufri Surya.N0-0 kg ha<sup>-1</sup>, N1-50 kg ha<sup>-1</sup>, N2-100 kg ha<sup>-1</sup>, N3- 150 kg ha<sup>-1</sup>,N4-200 kg ha<sup>-1</sup>, N5-250 kg ha<sup>-1</sup>.

270 **REFERENCES**

- 271 Cochran, W.G. and Cox, G.M. 1957.(2nd Edition).*Experimental Design*.John Wiley and Sons.  
272 New York.Pp: 615.
- 273 Government of India. 2018.*Monthly report potato*. Horticulture Statistics Division.  
274 Department of Agriculture, Cooperation and Farmers Welfare. Ministry of Agriculture  
275 and Farmers Welfare,New Delhi.
- 276 Handheld Photosynthesis System CI-340 from CID Bio-  
277 Science.2019.[http://phytoscience.com/biotique/plant/handheld/photosynthesis-system-](http://phytoscience.com/biotique/plant/handheld/photosynthesis-system-ci-340-from-cid-bio-science/)  
278 [ci-340-from-cid-bio-science/](http://phytoscience.com/biotique/plant/handheld/photosynthesis-system-ci-340-from-cid-bio-science/) Accessed 9 August 2019.
- 279 Kundu C.K., Bera P.S., Giri A., Das S. , Datta M.K. and Bandopadhyay P.2019. \_Effect of Different  
280 Doses of Nitrogen and Potassium on Growth and Yield of Potato (*Solanum tuberosum* L.)  
281 under New Alluvial Zone of West Bengal. *Current Journal of Applied Science and Technology*  
282 **36(2): 1-5**.
- 283 Mohammad, V. and Mohammadreza, N.2012.The effect of various nitrogen fertilizer amounts  
284 of yield and nitrate accumulation in tubers of two potato cultivars in cold–IV regions of  
285 Isfahan (Iran). *International Journal of Agriculture and Crop Sciences* .**4(22): 1688-**  
286 **1691**.
- 287 Nasab H.M. , Siadat S.A., Naderi A., Lack S.andModhej A.2014.The effects of Drought Stress  
288 and Nitrogen Levels on Yield , Stomatal Conductance and Temperature Stabily of  
289 Rapeseed (Canola) Genotypes. *Advances in Environmental Biology*.**8(10):1239-1247**.
- 290 Neshev N, Manolov I, Chalova V, Yordanova N.2014. Effect of nitrogen fertilization on yield and  
291 quality parameters of potatoes. *Journal of Mountain Agriculture on the Balkan*.**17(3): 615-627**.
- 292 Patel J.C., Patel B.K. 2001. Response of potato to nitrogen under drip and furrow methods of irrigation.  
293 *Indian Potato Assoc*.**28(2- 4):293-295**.
- 294 Sharma, B.D. and Sharma, U.C. 1991. Dry matter accumulation behaviour of different potato  
295 cultivars (*Solanumtuberosum* L.).*Indian J Hill Farming*.**4(1): 11-14**.
- 296 Sriom, Mishra D.P. , Rajbhar P. , Singh D., Singh R.K., and Mishra S.K. 2017. \_Effect of Different  
297 Levels of Nitrogen on Growth and Yield in Potato (*Solanum tuberosum* L.) CV. Kufri Khyati.  
298 *International Journal of Current Microbiology and Applied Sciences* **6(6): 1456-1460**.
- 299 Wolf, S.A., Marani and Rudich, J. 1990.Effect of temperature and photoperiod on assimilate  
300 partitioning in potato plants. *Annals of Botany*.**66:513-520**.

301

302