

Original Research Article

Characterization of Polyphenols and Mineral Contents in Three Medicinal Weeds

ABSTRACT

Aims: Common weeds *Rorippa palustris* (L.) Besser, *Euphorbia rothiana* Spreng. and *Schoenoplectiella articulata* (L.) Lye are used for food, medicinal, green biofertilizer and bio-sorbent applications. In this work, their polyphenol and mineral contents have been characterized.

Methodology: Samples from aforementioned three plants were manually collected in Raipur city (CG, India) and processed for the analyses. Folin-Ciocalteu and aluminum chloride were used for the spectrophotometric determination of polyphenols. The mineral contents were quantified by X-ray fluorescence.

Results: The total concentration of 20 elements (viz. P, S, Cl, As, Se, K, Rb, Mg, Ca, Sr, Ba, Al, Ti, Cr, Mn, Fe, Co, Zn, Mo and Pb), total polyphenol and flavonoid contents in the leaves ranged from 46372 to 71501, from 47877 to 73791 and from 1950 to 9400 mg/kg, respectively. Remarkable concentrations of several nutrients (P, S, Cl, K, Mg, Ca and Fe) were observed.

Conclusion: The biomass from medicinal weeds *R. palustris*, *E. rothiana* and *S. articulata* featured very high K, Ca and Fe contents. Other nutrients (polyphenols, flavonoids, P, S, Cl and Mg) were identified at moderate levels. These species may hold promise as bioindicators.

22 | **Keywords:** flavonoid; medicinal weeds; mineral elements; phenolic; XRF; ~~flavonoid; phenolic;~~
 23 | ~~mineral elements.~~

24 | 1. INTRODUCTION

25 | Aquatic plants not only provide food and habitat for many animals such as fish and waterfowl,
 26 | but also are potential bio-sorbents for the accumulation of trace elements [1], and may be used as
 27 | green manure to restore soil fertility [2]. Moreover, some wetland plants have medicinal value,
 28 | and are frequently used in Ayurveda and folk medicine [3].

29 | *Rorippa palustris* (bog yellowcress or marsh yellowcress) grows in many types of damp, wet,
 30 | and aquatic habitats. It has antiscorbutic properties; its roots are diuretic and it is used in the
 31 | treatment of measles [4].

32 | *Euphorbia rothiana* (white latex; in India: Chagul putputi, Merashupal chedi, and Palootti chedi)
 33 | is an annual erect, glabrous, profusely branched subshrub of ≈ 1 -m height, widely distributed in
 34 | India, Sri Lanka, China and Indonesia. The leaves of *E. rothiana* (*Rubus ellipticus* Sm. and *Rubus*
 35 | *racemosus* Roxb.) are chewed together to relieve symptoms of sudden sickness and giddiness
 36 | thought to be caused by evil spirits. Further, its biomass is used as a medicine for cattle. Its latex
 37 | is externally applied for healing sores, it is also believed to promote hair growth, and the whole
 38 | plant is used for household insect repellent [5-6].

39 | *Schoenoplectiella articulata* (in India: Chichora, Ciccodaka, Gaichira, Laghukaseruka, Pappati
 40 | Chickha, Tan-pokli, and Tsjeli) is an aquatic annual herb distributed through Indo-China to
 41 | Malaysia regions. Its tubers are effective for against diarrhea and vomiting, and its fruits and
 42 | leaves in bodyache, pain and fevers. It has been reported to have anthelmintic, antibacterial,

43 antiemetic, antifungal, carminative, contraceptive, digestive, febrifuge and sedative properties
44 [7].

45 Similar species, e.g., *Schoenoplectus lacustris* (L.) Palla, *Rorippa globosa* (Turcz.) Hayek and
46 *Euphorbia macroclada* Boiss. have been applied in the accumulation and phytoremediation of
47 heavy metals in contaminated soils [8-10]. The bioaccumulation of various elements (P, K, Mg,
48 Ca, Cr, Mn, Fe, Co, Zn, Pb, Cu and Cd) in macrophytes (e.g., *Azolla pinnata* R.Br., *Pistia*
49 *stratiotes* L., *Solvinia molesta* D.Mitch., *Trapa natans* L., *Persicaria maculosa* Gray and
50 *Nelumbo nucifera* Gaertn.) has also been reported in the literature [11-22].

51 In this work, the total polyphenols, flavonoids, and trace element (P, S, Cl, As, Se, K, Rb, Mg,
52 Ca, Sr, Ba, Al, Ti, Cr, Mn, Fe, Co, Zn, Mo and Pb) contents in the biomass from these three
53 weeds (*R. palustris*, *E. rothiana* and *S. articulata*) are described.

54 2. MATERIALS AND METHODS

55 *R. palustris* (RP), *E. rothiana* (ER) and *S. articulata* (SA), shown in **Fig. 1**, were botanically
56 recognized [23]. Samples were collected from a municipal waste dumping area in Raipur city,
57 CG, India (21.25°N 81.63°E) in March-June 2017. The plants biomass (including leaves, stems
58 and fruits) was washed thrice with deionized water and sun-dried in a glass room for one week.
59 The crushed samples were sieved out to separate particles of mesh size ≤ 0.1 mm, they were
60 further dried at 50 °C overnight, and they were finally stored in a deep freezer at -4 °C till the
61 analyses were conducted.



Fig. 1. (From left to right) Photographs of biomass from *Rorippa palustris*, *Euphorbia rothiana* and *Schoenoplectus articulata*.

The polyphenol content in the samples in powder form was determined by extraction in an acetone:water mixture (7:3, v/v), equilibrating 0.1 g of sample with the solvent, as indicated by Bertaud et al. [24]. For the analysis of the total polyphenol content, an aliquot of the extract was treated with Folin-Ciocalteu reagent (Sigma-Alrich, F9252) to form a blue colored polyacid [25]. A standard calibration curve was prepared by using tannic acid (Sigma-Alrich, 403040) as the reference material. The flavonoid content was determined by reacting an aliquot of the extract with aluminum chloride (Sigma-Alrich, 563919), as described in the literature [26], using quercetin (Sigma-Alrich, Q4951) as the reference material.

A Bruker Tracer 5i portable X-Ray Fluorescence (pXRF) instrument with a Rhodium tube was used for the analysis of trace elements, following the empirical calibrations described by Towett et al. [27].

All measurements were performed in triplicate, and their mean values were reported.

79 3. RESULTS AND DISCUSSION

80 3.1. Polyphenol Contents

81 The chemical characteristics of the biomass of RP, ER and SA are summarized in **Table 1**. The
82 total phenolic content (TPh) and the flavonoid content (Fla), expressed in tannic acid and
83 quercetin equivalents, ranged from 47877 to 73791 and from 1950 to 9400 mg/kg, respectively.

84 The mass concentration ratio of {[Fla]/[TPh]} varied from 0.04 to 0.13. The maximum
85 concentration of polyphenols was found in ER biomass.

86 The polyphenol contents in the examined weeds were lower than or comparable to the phenolic
87 contents reported by Březinová et al. [28] in other macrophytes (*Phragmites australis* Trin. ex
88 Steud., *Phalaris arundinacea* L., *Typha latifolia* L., *Glyceria maxima* Holmb., *Scirpus*
89 *sylvaticus* L., *Carex nigra* Reich. and *Juncus effusus* L.), in the 9020-28390 mg/kg interval. For
90 comparison purposes, the concentrations of total polyphenols and flavonoids in another set of
91 macrophytes (*A. pinnata*, *P. stratiotes*, *S. molesta*, *T. natans*, *P. maculosa* and *N. nucifera*) were
92 found to vary from 750 to 20800 and from 2420 to 11760 mg/kg, respectively [22].

93 |

Table 1. Concentration of polyphenols and trace elements in the three weeds under study, expressed in mg/kg (dw).

Weed	<i>Rorippa palustris</i>	<i>Euphorbia rothiana</i>	<i>Schoenoplectus articulata</i>
TPh	69623	73791	47877
Fla	5150	9400	1950
Mg	2286	2309	1550
Al	9319	1018	1123
P	1431	2321	1642
S	5129	3142	1672
Cl	2279	1725	9532
K	13038	17573	16537
Ca	13072	13086	3727
Rb	8	14	36
Ti	143	227	308
Cr	27	40	18
Mn	680	421	1452
Fe	19632	29486	8719
Co	67	44	1
Zn	30	37	5
As	3.5	1	1.5
Se	1	1	1
Sr	108	46	29
Mo	14	3	10
Ba	60	2	2
Pb	18	5	6

3.2. Trace Element Concentrations

The total concentration of 20 elements (P, S, Cl, As, Se, K, Rb, Mg, Ca, Sr, Ba, Al, Ti, Cr, Mn, Fe, Co, Zn, Mo and Pb), Σn_{20} , ranged from 46372 to 71501 mg/kg (**Table 1**). The maximum concentration corresponded to ER biomass.

The concentrations of P, K, Mg, Ca, Cr, Fe and Zn were in the 1431 to 2321, 13038 to 17573, 1550 to 2309, 3727 to 13086, 18 to 40, 8719 to 29486 and 5 to 37 mg/kg range, respectively. The maximum concentration of these elements was again observed in ER.

Contents of Cl, Rb, Ti and Mn varied from 1725 to 9532 mg/kg, from 8 to 36 mg/kg, from 143 to 308 mg/kg, and from 421 to 1452 mg/kg, respectively. Remarkable high concentrations of these elements were detected in SA biomass, suggesting their bioaccumulation as chlorine complexes.

The maximum concentration of the other elements: S, Al, As, Sr, Ba, Co, Pb and Mo was detected in RP biomass, probably due to their accumulation as sulfur compounds. The concentration of Se in the biomass of the three weeds was only detectable at trace levels (<1.0 mg/kg).

Similar bioaccumulation patterns have been reported for other macrophytes [11-22].

3.3. Toxicity

These weeds are used as food as well as in traditional medicine. Heavy metals were found to be bio-accumulated in the weeds. The permissible limits of As, Pb, Cr and Zn in medicinal plants are 1.0, 5.0, 2.3 and 20 mg/kg, respectively [29-30]. The concentrations of As and Pb were higher than the prescribed values in the biomass from two of the weeds (RP and SA), chromium concentration was above the prescribed limit in all weeds, and concentrations of Zn above 30 mg/kg were accumulated in RP and ER.

3.4. Correlation Coefficients

The correlation coefficients (r) matrix between the polyphenol contents and the studied trace elements (for average values across the three weeds) is shown in **Table 2**. The TPh and Fla contents showed a good correlation with K, Ti, Mn and Co, probably due to formation of complexes. The elements: S, Mg, Ca, Cr, Fe, Co and Zn had also high correlation, pointing to their accumulation as sulfur compounds. Similarly, a group of elements: Al, As, Sr, Mo, Ba and

Pb showed good correlation, expecting their accumulation as aluminum compounds. Elements: Cl, Rb, Ti and Mn showed good correlations, indicating their absorption as chloride complexes. Phosphorous showed good correlations with K, Cr and Fe, hinting at its role as a co-factor element.

Table 2. Correlation coefficients among the total polyphenol (TPh), flavonoid (Fla) and trace element content present in the biomass from *R. palustris*, *E. rothiana* and *S. articulata*.

	Mg	Al	P	S	Cl	K	Ca	Rb	Ti	Cr	Mn	Fe	Co	Zn	As	Sr	Mo	Ba	Pb	TPh	Fla
Mg	1.00																				
Al	0.47	1.00																			
P	0.32	-0.69	1.00																		
S	0.80	0.90	-0.31	1.00																	
Cl	-1.00	-0.43	-0.35	-0.78	1.00																
K	-0.27	-0.98	0.83	-0.79	0.24	1.00															
Ca	1.00	0.49	0.29	0.82	-1.00	-0.30	1.00														
Rb	-0.97	-0.66	-0.09	-0.92	0.96	0.49	-0.98	1.00													
Ti	-0.85	-0.87	0.24	-1.00	0.83	0.74	-0.86	0.95	1.00												
Cr	0.82	-0.12	0.80	0.33	-0.84	0.32	0.81	-0.67	-0.40	1.00											
Mn	-0.98	-0.27	-0.51	-0.66	0.98	0.06	-0.97	0.90	0.71	-0.93	1.00										
Fe	0.89	0.02	0.71	0.45	-0.91	0.19	0.88	-0.77	-0.52	0.99	-0.97	1.00									
Co	0.93	0.76	-0.06	0.97	-0.92	-0.61	0.94	-0.99	-0.98	0.56	-0.83	0.66	1.00								
Zn	0.98	0.30	0.48	0.68	-0.99	-0.09	0.98	-0.92	-0.74	0.91	-1.00	0.96	0.85	1.00							
As	0.30	0.98	-0.81	0.81	-0.27	-1.00	0.33	-0.51	-0.76	-0.29	-0.09	-0.16	0.63	0.12	1.00						
Sr	0.65	0.98	-0.52	0.97	-0.62	-0.91	0.67	-0.80	-0.95	0.10	-0.47	0.23	0.88	0.50	0.92	1.00					
Mo	-0.18	0.78	-0.99	0.44	0.22	-0.90	-0.16	-0.05	-0.37	-0.71	0.39	-0.61	0.19	-0.36	0.88	0.63	1.00				
Ba	0.48	1.00	-0.68	0.91	-0.44	-0.98	0.50	-0.67	-0.87	-0.10	-0.28	0.03	0.77	0.31	0.98	0.98	0.78	1.00			
Pb	0.41	1.00	-0.73	0.87	-0.38	-0.99	0.44	-0.61	-0.84	-0.17	-0.21	-0.04	0.72	0.24	0.99	0.96	0.82	1.00	1.00		
TPh	0.99	0.35	0.43	0.72	-1.00	-0.15	0.99	-0.94	-0.77	0.89	-1.00	0.94	0.88	1.00	0.18	0.55	-0.30	0.36	0.30	1.00	

Fla 0.84 -0.09 0.78 0.35 -0.86 0.30 0.82 -0.69 -0.42 1.00 -0.94 0.99 0.58 0.92 -0.27 0.12 -0.69 -0.08 -0.15 0.90 1.00

131

132 4. CONCLUSIONS

133 The polyphenols and mineral contents of three weeds that grow in wetlands in India (*Euphorbia*
 134 *rothiana*, *Rorippa palustris* and *Schoenoplectus articulata*) were characterized. Total polyphenol
 135 and flavonoid contents ranged from 47877 to 73791 and from 1950 to 9400 mg/kg, respectively,
 136 lower than (or comparable to) those reported for other macrophytes. Although the three species
 137 were found to feature high contents of nutrients along with Cl and Al, the concentrations of
 138 heavy metals beyond safety limits advice against their use for food purposes. Nonetheless, these
 139 species may hold promise to control water and soil alkalinity, hardness and metal toxicity. The
 140 biomass from *R. palustris* for Ca, Mg, Al, S, Co, As and Pb; that of *E. rothiana* for P, K, Cr, Fe
 141 and Zn; and the *S. articulata* for Cl, Rb, Ti and Mn can be used as a bioindicator for their
 142 contamination detection, respectively.

143 CONSENT

144 Not applicable.

145 ETHICAL APPROVAL

146 Not applicable.

147

148 References

- 149 1. Jain A, Roshnibala S, Kanjilal PB, Singh RS, Singh HB. Aquatic/semi-aquatic plants used in
 150 herbal remedies in the wetlands of Manipur. Indian Journal of Traditional Knowledge.
 151 2007;6(2):345–351. <http://hdl.handle.net/123456789/941>.
- 152 2. Kamal M, Ghaly AE, Mahmoud N, Côté R. Phytoaccumulation of heavy metals by aquatic
 153 plants. Environment International. 2004;29(8):1029–1039. [https://doi.org/10.1016/S0160-](https://doi.org/10.1016/S0160-4120(03)00091-6)
 154 4120(03)00091-6.
- 155 3. Chukwuka KS, Omotaya OE. Soil Fertility Restoration Potentials of Tithonia Green Manure
 156 and Water Hyacinth Compost on a Nutrient Depleted Soil in South Western Nigeria Using
 157 *Zea mays* L. as Test Crop. Research Journal of Soil Biology. 2009;1(1):20–30.
 158 <http://dx.doi.org/10.3923/rjsb.2009.20.30>.
- 159 4. Marzouk MM, Hussein SR, Elkhateeb A, Farid MM, Ibrahim LF, Abdel-Hameed ESS.
 160 Phenolic profiling of *Rorippa palustris* (L.) Besser (Brassicaceae) by LC-ESI-MS:
 161 Chemosystematic significance and cytotoxic activity. Asian Pacific Journal of Tropical
 162 Disease. 2016;6(8):633–63. [https://doi.org/10.1016/S2222-1808\(16\)61100-3](https://doi.org/10.1016/S2222-1808(16)61100-3).
- 163 5. Deepak P, Gopal GV. Nilgiris: A Medicinal Reservoir. The Pharma Innovation Journal.
 164 2014;3(8):73–79. http://www.thepharmajournal.com/vol3Issue8/Issue_oct_2014/8.1.pdf.
- 165 6. Rani KRV, Prakash SEL, Geetha K, Dhandapani B, Dhanabal K. Evaluation of
 166 pharmacognostic, phytochemical and antimicrobial activity of *Euphorbia rothiana*.
 167 Pharmacognosy Journal. 2009;1(1):22–24. [http://www.phcogfirst.com/sites/default/files/](http://www.phcogfirst.com/sites/default/files/PJ_1_1_Evaluation%20of%20Pharmacognostic.....pdf)
 168 PJ_1_1_Evaluation%20of%20Pharmacognostic.....pdf.
- 169 7. Roy DK, Sinha BK, Singh B. Less known uses of *Schoenoplectus articulatus* (L.) Palla of
 170 lower Assam. Indian Journal of Traditional Knowledge. 2010;9(3):599–600.

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

- 171 [https://www.niscair.res.in/Sciencecommunication/ResearchJournals/rejour/ijtk/Fulltextsearch/2010/July%202010/IJTK-Vol%209\(3\)-July%202010-pp%20599-600.htm](https://www.niscair.res.in/Sciencecommunication/ResearchJournals/rejour/ijtk/Fulltextsearch/2010/July%202010/IJTK-Vol%209(3)-July%202010-pp%20599-600.htm).
- 172
- 173 8. Hunashal RD, Bagewadi RM, Satyanarayana D. Physico-chemical and pharmacological
- 174 characteristics of *Euphorbia rothiana* seed oil. Journal of Chemical and Pharmaceutical
- 175 Research. 2011;3(5):124–129. [http://www.jocpr.com/articles/physicochemical-and-](http://www.jocpr.com/articles/physicochemical-and-pharmacological-characteristics-ofeuphorbia-rothiana-seed-oil.pdf)
- 176 [pharmacological-characteristics-ofeuphorbia-rothiana-seed-oil.pdf](http://www.jocpr.com/articles/physicochemical-and-pharmacological-characteristics-ofeuphorbia-rothiana-seed-oil.pdf).
- 177 9. Duman F, Cicek M, Sezen G. Seasonal changes of metal accumulation and distribution in
- 178 common club rush (*Schoenoplectus lacustris*) and common reed (*Phragmites australis*)
- 179 Ecotoxicology. 2007;16(6):457–463. <https://doi.org/10.1007/s10646-007-0150-4>.
- 180 10. Girdhar M, Sharma NR, Rehman H, Kumar A, Mohan A. Comparative assessment for
- 181 hyperaccumulatory and phytoremediation capability of three wild weeds. 3 Biotech.
- 182 2014;4(6):579–589. <https://doi.org/10.1007/s13205-014-0194-0>.
- 183 11. Girdhar M, Sharma NR, Rehman H, Kumar A, Mohan A. Comparative assessment for
- 184 hyperaccumulatory and phytoremediation capability of three wild weeds. 3 Biotech.
- 185 2014;4(6):579–589. <https://doi.org/10.1007/s13205-014-0194-0>.
- 186 12. Nematian MA, Kazemeini F. Accumulation of Pb, Zn, Cu and Fe in plants and
- 187 hyperaccumulator choice in Galali iron mine area, Iran. International Journal of Agriculture
- 188 and Crop Sciences. 2013;5(4):426–432. [https://pdfs.semanticscholar.org/f099/](https://pdfs.semanticscholar.org/f099/1c015a613dc9536b77c06c048191d6b52fbf.pdf)
- 189 [1c015a613dc9536b77c06c048191d6b52fbf.pdf](https://pdfs.semanticscholar.org/f099/1c015a613dc9536b77c06c048191d6b52fbf.pdf).
- 190 13. Hameed I, Dastagir G, Hussain F. Nutritional and elemental analyses of some selected
- 191 medicinal plants of the family polygonaceae. Pakistan Journal of Botany. 2008;40(6):2493–

2502. [https://www.researchgate.net/publication/260387195_Nutritional_and](https://www.researchgate.net/publication/260387195_Nutritional_and_elemental_analyses_of_some_selected_medicinal_plants_of_the_family_Polygonaceae)
 _elemental_analyses_of_some_selected_medicinal_plants_of_the_family_Polygonaceae.
14. Lee KS, Kwon YJ, Lee KY. Analysis of chemical composition, vitamin, mineral and
 antioxidative effect of the Lotus leaf. Journal of the Korean Society of Food Science and
 Nutrition. 2008;37(12):1622–1626. <http://dx.doi.org/10.3746/jkfn.2008.37.12.1622>.
15. Pascal L, Angela ML, Jaime EM, Jeimmy S, Carlos AB, Wolfgang BS, Andre B. Nutritional
 value of aquatic ferns (*Azolla filiculoides* Lam. and *Salvinia molesta* Mitchell) in pigs.
 Animal Feed Science and Technology. 2009;149(1-2):135–148. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.anifeedsci.2008.04.013)
 anifeedsci.2008.04.013.
16. Qin L, Zhenli LH, Donald AG, Peter JS, Xiaoe Y. Uptake and distribution of metals by
 water lettuce (*Pistia stratiotes* L.). Environmental Science and Pollution Research.
 2011;18(6):978–986. <https://doi.org/10.1007/s11356-011-0453-0>.
17. Babu M, Dwivedi DH, Lata R. Distribution of heavy metals in edible aquatic plant: water
 chestnut (*Trapa natans* var. *bispinosa* Roxb.). International Journal of Plant Sciences.
 2011;6(2):270–273. [http://www.researchjournal.co.in/online/IJPS/IJPS%206\(2\)/6_A-270-](http://www.researchjournal.co.in/online/IJPS/IJPS%206(2)/6_A-270-273.pdf)
 273.pdf.
18. Zhang J, Wang N, Zhang F. Analysis of Accumulating Ability of Heavy Metals in Lotus
 (*Nelumbo nucifera*) Improved by Ion Implantation. Plasma Science and Technology.
 2012;14(5):424–426. <https://doi.org/10.1088/1009-0630/14/5/21>.
19. Petrović D, Jančić D, Furdek M, Mikac N, Krivokapić S. Aquatic plant *Trapa natans* L. as
 bioindicator of trace metal contamination in a freshwater lake (Skadar Lake, Montenegro).
 Acta Botanica Croatica. 2016;75(2):236–243. <https://doi.org/10.1515/botcro-2016-0031>.

- 214 20. Anitha KC, Rajeshwari YB, Prasanna SB, Shilpa SJ. Nutritive evaluation of azolla as
215 livestock feed. *Journal of Experimental Biology and Agricultural Sciences*. 2016;4(6):670–
216 674. [http://dx.doi.org/10.18006/2016.4\(Issue6\).670.674](http://dx.doi.org/10.18006/2016.4(Issue6).670.674).
- 217 21. Verma A, Karki D, Meghwal S. Heavy metal extraction potential of various plant of
218 *Nelumbo nucifera*, An aquatic macrophyte in tropical freshwater systems. *International*
219 *Journal of Research and Analytical Reviews*. 2018;5(4):786–790.
220 <https://www.researchgate.net/publication/329117000>.
- 221 22. Sahu YK, Deb MK, Patel KS, Martín-Ramos P, Towett EK, Tarkowska-Kukuryk M.
222 Bioaccumulation of nutrients and toxic elements with macrophytes. *Journal of Toxic,*
223 *Hazardous and Radio Active Waste*. 2019; (In press).
- 224 23. Bowden WB, Glime JM, Riis T. Methods in stream ecology, *Ecosystem Structure*, Chapter
225 13 - Macrophytes and Bryophytes, 2017, p. 243–271. [https://doi.org/10.1016/B978-0-12-](https://doi.org/10.1016/B978-0-12-416558-8.00013-5)
226 [416558-8.00013-5](https://doi.org/10.1016/B978-0-12-416558-8.00013-5).
- 227 24. Bertaud F, Tapin-Lingua S, Pizzi A, Navarrete P, Petit-Conil M. Characterisation of
228 industrial barks for their tannin contents for further green-wood based adhesives
229 applications. *InTech Fibres COST FP0901-Hamburg*. 2010.
- 230 25. Singleton VL, Orthofer R, Lamuela-Raventos RM. Analysis of total phenols and other
231 oxidation substances by means of Folin-Ciocalteu reagent. *Methods in Enzymology*.
232 1999;299:152–178. [https://doi.org/10.1016/S0076-6879\(99\)99017-1](https://doi.org/10.1016/S0076-6879(99)99017-1).
- 233 26. Chang, CC, Yang MH, Wen HM, Chern JC. Estimation of total flavonoid content in
234 propolis by complementary colorimetric methods. *Journal of Food and Drug Analysis*.
235 2002;10(3):178–182. <https://www.fda.gov.tw/en/publishjfdalistContent.aspx?id=27>.

- 236 27. Towett EK, Shepherd KD, Drake BL. Plant elemental composition and portable X-ray
237 fluorescence (pXRF) spectroscopy: quantification under different analytical parameters. X-
238 Ray Spectrometry. 2016;45:117–124. <https://doi.org/10.1002/xrs.2678>.
- 239 28. Březinová TD, Vymazal J. Phenolic Compounds in Wetland Macrophytes. Plant Science.
240 2018;49(1):1–8. <https://doi.org/10.2478/sab-2018-0001>.
- 241 29. FAO/WHO. Joint FAO/WHO Food Standards Programme Codex Committee on
242 Contaminants in Foods Food. CF/5 INF/1. 2011. [http://www.fao.org/tempref/codex/](http://www.fao.org/tempref/codex/Meetings/CCCF/CCCF5/cf05_INF.pdf)
243 [Meetings/CCCF/CCCF5/cf05_INF.pdf](http://www.fao.org/tempref/codex/Meetings/CCCF/CCCF5/cf05_INF.pdf).
- 244 30. WHO. Quality control methods for medicinal plant materials. World Health Organization.
245 1998. <https://apps.who.int/iris/handle/10665/41986>.