

# Original Research Article

## **Mapping a climate change vulnerability index: An assessment in agricultural, geological and demographic sectors across the districts of Karnataka state (India)**

### **Abstract**

Climate change is a continuous phenomenon and over hundreds of years, the atmosphere has changed considerably around the world. Karnataka has the second largest drought prone area in the country next only to Rajasthan. Assessment of vulnerability index could play a major role in designing appropriate mitigation and adaptation policies to overcome the impacts of climate change. The vulnerability assessment is an exhaustive procedure determined by a large number of indicators. This study attempted to capture a picture of composite vulnerability index of different districts of Karnataka by considering agronomic, climatic and demographic indicators. The secondary data on climatic, agronomic and demographic factors were collected from various sources for the year 2017-18. The findings of the study as shown that the average vulnerability index for 30 districts is 0.577 and 16 districts placed above the average composite vulnerability index level. Bidar (0.655) is the most vulnerable district followed by Kolar (0.658) and Yadgir (0.638) districts. Shivamogga (0.440), Davanagere (0.486) and Udupi (0.486) districts exhibit the least vulnerability to changing climate. The results suggest that agricultural and climatic indicators are the major factors which influence vulnerability. So special attention should be given to agricultural and climatic sectors to minimize the impacts of climatic change in the most vulnerable districts.

**Key words:** Vulnerability index, Climate change, Per capita income, Sensitivity, Exposure and Adaptability

**Introduction:**

28 Agricultural economy in Karnataka is largely influenced by agro-climatic factors, water and  
29 other resource contributed by farmers, technology, infrastructure, tradition and social capital as  
30 also the market forces of demand and supply. Karnataka has the second largest drought prone  
31 area in the country next only to Rajasthan and water availability is one of the major concerns in  
32 the state. Karnataka's annual rainfall is 1,151 mm on an average, of which 80 per cent is received  
33 during the southwest monsoon, 12 per cent in the post monsoon period, 7 per cent during  
34 summer and 1 percent in rabi season. Groundwater potential of the area depends on rainfall and  
35 efforts to recharge. Change in climatic conditions directly affects the hydrological cycle and  
36 gradually the groundwater table. Obviously the economic impact of climate change will severely  
37 affect the food security as well as livelihood security including health security of farmers  
38 (Chandrakanth, 2015).

39 Climate change is a continuous phenomenon and over hundreds of years, the atmosphere  
40 has changed considerably around the world. However, the pace and pattern of changes in  
41 climatic factors in recent decades have turned into a matter of concern. Especially, since it is  
42 very hard to comprehend the effect of change in climatic factors at the small scale level even,  
43 say, at block or district levels (Raju *et al.*, 2017). The Intergovernmental Panel on Climate  
44 Change (IPCC), in its second evaluation report (Anonymous, 1996), characterizes vulnerability  
45 as the degree to which environmental change may harm or damage a system. It infers that  
46 vulnerability not only depends on a system of sensitivity, but also in addition, on its capacity to  
47 adjust to new climatic conditions, the level of economic development and institutions.

48 It is well known that poor people in the least developed nations are the most vulnerable  
49 against the effects of anthropogenic environmental change (Stern *et al.*, 2006). The poor are  
50 antagonistically affected by the environmental change since they live in vigorously affected  
51 nations and areas inside those nations, rely upon natural resource-based livelihood that are  
52 lopsidedly influenced by climate change.

53 People who live in the semi-arid and arid region, in low-lying seaside regions, in water-  
54 restricted or flood-inclined zones or on little islands are especially vulnerable to environmental  
55 change (Watson *et al.*, 1996). Obviously climate change will, in many parts of the world,  
56 antagonistically influence socio-economic status, including water resources, farming, forestry,

57 fisheries and human settlements, natural resources and human wellbeing with creating nations  
58 being the most vulnerable (IPCC, 2001).

59 There is a huge demand to create indicators of vulnerability and of adaptive capacity to  
60 decide the robustness of methodologies over time (Adger *et al.*, 2004). At the district level,  
61 vulnerability appraisals add to setting development needs and monitoring progress. Sectoral  
62 evaluations give details and focus to key improvement plans. In Karnataka, farmers and  
63 agriculture workers constitute 56 per cent of the aggregate workforce (Government of Karnataka,  
64 2005) and this is viewed as one of the main thrusts in deciding the vulnerabilities of farming  
65 families in Karnataka.

## 66 2. Methodology:

67 The key target of this assessment is to analyse the climate vulnerability of different  
68 sectors across the districts of Karnataka (Fig 1). Keeping in view of this appraisal the  
69 information relating to different indicators pertaining to agriculture year 2013-14 to 2017-18  
70 were collected from various sources such as Karnataka State Natural Disaster Monitoring Centre  
71 (KSNDMC), Directorate of Economics and Statistics (DES) and Central Groundwater Board  
72 (CGB).

73 The vulnerability assesment is an exhaustive procedure influenced by a large number of  
74 indicators. However only the most significant and appropriate indicators were chosen for  
75 calculation of vulnerability index based on exposure, sensitivity and adaptability to varied  
76 climate. Parameters used in this study include

77 Climatic components: Variance of annual rainfall ( $\text{mm}^2$ ), Variance of South-West monsoon  
78 ( $\text{mm}^2$ ), Variance of maximum temperature, Variance of minimum temperature and Variance of  
79 average temperature.

80 Agricultural Components: Geographical area (GA) (ha), Forest area (% of GA), Area under food  
81 crops (% of Gross Cropped Area(GCA)), Net sown area (% of GA), Livestock population (No.  
82 per ha of GCA), Irrigated area (% of GCA), Cropping intensity (%), Productivity of major crops  
83 (Paddy, Ragi, Jowar, Sugarcane, Maize, Groundnut, Sunflower, Cotton, Arecanut, Coconut,  
84 Redgram, Cowpea, Chilli), Depth of Groundwater (meter below ground level), Per capita income  
85 (Rs per person).

86 Demographic components: Density of male population (Persons per sq. ha of GA), Density of  
 87 female population (Persons per sq. ha of GA), Literacy rate of male (%) and Literacy rate of  
 88 female (%).

89 Composite Vulnerability Index (CVI) is assessed for each district by using Iyenger and  
 90 Sudarshan (1982) technique for unequal weight. The assessed CVI is a total of three sub-sectors  
 91 specifically Climatic Vulnerability, Agriculture Vulnerability and Demographic Vulnerability.  
 92 Development of vulnerability index and Composite Vulnerability Index comprises of several  
 93 steps.

94 Step 1: The information compiled pertaining to three components was transformed into suitable  
 95 estimation units and arranged in a rectangular matrix with rows representing districts and  
 96 columns representing indicators.

97 Step 2: Since every one of the sub-component is measured using different units and scale, they  
 98 need to normalized first. The procedure developed by Anand and Sen (1994) for construction of  
 99 the Human Development Index (HDI) is used to normalize indicators. In any case, before doing  
 100 normalization, it is imperative to distinguish the functional relationship between the indicators  
 101 and vulnerability. Two kinds of practical relationships, vulnerability increases with the increase  
 102 (decrease) in the value of indicators are conceivable.

103 **Table 1: List of indicators and their functional relationship with vulnerability**

Components	Indicators	Functional relationship	Reference
Demographic	Density of population	Direct (↑)	Palanisami <i>et al.</i> , 2009
	Literacy rate	Inverse (↓)	Palanisami <i>et al.</i> , 2009
Climatic	Variance of rainfall	Direct (↑)	Ravindranath <i>et al.</i> , 2011
	Variance of Temperature	Direct (↑)	Ravindranath <i>et al.</i> , 2011
Agricultural	Productivity of major crops	Inverse (↓)	Hiremath and Shiyani, 2013
	Cropping intensity	Inverse (↓)	Hiremath and Shiyani, 2013
	Irrigated area	Inverse (↓)	Hiremath and Shiyani, 2013
	Forest area	Inverse (↓)	Hiremath and Shiyani, 2013
	Net sown area	Inverse (↓)	Hiremath and Shiyani, 2013
	Livestock population	Inverse (↓)	Hiremath and Shiyani, 2013

	Geographical area	Inverse (↓)	Palanisami <i>et al.</i> , 2009
	Depth of Groundwater	Direct (↑)	Suresh <i>et al.</i> , 2016
	Per capita income	Inverse (↓)	Suresh <i>et al.</i> , 2016

104

105 For direct relationship:

$$Y_{ij} = \frac{X_{ij} - \text{Min}(X_{ij})}{\text{Max}(X_{ij}) - \text{Min}(X_{ij})}$$

106

107 For indirect relationship:

$$Y_{ij} = \frac{\text{Max}(X_{ij}) - X_{ij}}{\text{Max}(X_{ij}) - \text{Min}(X_{ij})}$$

108

109 Where,

110

111  $Y_{ij}$  is the normalized value

112  $X_{ij}$  is the actual value of the indicator

113  $\text{Min}(X_{ij})$  and  $\text{Max}(X_{ij})$  are the minimum and maximum actual values

114 Step 3: The degree of vulnerability ( $\bar{y}_i$ ) is assumed to be the linear sum of  $X_{ij}$  as

$$\bar{y}_i = \sum_{j=1}^k w_j Y_{ij}$$

115 Where  $w_j$ 's are weights and are determined by

$$w_j = \frac{c}{\sqrt{\text{var}(Y_{ij})}}$$

116 Where  $c$  is the normalizing constant

$$c = \left[ \sum_{j=1}^k \frac{1}{\sqrt{\text{var}(Y_{ij})}} \right]^{-1}$$

117

118 The vulnerability index lies in the range of 0 and 1. A value of 1 indicates greatest vulnerability

119 and 0 shows absence of vulnerability.

120

### 121 3. Results and Discussion:

122 The Sector wise vulnerability indices and composite index were constructed for all the 30

123 districts of Karnataka. The districts were ranked based on extent of vulnerability index.

#### 124 3.1 Component wise vulnerability index

### 125 **3.1.1 Climatic Vulnerability index**

126 To construct district level vulnerability index five climatic variables were used and the  
127 results are presented in the Table 2. The results show that the Kalaburagi district has the highest  
128 climate vulnerability index of 0.747 followed by Kolar (0.720), Bidar (0.720), Raichur (0.712)  
129 and Yadgir (0.711) districts. The districts of Kodagu and Udupi have only 0.278 and 0.215  
130 vulnerability index respectively, the least in Karnataka state. We can observe highest  
131 vulnerability index values in northern districts of Karnataka which is due to large variations in  
132 rainfall and temperature during the year. These are the key determinant indicators which explain  
133 high climatic fluctuations among districts.

134

135 For instance, Prevalence of a high degree of anticipated change in mean precipitation and  
136 high inconsistency in minimum and maximum temperatures drove Kalaburagi district to the top  
137 of the chart.

### 138 **3.1.2 Agriculture Vulnerability index**

139 Based on functional relationship of the indicators, Vulnerability index for agricultural parameters  
140 were calculated for each district and is presented in Table 3.

141 Kodagu district secures first place with a total vulnerability index value of 0.787 followed  
142 by Bidar (0.761), Kolar (0.741) and Chitradurga (0.732) districts. Whereas Davanagere has been  
143 rated as least vulnerable district (0.524). Lower productivity, declined forest area, high  
144 groundwater table level, lower cropping intensity and low per capita income are the major  
145 factors which influence the high level of sensitivity leading to higher vulnerability index.

146 In general Kodagu, Bidar, Kolar and Chitradurga districts are most sensitive districts and  
147 highly vulnerable to climate change. On the contrary, Davanagere, Shivamogga, Bellary and  
148 Bengaluru Urban districts are less sensitive and least vulnerable to changing climate.

### 149 **3.1.3 Demographic vulnerability index**

150 The districts having high population density coupled with a lower rate of literacy were  
151 identified as vulnerable districts with respect to demographic features.

152 Bengaluru Urban (0.579) district occupied the first place whereas Dakshina Kannada  
153 (0.039) district is placed in the last position with respect to demographic vulnerability (Table 4).  
154 Yadgir (0.449), Raichur (0.353), Chamarajnaraga (0.335) and Kalaburagi (0.294) are the districts

155 having higher degree of vulnerability index next to Bengaluru Urban district. The coastal  
156 districts of Dakshina Kannada, Udupi (0.051) and Uttara Kannada (0.055) are having lower  
157 vulnerability index and higher adaptive capacity to changing climate because of high literacy rate  
158 and lower population density.

### 159 **3.2 Composite vulnerability index**

160 Agricultural indicators, climatic indicators and demographic indicators were used to  
161 construct composite vulnerability index. Table 5 shows district wise composite vulnerability  
162 index which is calculated using all the three sub-components (Agricultural, Climatic and  
163 Demographic). Average composite vulnerability index for 30 districts is 0.584 and 17 districts  
164 placed above the average composite vulnerability index level. Districts having high composite  
165 vulnerability index will be highly vulnerable to climate change. Bidar (0.577) district is having  
166 the highest composite vulnerability index followed by Kolar (0.658) and Yadgir (0.638). These  
167 districts are most vulnerable districts and the results are inline with the report submitted by  
168 Anonymous (2011) which used composite vulnerability index. They reported that Kalaburagi  
169 and Dakshina Kannada districts were the most and the least vulnerable districts, respectively.  
170 Higher composite index is observed mainly due to higher sensitivity of agricultural sector and  
171 larger exposure to climate change. Composite vulnerability index is lower for Shivamogga  
172 (0.440), Davanagere (0.486) and Udupi (0.486) districts because these districts are showing less  
173 vulnerability in terms of agriculture and climatic indicators. In addition also demographic  
174 variables such as population density and literacy rate have contributed to lowering of composite  
175 vulnerability index . At district level, contribution of each sub-component to composite index is  
176 not uniform. In general agricultural indicators contributed foremost, followed by climatic and  
177 demographic indicators. A study conducted by Hiremath and Shiyani (2013) reported that  
178 agriculture and occupation sector were the major sectors which have contributed most to  
179 composite vulnerability index in Saurashtra.

180

### 181 **4. Conclusion:**

182 Karnataka is the second most drought prone state after Rajasthan. District wise  
183 vulnerability mapping was carried out to calculate the vulnerability index of each district. Sector  
184 wise indicators were selected based on exposure, sensitivity and adaptive capacity to climate  
185 change. All the indicators were considered to calculate composite vulnerability index. Findings

186 of the analysis shows that Bidar is the most vulnerable district and Shivamogga is the least  
187 vulnerable. Major component which is contributing to composite index is the Agricultural  
188 vulnerability. The results of agricultural vulnerability index analysis has highlighted the  
189 indicators such as productivity of the major crops, cropping intensity and per capita income are  
190 the major drivers in determining the vulnerability of districts. Therefore, it is suggested that  
191 Bidar, Kolar, Yadgir, Koppal and Chtradurga districts should be considered under on priority to  
192 minimize degree of vulnerability. There is a need to take up adaptive practices such as varietal  
193 selection according to prevailing weather, contingent cropping, soil and water conservation  
194 measures, in-situ moisture conservation, rainwater harvesting and augmenting recharging of  
195 groundwater for supplementary irrigation. In addition, better education and infrastructure  
196 development in rural areas will also play a catalytic role in enhancing adaptive capacity of these  
197 districts.

198 Ethical: NA

199 Consent: NA

## 200 5. References:

201 Adger, W., Neil., Brooks, Nick, Bentham, Graham, Agnew, Maureen, and Eriksen, S., 2004.  
202 New indicators of vulnerability and adaptive capacity (Technical Report 7). Norwich,  
203 UK: Tyndall Centre for Climate Change Research.

204 Anand, S. and Sen, A., 1994, Human development index : Methodology and measurement.  
205 United Nations Development Programme, New York : Human Development Report  
206 Office.

207 Anonymous, 1996, Climate change second assessment report (1995). Intergovernmental panel on  
208 climate change. Geneva, Switzerland.

209 Anonymous, 2011, Karnataka Climate Change Action Plan, Bangalore Climate Change  
210 Initiative–Karnataka, Bengaluru (India) Final Report, Government of Karnataka.

211 Chandrakanth, M. G., 2015, *Water resource economics: Towards sustainable use of water for*  
212 *irrigation in India*, Springer, Switzerland.



213 Government of Karnataka. 2005, Human development report, Karnataka. Bangalore:Government  
214 of Karnataka.

215 Hiremath, D. B. and Shiyani, R. L., 2013, Analysis of vulnerability indices in various agro-  
216 climatic zones of Gujarat. *Indian Journal of Agricultural Economics*, **68**(1):122-137.

217 IPCC, 2001. Climate change 2001: The scientific basis—Contribution of working group I to the  
218 third assessment report of the intergovernmental panel on climate change. Cambridge,  
219 UK: Cambridge University Press.

220 Iyengar, N. S. and Sudharshan, P., 1982, A method of classifying regions from multivariate data,  
221 *Economic and Political Weekly, Special Article* : 2048-1052.

222 Palanisami, K., Paramasivam, P., Ranganathan, C.R., Aggarwal, P.K., and Senthilnathan, S.  
223 (2009). Quantifying vulnerability and impact of climate change on production of major  
224 crops in Tamil Nadu, India. In: Taniquchi, M., Burnett, W.C., Fukushima, Y., Haigh, M.,  
225 and Y. Umezawa, From headwaters to the ocean: Hydrological changes and watershed  
226 management. London: Taylor and Francis Group.pp.509-521.

227 Raju, K. V., Deshpande, R. S. and Satyasiba Bedamatta, 2017, Vulnerability to Climate Change:  
228 A Sub-regional Analysis of Socio-economic and Agriculture Sectors in Karnataka, India,  
229 *Journal of Development Policy and Practice*, **2**(1):24–55.

230 Ravindranath, N.H., Rao, S., Sharma, N., Nair, M., Gopalakrishnan, Rao, A.S. and Baka, G.,  
231 2011, Climate change and vulnerability profiles for north-east India. *Current Science*,  
232 **101**(3): 384-394.

233 Stern, N., Peters, S., Bakhshi, V., Bowen, A., Cameron, C., Catovsky, S. and Dietz, S., 2006,  
234 *Stern review: The economics of climate change*. Oxford, UK: OUP.

235 Suresh, K., Biswas, H., Raizada, A. and Srinivas, S., 2016, Assesment of vulnerability to climate  
236 change:A case study of Karnataka, *Journal of Soil and Water Conservation*, Vol. 44, No.  
237 **3**, pp 314-320.

238 Watson, R.T., Zinyoera, M.C. and Moss, R.H., 1996, The regional impact of climate change: An  
 239 assessment of vulnerability (A Special Report of IPCC Working Group II).Cambridge,  
 240 UK: Cambridge University Press.

241

242 **Tables**

243 **Table 2: Index of climate vulnerability across the various districts of Karnataka**

Sl. No	Districts	Annual rainfall	S-W monsoon	Max Temp	Min Temp	Avg Temp	Index total
1	KALABURAGI	0.177	0.174	0.102	0.169	0.125	0.747
2	KOLAR	0.170	0.183	0.189	0.107	0.073	0.720
3	BIDAR	0.173	0.169	0.120	0.189	0.069	0.720
4	RAICHUR	0.183	0.183	0.082	0.127	0.138	0.712
5	YADGIR	0.181	0.177	0.088	0.130	0.135	0.711
6	VIJAYAPURA	0.185	0.182	0.087	0.147	0.100	0.701
7	RAMANAGARA	0.163	0.178	0.165	0.085	0.103	0.693
8	BALLARI	0.186	0.187	0.100	0.054	0.154	0.681
9	KOPPALA	0.183	0.184	0.063	0.096	0.141	0.667
10	BAGALKOTE	0.188	0.185	0.078	0.109	0.095	0.656
11	DHARWAD	0.181	0.179	0.118	0.094	0.070	0.643
12	DAVANAGERE	0.178	0.179	0.066	0.046	0.125	0.593
13	GADAG	0.189	0.186	0.061	0.075	0.082	0.593
14	CHITRADURGA	0.183	0.185	0.060	0.044	0.119	0.591
15	CHIKKABALLAPURA	0.176	0.183	0.055	0.077	0.094	0.585
16	BELAGAVI	0.174	0.166	0.058	0.103	0.083	0.584
17	TUMAKURU	0.176	0.181	0.044	0.055	0.105	0.561
18	HAVERI	0.179	0.174	0.072	0.051	0.082	0.559
19	MANDYA	0.172	0.185	0.038	0.026	0.116	0.537
20	MYSURU	0.173	0.182	0.023	0.036	0.101	0.514
21	CHAMARAJANAGARA	0.172	0.189	0.000	0.028	0.086	0.475
22	UTTARA KANNADA	0.078	0.074	0.139	0.077	0.106	0.474
23	BENGALURU RURAL	0.164	0.175	0.023	0.052	0.056	0.470
24	BENGALURU URBAN	0.157	0.168	0.015	0.038	0.055	0.431
25	HASSAN	0.152	0.149	0.024	0.043	0.052	0.421
26	CHIKKAMAGALURU	0.128	0.126	0.076	0.039	0.042	0.412
27	SHIVAMOGGA	0.103	0.082	0.053	0.058	0.076	0.372
28	DAKSHINA KANNADA	0.031	0.030	0.045	0.000	0.189	0.294
29	KODAGU	0.080	0.082	0.094	0.022	0.000	0.278
30	UDUPI	0.000	0.000	0.024	0.022	0.170	0.215

244

245

246

247

248

249 **Table 3: Agricultural vulnerability index across the districts of Karnataka**

<b>District</b>	<b>Geographical area(Ha)</b>	<b>Forest area(% to GA)</b>	<b>Total food crops(% to GCA)</b>	<b>Net sown area(% to GA)</b>	<b>Livestock pon(No. per Ha of GCA)</b>
Kodagu	0.008	0.032	0.055	0.027	0.065
Bidar	0.013	0.051	0.026	0.016	0.063
Kolar	0.007	0.051	0.012	0.027	0.057
Chitradurga	0.026	0.048	0.031	0.024	0.059
Koppal	0.014	0.051	0.018	0.010	0.062
Hassan	0.019	0.048	0.022	0.021	0.061
Gadag	0.010	0.049	0.020	0.001	0.064
Dakshin Kannada	0.011	0.036	0.013	0.034	0.063
Dharwad	0.008	0.049	0.023	0.003	0.064
Haveri	0.011	0.048	0.021	0.006	0.062
Chikballapura	0.008	0.046	0.016	0.024	0.058
Bengaluru Rural	0.000	0.051	0.026	0.022	0.061
Kalaburagi	0.036	0.052	0.006	0.009	0.064
Raichur	0.026	0.053	0.014	0.018	0.061
Tumkuru	0.035	0.051	0.037	0.024	0.059
Mysuru	0.017	0.047	0.021	0.016	0.062
Chamarajanagara	0.014	0.021	0.017	0.034	0.061
Yadgir	0.013	0.050	0.028	0.014	0.061
Ramanagara	0.005	0.041	0.019	0.024	0.060
Chikkamagaluru	0.021	0.035	0.032	0.027	0.063
Vijayapura	0.034	0.054	0.003	0.000	0.064
Bagalkot	0.018	0.046	0.005	0.010	0.060
Uttar Kannada	0.033	0.000	0.004	0.044	0.059
Udupi	0.006	0.035	0.015	0.035	0.061
Mandya	0.011	0.051	0.010	0.026	0.058
Belagavi	0.046	0.045	0.014	0.016	0.065
Bellari	0.026	0.046	0.019	0.020	0.057
Bengaluru Urban	0.000	0.053	0.017	0.041	0.000
Shivamogga	0.026	0.032	0.000	0.035	0.060
Davanagere	0.015	0.044	0.005	0.013	0.062

250

251

252  
 253  
 254  
 255  
 256  
 257  
 258

Contd....

<b>Irrigated area(% to GCA)</b>	<b>Cropping intensity (%)</b>	<b>Productivity</b>	<b>Per capita income</b>	<b>Depth of groundwater (mbgl)</b>	<b>Index total</b>
0.049	0.045	0.443	0.046	0.018	<b>0.787</b>
0.041	0.034	0.438	0.051	0.030	<b>0.761</b>
0.038	0.056	0.442	0.045	0.006	<b>0.741</b>
0.032	0.037	0.410	0.048	0.017	<b>0.732</b>
0.027	0.040	0.426	0.050	0.012	<b>0.708</b>
0.032	0.037	0.378	0.042	0.038	<b>0.697</b>
0.035	0.029	0.402	0.046	0.038	<b>0.694</b>
0.012	0.041	0.449	0.014	0.011	<b>0.684</b>
0.041	0.000	0.429	0.041	0.024	<b>0.683</b>
0.028	0.039	0.394	0.048	0.020	<b>0.677</b>
0.029	0.048	0.381	0.046	0.012	<b>0.669</b>
0.034	0.057	0.358	0.037	0.020	<b>0.667</b>
0.040	0.041	0.332	0.051	0.033	<b>0.663</b>
0.023	0.039	0.379	0.049	0.000	<b>0.663</b>
0.025	0.044	0.345	0.040	0.003	<b>0.662</b>
0.026	0.024	0.371	0.045	0.030	<b>0.661</b>
0.019	0.037	0.392	0.044	0.017	<b>0.657</b>
0.023	0.034	0.379	0.051	0.003	<b>0.655</b>
0.034	0.053	0.352	0.039	0.028	<b>0.654</b>
0.036	0.035	0.353	0.029	0.021	<b>0.651</b>
0.025	0.053	0.348	0.050	0.018	<b>0.649</b>
0.014	0.036	0.378	0.041	0.037	<b>0.646</b>
0.024	0.050	0.364	0.043	0.021	<b>0.644</b>
0.027	0.045	0.386	0.024	0.007	<b>0.641</b>
0.005	0.033	0.356	0.039	0.039	<b>0.628</b>
0.011	0.027	0.325	0.049	0.025	<b>0.623</b>
0.014	0.032	0.297	0.040	0.013	<b>0.564</b>
0.029	0.049	0.361	0.000	0.003	<b>0.552</b>
0.000	0.047	0.266	0.036	0.025	<b>0.527</b>
0.012	0.049	0.267	0.048	0.010	<b>0.524</b>

259

260  
261  
262  
263  
264  
265  
266  
267

**Table 4: Demographic vulnerability index across the districts of Karnataka**

<b>District</b>	<b>Density of male population</b>	<b>Density of female population</b>	<b>Literacy rate of male (%)</b>	<b>Literacy rate of female (%)</b>	<b>Index total</b>
Bengaluru Urban	0.281	0.281	0.015	0.001	0.579
Yadgir	0.006	0.006	0.225	0.212	0.449
Raichur	0.006	0.006	0.165	0.176	0.353
Chamarajanagara	0.003	0.003	0.184	0.145	0.335
Kalaburagi	0.007	0.007	0.137	0.144	0.294
Bellari	0.010	0.010	0.120	0.129	0.270
Vijayapura	0.005	0.005	0.116	0.136	0.262
Ramanagara	0.011	0.012	0.119	0.112	0.255
Koppal	0.007	0.008	0.106	0.132	0.253
Bagalkot	0.010	0.010	0.101	0.128	0.249
Mandya	0.015	0.016	0.108	0.107	0.246
Chikkaballapura	0.011	0.011	0.112	0.112	0.245
Bidar	0.012	0.012	0.102	0.112	0.238
Mysuru	0.022	0.023	0.107	0.085	0.237
Belagavi	0.014	0.015	0.080	0.097	0.206
Kolar	0.016	0.017	0.083	0.086	0.202
Chitradurga	0.004	0.004	0.086	0.091	0.184
Davanagere	0.013	0.013	0.078	0.075	0.179
Tumakuru	0.008	0.008	0.075	0.083	0.174
Gadag	0.006	0.006	0.062	0.093	0.167
Bengaluru Rural	0.020	0.020	0.061	0.067	0.167
Hassan	0.008	0.009	0.069	0.077	0.163
Haveri	0.013	0.013	0.067	0.068	0.160
Dharwad	0.019	0.020	0.049	0.053	0.142
Chikkamagaluru	0.001	0.002	0.056	0.054	0.114
Shivamogga	0.005	0.005	0.052	0.046	0.107
Kodagu	0.000	0.000	0.043	0.030	0.073
Uttara Kannada	0.000	0.000	0.026	0.028	0.055
Udupi	0.011	0.014	0.013	0.013	0.051
Dakshina Kannada	0.018	0.021	0.000	0.000	0.039

268

269

270

271

272 **Table 5: Composite index of vulnerability**

Sl. No	Districts	Composite index	Sl. No	Districts	Composite index
1	BIDAR	0.677	16	CHAMARAJANAGAR	0.579
2	KOLAR	0.658	17	MYSURU	0.574
3	YADGIR	0.638	18	TUMKUR	0.573
4	KOPPAL	0.636	19	HASSAN	0.571
5	RAICHUR	0.628	20	BENGALURU RURAL	0.558
6	CHITRADURGA	0.628	21	MANDYA	0.557
7	KALABURAGI	0.625	22	BELAGAVI	0.555
8	RAMANAGARA	0.604	23	BALLARI	0.543
9	VIJAYAPURA	0.602	24	BENGALURU URBAN	0.538
10	GADAG	0.599	25	CHIKKKAMAGALURU	0.531
11	DHARWAD	0.596	26	UTTARA KANNADA	0.530
12	KODAGU	0.594	27	DAKSHINA KANNADA	0.528
13	CHIKBALLAPUR	0.593	28	UDUPI	0.486
14	BAGALKOT	0.590	29	DAVANGERE	0.486
15	HAVERI	0.580	30	SHIVAMOGGA	0.440
					<b>Average=0.577</b>

273

274

275

276

277

278

279

280

281


282

283 **Fig 1: Map showing Karnataka state in India**

UNDER PEER REVIEW

India

 Karnataka



284

Created with mapchart.net ©

285

286



287 **Fig:2 Climate Vulnerability of different districts of Karnataka (India)**

288  
289  
290  
291  
292

