

1 **Evaluation of Different Nutritional and Soil Sources **Fertilizers** on the Early**  
2 **Growth of *Moringa oleifera* (Lam)**

3  
4 **ABSTRACT**

5 In any plantation establishment programme **there should** be adequate number of healthy  
6 seedlings and **this depends** on the nutrition and care given to them at the nursery stage.  
7 Pot experiment was conducted to assess the response of *Moringa oleifera* (Lam)  
8 seedlings to different soil and nutritional sources at nursery stage. One hundred seeds  
9 were obtained from Centre for Environmental Renewable Resources Research and  
10 Development (CENRAD) Ibadan, Nigeria and sown in germination trays. Seed  
11 germination was completed between 10-15 days, 36 uniformly growing seedlings were  
12 transplanted into polythene pots of size 29×25cm and were filled with different soil  
13 sources at 500gm (arable) soil, forest reserve soil and natural forest mixed with  
14 nutritional sources of the same ratio (10gm) i.e. poultry manure, cow dung and N.P.K  
15 15.15.15. The experiment was factorial arranged in a completely randomized design.  
16 Result showed that pots with natural forest soil mixed with 10gm N.P.K ( $s_3f_3$ ) produced  
17 seedlings with highest value In plant height (76.30cm), stem diameter (3.47mm) and  
18 number of branches (10.00) which was significantly different ( $p < 0.01$ ) from the other  
19 treatments. The least value in plant height (30.70cm) stem diameter (1.50mm) and  
20 number of branches (3.3) were recorded in pots with forest reserve soil mixed with cow  
21 dung ( $s_2f_2$ ). Interaction effect of soil and nutritional sources were significant for all the  
22 growth parameters assessed ( $p < 0.01$ ). **Results showed that Natural forest soils treated**  
23 **with N.P.K 15.15.15 could be** employed in raising *Moringa oleifera* seedlings at nursery  
24 stage for optimum performance.

25 *Keywords: Moringa oleifera, growth, nutritional sources, different soil type*

26 **1. INTRODUCTION**

27 The forest plays an important role in protecting the soil, ameliorating the environment  
28 and protecting water resources. Non-timber forest products are very essential in urban  
29 and rural life, under which *Moringa oleifera* (Lam) belong (Leone *et al.*, 2015). *Moringa*  
30 *oleifera* is the most widely cultivated species of the genus *Moringa*, is the only genus in  
31 the family Moringaceae. English common names include: Moringa, (Olson, 2010)  
32 drumstick tree (from the appearance of the long, slender, triangular seed pods),  
33 horseradish tree (from the taste of the roots, which resembles horseradish, ben oil tree, or  
34 benzoil tree (from the oil which is derived from the seeds). It is a fast-growing, drought-  
35 resistant tree, native to the southern foothills of the Himalayas in the northwestern Indian,  
36 and widely cultivated in tropical and subtropical areas where its young seed pods and  
37 leaves are used as vegetables. It can also be used for water purification and washing, and  
38 is sometimes used in herbal medicine (Torondel, *et al.*, 2014).

39 It has become a clear issue that man cannot sufficiently sustain its existence without  
40 adequately improving the level of food and fiber production as raw material for industrial  
41 uses. Most tropical soils are deficient in nitrogen and other macronutrients and uptake of  
42 these limited quantities of nutrients by plant roots from litter alone is insufficient (Jose,  
43 2003). Nitrogen allows plants to produce proteins needed to build living tissues for green  
44 stems, leaves and strong roots, phosphorus helps move energy throughout the plant while  
45 potassium aids plants in assimilating sugars needed for growth. The application of  
46 fertilizers is the only way to supply nutrients within a short period of time. Adegbidi et al,  
47 (2003) reported that the effects of the mixed use of chemical fertilizer and organic matter  
48 on the growth of trees and soil fertility vary substantially according to the fertilizer  
49 amounts and the organic manure characteristics. The need to investigate the response of  
50 *Moringa oleifera* seedlings to different ratios of inorganic and organic fertilizer  
51 application on soil sources is essential as this will determine its optimum growth  
52 performance at the nursery stage. The objective of this study was thus; to investigate the  
53 effect of different nutritional and soil sources of fertilizers on early growth of *M. oleifera*  
54 so as to find the optimum dose of fertilizer for raising quality seedlings.

## 55 2. MATERIALS AND METHODS

56 This experiment was carried out at the West African Hardwood Improvement Project  
57 (WAHIP) of the Forestry Research Institute of Nigeria (FRIN), Ibadan (Latitude 7°39'13"  
58 and longitude 3°8'28"E. The Institute is situated at Jericho Hills in Ibadan North West  
59 Local Government Area of Oyo State. The climate of the area is tropically dominated by  
60 rainfall pattern ranging between 1400 - 1500mm, average temperature is 30°C. It has 2  
61 distinct seasons rainy season (April – October) and dry season (November – March)  
62 (FRIN, 2015). The black polythene pots were purchased from CENRAD, Ibadan. Cow  
63 dung and poultry manure were collected from Federal College of Forestry Teaching and  
64 Research Farm Ibadan. N. P. K 15.15.15 fertilizer was obtained from Centre for  
65 Environmental Renewable Resources and Management Development (CENRAD),  
66 Jericho, Ibadan. The natural forest soil was collected from FRIN arboretum, forest  
67 reserve soil was collected from *Pinus caribea* Morelet plantation FRIN while the  
68 cultivated soil used was collected from Federal College of Forestry Ibadan farm. One  
69 hundred *Moringa oleifera* seeds were obtained from CENRAD and sown in germination  
70 trays filled with sterilized river sand.

71 Cow dung and poultry manure were dried, crushed and sieved with 2 mm diameter mesh  
72 sieve while the soil samples were also sieved. The same ratio of cow dung, poultry  
73 manure and fertilizer (N.P.K. 15.15.15) were measured in grams (10 gm each) and mixed  
74 with the soil samples, each treatment contained the same level of organic and inorganic  
75 fertilizers. 10gm of organic fertilizer, (cow dung and poultry manure of same ratio) and  
76 10gm of N.P.K. 15.15.15 were weighed in the soil laboratory of Forestry Research  
77 Institute of Nigeria. Cow dung and also 10 mg of poultry manure were applied on 36  
78 seedlings i.e. cow dung and another 18 seedlings and 18 seedlings contained 10 gm of  
79 poultry manure, while other 18 seedlings contained 10 gm of N.P.K. 15.15.15. Eighteen  
80 seedlings were used as control (without fertilizer). A total of 72 seedlings were  
81 transplanted after four weeks into polythene pots of 29cm×25cm size filled with the  
82 different potting mixtures. Watering of the seedlings was done once daily. Seedling

83 height (cm), stem diameter (mm) and number of branches were assessed after 4 weeks  
 84 after transplanting for 12 weeks.

85 **Table 1:** Laboratory Analysis of Organic and Inorganic Fertilizer

86 <b>Sample Code</b>	%N	%P	%K
87 Poultry manure	3.47	1.18	1.38
88 Cow dung	4.06	0.33	0.77
89 N.P.K. 15.15.15	15.00	15.00	15.00

90 Treatment Combinations

- 91 Where; (1) (i)  $S_1f_0$  - Arable soil without fertilizer (control)  
 92 (ii)  $S_1f_1$  - Arable soil with poultry manure  
 93 (iii)  $S_1f_2$  - Arable soil with cowdung  
 94 (iv)  $S_1f_3$  - Arable soil with N.P.K (15.15.15)  
 95 (2) (i)  $S_2f_0$  - Forest reserve soil without fertilizer (control)  
 96 (ii)  $S_2f_1$  - Forest reserve soil with poultry manure  
 97 (iii)  $S_2f_2$  - Forest reserve soil with cowdung  
 98 (iv)  $S_2f_3$  - Forest reserve soil with N.P.K. (15.15.15)  
 99 (3) (i)  $S_3f_0$  - Natural Forest soil without fertilizer (control)  
 100 (ii)  $S_3f_1$  - Natural Forest soil with poultry manure  
 101 (iii)  $S_3f_2$  - Natural Forest soil with cowdung  
 102 (iv)  $S_3f_3$  - Natural forest soil with N.P.K (15.15.15)

103 **Data Analysis**

104 Analysis of variance was used to analyze the data obtained where significant, using SAS  
 105 statistical package; Least Significant Differences (LSD) was used to separate the means.

106 **RESULTS AND DISCUSSION**

107 **Table2:** Effect of treatment on stem diameter (mm) of *Moringa oleifera* seedlings

Treatment	2WAT	4WAT	6WAT	8WAT	10WAT	12WAT
-----------	------	------	------	------	-------	-------

S <sub>1</sub> F <sub>0</sub>	1.90	2.06	2.00	2.03	2.17	3.19
S <sub>1</sub> F <sub>1</sub>	1.93	2.07	1.97	1.97	2.17	3.20
S <sub>1</sub> F <sub>2</sub>	2.03	2.10	2.00	2.03	2.20	2.84
S <sub>1</sub> F <sub>3</sub>	2.07	2.20	2.40	2.63	3.17	4.28
S <sub>2</sub> F <sub>0</sub>	1.97	2.03	0.63	0.63	0.63	1.45
S <sub>2</sub> F <sub>1</sub>	2.10	2.13	1.97	1.93	2.00	3.17
S <sub>2</sub> F <sub>2</sub>	2.07	1.40	1.40	1.43	1.50	2.42
S <sub>2</sub> F <sub>3</sub>	2.03	2.03	0.00	0.00	0.00	0.00
S <sub>3</sub> F <sub>0</sub>	1.90	2.03	1.40	3.80	1.77	2.85
S <sub>3</sub> F <sub>1</sub>	2.03	2.07	2.13	2.17	2.60	3.00
S <sub>3</sub> F <sub>2</sub>	1.90	2.03	2.17	2.22	2.60	3.00
S <sub>3</sub> F <sub>3</sub>	1.93	2.00	2.07	2.47	3.47	4.53
Mean	1.99	2.01	1.68	1.94	2.02	2.83
Significance	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***
Se±	0.1023	0.1023	0.1023	0.1023	0.1023	0.1023
LSD	0.9105	0.9105	0.9105	0.9105	0.9105	0.9105
C.V%	5.3	5.3	5.3	5.3	5.3	5.3

108 **Table 2:** Effect of treatment on stem diameter (mm) of *Moringa oleifera* seedlings  
109 Where: - WAT= Week after transplanting\*\*\* significantly difference (p<0.001).

110 Table 2 shows that there was no significant difference in stem diameter of the seedlings  
111 subjected to various treatments at P<0.001 while N.P.K. 15.15.15 in natural forest soil  
112 (S<sub>3</sub>F<sub>3</sub>) had the highest stem diameter of 4.53mm. There was a continuous increase in stem  
113 diameter across the weeks. This was followed by N.P.K. 15.15.15 in arable soil (S<sub>1</sub>F<sub>3</sub>)  
114 that had value of 4.28mm against cow dung in forest reserve soil (S<sub>2</sub>F<sub>2</sub>) which had the  
115 lowest stem diameter of 1.5mm across the weeks. This confirms the findings of Waheed  
116 *et al.*, (2001) for *Camelia sinensis* that nitrogen containing fertilizers such as N. P.K had  
117 a significant effect on seedling growth parameters. Almeida (1997) had earlier reported  
118 that N.P.K fertilizer gave a positive response in seedling growth of cashew. This also  
119 supports the findings of Larcheveque *et al.*, (2011) that chemical fertilizers promote

120 higher growth and root development compared to livestock organic manure in a Poplar  
 121 plantation.

122 **Table 3:** Effect of soil mixture on the height (cm) of *Moringa oleifera* seedlings

Treatment	2WAT	4WAT	6WAT	8WAT	10WAT	12WAT
S <sub>1</sub> F <sub>0</sub>	32.40	34.50	36.40	37.40	40.00	43.04
S <sub>1</sub> F <sub>1</sub>	28.40	37.33	41.33	42.33	46.00	54.25
S <sub>1</sub> F <sub>2</sub>	32.00	36.67	39.50	34.17	46.67	55.80
S <sub>1</sub> F <sub>3</sub>	40.17	41.33	46.00	54.33	66.67	74.55
S <sub>2</sub> F <sub>0</sub>	32.33	31.67	11.00	11.67	12.33	15.61
S <sub>2</sub> F <sub>1</sub>	34.73	33.73	35.17	37.83	42.50	48.72
S <sub>2</sub> F <sub>2</sub>	24.83	20.67	28.77	30.00	30.67	35.74
S <sub>2</sub> F <sub>3</sub>	26.33	34.33	0.00	0.00	0.00	0.00
S <sub>3</sub> F <sub>0</sub>	28.50	31.50	32.17	35.00	36.00	39.50
S <sub>3</sub> F <sub>1</sub>	34.83	35.57	46.50	50.33	52.33	60.28
S <sub>3</sub> F <sub>2</sub>	34.50	34.17	42.33	49.33	54.67	64.45
S <sub>3</sub> F <sub>3</sub>	30.90	32.33	43.33	58.00	73.33	89.67
Mean	31.66	33.81	3.38	37.45	41.74	48.47
significance	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***
Se±	3.638	3.638	3.638	3.638	3.638	3.638
LSD	0.9105	0.9105	0.9105	0.9105	0.9105	0.9105
C.V%	5.3	5.3	5.3	5.3	5.3	5.3

123 Where:- WAT = Week after transplanting\*\*\* significantly difference (<0.001).

124 Table 3 revealed that there was no significant difference in the height of Moringa  
 125 seedlings among the treatments at P<0.001. N.P.K. 15.15.15 in natural forest soil (S<sub>3</sub>F<sub>3</sub>)  
 126 had highest height of 89.67cm (P<0.001), this was achieved due to the increase in the  
 127 growth rate (height) of Moringa across the weeks. This was followed by N.P.K. 15.15.15  
 128 in arable soil (S<sub>1</sub>F<sub>3</sub>) that had a value of 74.55cm, the trend was also maintained across the  
 129 weeks while cow dung in forest reserve soil (S<sub>1</sub>F<sub>3</sub>) had the lowest height of 35.74cm.  
 130 Similar positive results has been reported by (Hoque, *et al.*, 2004) seedling growth was  
 131 enhanced significantly with the application of N.P.K fertilizer. Tree seedlings need

132 nutrients to grow, nitrogen for lots of green leaves, phosphorus for new tissues  
 133 particularly the roots and potassium for seedling vigour.

134 **Table 4:** Effect of soil mixture on number of branches of *Moringa oleifera* seedlings

Treatment	2WAT	4WAT	6WAT	8WAT	10WAT	12WAT
S <sub>1</sub> F <sub>0</sub>	6.00	6.00	5.67	6.33	6.67	6.67
S <sub>1</sub> F <sub>1</sub>	7.00	6.00	6.33	6.00	6.00	6.00
S <sub>1</sub> F <sub>2</sub>	5.67	4.67	6.00	7.33	6.67	7.52
S <sub>1</sub> F <sub>3</sub>	4.00	6.00	8.67	9.67	9.67	10.23
S <sub>2</sub> F <sub>0</sub>	5.00	2.00	1.33	1.33	1.33	1.33
S <sub>2</sub> F <sub>1</sub>	6.33	3.67	4.00	5.00	5.33	6.50
S <sub>2</sub> F <sub>2</sub>	5.67	3.00	3.00	3.67	3.33	3.33
S <sub>2</sub> F <sub>3</sub>	5.67	4.33	0.00	0.00	0.00	4.33
S <sub>3</sub> F <sub>0</sub>	5.00	3.67	4.67	4.67	4.00	4.00
S <sub>3</sub> F <sub>1</sub>	6.67	6.33	7.33	6.67	6.67	6.67
S <sub>3</sub> F <sub>2</sub>	6.33	7.00	7.00	8.67	8.00	8.00
S <sub>3</sub> F <sub>3</sub>	6.33	6.33	8.67	10.33	10.00	11.00
<b>Mean</b>	5.81	4.91	5.58	6.17	6.00	6.33
<b>significance</b>	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***	<0.001***
<b>Se±</b>	0.570	0.570	0.570	0.570	0.570	0.570
<b>LSD</b>	2.830	2.830	2.830	2.830	2.830	2.830
<b>C.V%</b>	10.3	10.3	10.3	10.3	10.3	10.3

135 Where: WAT= week after transplanting\*\*\* significantly different (P < 0.001)

136 Table 4 shows that there were significant differences in number of branches of Moringa  
 137 seedlings among the treatments at P<0.001. N.P.K. 15.15.15 in natural forest soil (S<sub>3</sub>F<sub>3</sub>)  
 138 had the highest value of 11.00. This was achieved due to continuous production in  
 139 number of branches across the weeks. This was followed by N.P.K. 15.15.15 in arable  
 140 soil (S<sub>1</sub>F<sub>3</sub>) that had the value of 10.23 which was also maintained across the weeks while  
 141 cow dung in forest reserve soil (S<sub>2</sub>F<sub>2</sub>) had the lowest value of 3.33 across the weeks. This  
 142 supports the findings of Jaenicke (1999) who stated that cow dung contains 0.3%  
 143 Nitrogen, 0.2% phosphoric acid and 0.1- 0.5% while Ajay, (2017) also reported that cow  
 144 dung is not as rich in nitrogen as many other types of fertilizers. He reported that cow  
 145 dung has about 8% nitrogen, 2% phosphorus and 1% potassium. These nutrients are also  
 146 slowly infused into the soil.

147

148

149

## 150 **Conclusion**

151 Based on the findings of these studies, seedlings raised with N.P.K. 15:15:15 mixed with  
152 natural forest soil had the highest plant height (76.30 cm), stem diameter (3.47 mm) and  
153 number of branches (11.00).

## 154 **Recommendation**

155 Natural forest soil mixed with N.P.K. 15:15:15 could be recommended for use in raising  
156 seedlings of *Moringa oleifera* for optimum growth performance since the seedlings  
157 presented most noticeable growth at the nursery stage.

158

159

## 160 **References- can include more recent year references**

161 Adegbidi, H.G.; Briggs, R.D.; White, E.H.; Abrahamson, L.P.; Volk, T.A. Effects of  
162 organic amendments and slow release N<sub>2</sub> fertilizer on willow stem biomass  
163 production and soil chemical characteristics. **Biomass Bioenergy** v.25,  
164 no.4, p. 389-398, 2003.

165 Ajay, M.R. Effective use of cow dung manure for healthy plant growth. **International**  
166 **Journal of Advanced Research and Development** v, 2 5, no. p 218-221,  
167 2017.

168 Almeida, A.G.; Almeida, F.C.G.; Meneses, J. J. Effect of N, P and K fertilization on  
169 productivity in dwarf early air-layered Cashew under irrigation. **Turri alba**  
170 v. 44 no.3, p. 168-178,1997.

171 FRIN-Forestry Research Institute of Nigeria. Annual **Metereological** Report. 2013.

172 Hoque, A. T.; M. R.; Hossian, M.K.; Mohiuddin, M.; Hoque, M. M. (2004). Effect of  
173 inorganic fertilizers on initial growth performance of *Michelia champaca*  
174 Linn. Seedlings Nursery. **J.Biol.Sci** v 4 p. 489-497.

175 Jaenicke, H. Good tree nursery practices: practical guidelines for research nurseries.  
176 Nairobi ICRAF, 1999

177 Jose, L.M (2003). Nitrogen and phosphorus resorption in tree of neotropical rainforest.  
178 **J.Trop Ecol** v, 19 p.465-468.

- 179 Larcheveque, M.; Desroches, A.; Larocque, G.R. Comparison of manure compost and  
180 mineral fertilizer for hybrid poplar plantation establishment on boreal clay  
181 soils . **Ann. For, Sci** v. 68, no.4, p 849-860. 2011.
- 182 Leone, A.; Spada, A.; Battezzati, A.; Schiraldi, A.; Aristil, J.; Bertoli, S. "Cultivation,  
183 Genetic, Ethnopharmacology, Phytochemistry and Pharmacology of  
184 *Moringa oleifera* Leaves: An Overview" .**Int J Mol Sci**.v, 16 no.6: 12.  
185 p791~835. 2015.
- 186 Olson, M.E. Flora of North America Committee, ed.Flora summary :Moringaceae  
187 Drumstick Family. **Flora of North America North of Mexico**.7.New York  
188 and Oxford.p 167-169. 2010
- 189 Torondel, B.; Opare, D.; Brandberg, B.; Cobb, E.; Cairncross, S. (2014). "Efficacy of  
190 *Moringa oleifera* leaf powder as a hand- washing product: A crossover  
191 controlled study among healthy volunteers". **BMC Complementary and  
192 Alternative Medicine**.v,14 no, 57. doi : 10.1186/1472-6882-14-57 .2014.
- 193 Waheed, A.; Jamal, A.; Hamad, F.S; Ahmad, N. Growth of tea seedlings affected by  
194 different levels and application of N.P.K Urea. **Online Journal of  
195 Biological Sciences** v, 1 no.3.p101-102.2001.
- 196
- 197