

Diagnostic significance of Beclin-1 and Transforming growth factor β in Breast Cancer

5
7

ABSTRACT

Background: The main cause of cancer deaths amongst women breast cancer remains a clinical and social challenge, and a serious public health problem. On a worldwide level, it continues to be a devastating disorder. BECN1 is a tumor suppressor gene implicated in the initiation of autophagy. It encodes beclin-1 protein that inhibits cancer growth. There is wide disputation concerning its role in initiation, promotion of tumor and predictive importance of autophagic molecules. Transforming growth factor β (TGF- β) induces process of epithelial-mesenchymal transition (EMT) keeping, epithelial cells more motile and invasive resulting in cancer progression and metastasis.

Aim: Detection of beclin-1 expression level in metastatic and non-metastatic breast cancer patients and study its role in tumorigenesis of breast cancer through attainable association with the inflammatory cytokine, TGF- β .

Methods: Expression levels of beclin-1 and TGF- β were assessed in 70 breast cancer female patients and 20 controls using quantitative real-time PCR.

Results: Beclin-1 expression levels as well as TGF- β were significantly higher in metastatic breast cancer patients and non-metastatic patients compared to controls. Positive correlation was found between beclin-1 expression level and TGF- β expression level in breast cancer patients.

Conclusion: Our results indicated that over-expression of both beclin-1 and TGF- β was associated with aggressive clinical outcomes of breast cancer patients and tumor growth. These findings suggest that beclin-1 and TGF- β are associated with tumorigenesis of breast cancer.

Keywords: autophagy, BECN1, TGF- β , breast cancer.

1. INTRODUCTION

Breast carcinoma is the y main cancer that influences females accounting for 37.7% of their overall malignancies in Egypt.

Moreover, it is important reason of cancer related mortality in Egypt, constituting about 29.1% of all most cancer associated mortality [1]. Autophagy is a cellular degradation or "self-eating" pathway extremely preserved throughout all

existence kingdoms. [2]. Considering the truth that autophagy is implicated in numerous cellular processes, and keeping in mind the complexness complexity of the molecular mechanisms of tumor initiation and development, it is no longer

shocking that the interruption of autophagy plays role in tumor inatition initiation and progression. Indeed, cancer was the primary disease allied with disturbed autophagy besides, the first for which medical trials in humans were performed [3].

Beclin-1 (BECN1) is a rate-limiting element of autophagy and a haplosufficient cancer suppressor that is fundamental for embryonic development. Moreover, this protein is a core factor essential for autophagosome formation [4]. Transforming

growth factor-beta (TGF- β) is a protein that is essential regulator in homeostasis in various organs [5]. (As a signaling molecule, TGF- β has a variety of biological function and participates in several physiological and pathological processes

[6].

2. MATERIAL AND METHODS

2.1 Ethical approval

The study was approved by the Institutional Review Board (IRB) of the NCI, Cairo University and was conducted according to the rules of Helsinki declaration for human studies. A Written informed consent was obtained from all study subjects.

2.2. Subjects

The study included 70 Egyptian women with breast cancer at distinctive stages, their age ranged from (23-76 years). They were collected from Kasr Alainy hospital. Patients were diagnosed by clinical examination and affirmed by mammography and surgical biopsies. Twenty clinically normal adjoining specimens were moreover collected from adjacent healthy breast tissue of non-metastatic breast cancer cases. All specimens were subjected to estimation of expression levels of beclin-1 and TGF- β in breast tissue.

The studied subjects were separated into three groups as following follows:

Group I: (n=26) metastatic breast cancer female patients with distant metastasis (bone, liver, and lung).

Group II: (n=44) non-metastatic breast cancer female patients.

Group III: (n=20) healthy adjacent normal tissues of non-metastatic breast cancer cases as a control group.

2.3. Tissue sampling and RT PCR analysis

Total RNA was extracted from breast tissues using Qiagen tissue extraction kit (Qiagen, USA) according to instructions of manufacture. The first strand cDNA was derived from total RNA (0.5 μ g) and oligo (dT) using the Reverse Transcription System (The QuantiTect Reverse Transcription Kit). For real-time quantitative RT-PCR, gene specific primers listed in (Table 1) and QuantiTect SYBR Green PCR Kit were used. GAPDH was used to normalize the quantity of specific mRNA. The amplification efficiency determined for both target and housekeeping genes was equal. Relative expression levels were calculated by the $2^{-\Delta\Delta C_t}$ method.

Table 1: List of primers sequences applied in real time-PCR.

Genes	Forward primer	Reverse primer
GAPDH	CAATGACCCCTTCATTGACC	TTGATTGGAGGGATCTCG
Beclin-1	GGCTGAGAGACTGGATCGG	CTGCGTCTGGGCATAACG
TGF- β	ACATTGACTTCCGCAAGCAC	GTCCAGGCTCCAAATGTAGG

2.4. Statistical analysis:

Statistical analysis of data was performed by Sigma Plot version 12.5 and Graphpad Prism 5. Data were illustrated as mean \pm standard deviation. Differences among groups were analyzed by Shapiro-Wilk test. Post-hoc testing was performed by the Tukey test to compare the difference among the groups. Simple linear correlation (Pearson correlation coefficient test) (r) was also done to test for linear relations between beclin-1 and TGF- β with other variables. P-value is

considered significant if < 0.05 . Receiver operating characteristics curves (ROC curves) were utilized to assess the diagnostic performance of the studied parameters.

3. RESULTS

3.1 Expression levels of beclin-1 and TGF- β in different clinicopathological factors in patients with breast cancer

The associations between the levels of beclin-1 and TGF- β with clinicopathological factors were analyzed. The levels of beclin-1 and TGF- β were increased with TNM stage ($p < 0.01$) but showed no significant correlation with the other factors (Table 2).

Table (2): Correlation between expression levels of beclin-1 and TGF- β with clinicopathological factors in the malignant groups

Clinicopathological Factors	Malignant patients (70)	Beclin-1 P value	TGF- β P value
Metastasis Metastatic Non-metastatic	26 (37.1 %) 44 (62.9%)	>0.05	>0.05
Family history No family history Family history	54 (77.1%) 16 (22.9%)	>0.05	>0.05
Menopause Pre-menopause Post-menopause	47 (67.1%) 23 (32.9%)	0.05>	0.05>
Site Right Left Bilateral	35 (50%) 32 (45.7%) 3 (4.3%)	>0.05	>0.05
Pathology IDC II Medullary IDC III IDC + ILC ILC	58 (82.8%) 1 (1.4%) 6 (8.6%) 3 (4.3%) 2 (2.9 %)	>0.05	>0.05
ER sensitivity Positive Negative	49 (70%) 21 (30%)	>0.05	>0.05
PR sensitivity Positive Negative	42 (60%) 28 (40%)	>0.05	>0.05
HER sensitivity Positive Negative	29 (41.4%) 41 (58.6%)	>0.05	>0.05
Molecular diagnosis HER 2 enriched	29 (41.4%)	>0.05	>0.05

Triple negative	11 (15.7 %)		
Luminal B	26 (37.15%)		
Luminal A	4 (5.75%)		
Stage			
Stage II	18 (25.7%)	<0.05**	<0.05**
Stage III	26 (37.15%)		
Stage IV	26 (37.15%)		

a* significant from stage II

3.2 Expression levels of beclin-1 and TGF- β in patients with breast cancer compared to control group

Expression Level of beclin-1 was highly significant in metastatic malignant group (11 ± 3.6) and non-metastatic malignant group (9.3 ± 4.5) compared to the control group (1.2 ± 0.34) with (p value < 0.05), as well as expression Level of TGF- β was highly significant in metastatic malignant group (9.5 ± 7.9) and non-metastatic malignant group (9.3 ± 7.7) compared to the control group (1.1 ± 0.30) with (p value < 0.05) (Figure 1).

a* significant from controls




Figure (1): Mean \pm SD of beclin-1 and TGF- β expression levels for the studied groups.

3.3 Receiver operating characteristics (ROC) curves

Receiver operating characteristic curves were carried out to assess the diagnostic performance of beclin-1 and TGF- β and their sensitivity and specificity independently or in combination. The best cut-off value for beclin-1 in malignant patients was > 1.9 ($P < 0.0001$) with 95 % sensitivity and 98.5% specificity producing area under the curve (AUC) 0.997 (Figure 2). For tissue expression of TGF- β in malignant patients; the best cut-off point was > 3.98 ($P < 0.001$) with 100% sensitivity and 98.5% specificity producing AUC=1 (Figure 3). On the other hand, the best cut-off point for both beclin-1 and TGF- β in combination was > 1.9 ($P < 0.001$) with 97.5 % sensitivity and 99.2 % specificity and AUC= 0.99 (Figure 4)

ROC curve of beclin 1




Figure (2): ROC curve of beclin-1.

ROC curve of TGF- β




Figure (3): ROC curve of TGF- β .

ROC curve of beclin 1 & TGF- β





Figure (4): ROC curve of beclin1 and TGF- β .

3.4 Pearson's correlation analysis

116 There was a significant positive association between expression level of beclin-1 in breast cancer patients and TGF- β (r=
117 0.241 and $P < 0.05$) (Figure 5).
118
119
120
121



122 **Figure (5): Correlation between beclin-1 and TGF- β expression levels.**
123
124
125

126 4. DISSCUSSION

127 Autophagy is utilized by ordinary and cancer cells. However, the mechanism of autophagy in different cancers including
128 breast cancer remains not clear in oncogenic/protumorigenic and tumor-suppressor feature [7]. The purpose of our study
129 was to assess expression levels of beclin-1 and TGF- β in breast carcinoma Egyptian women compared with to normal
130 controls and to determine the potential value of beclin-1 and TGF- β as molecular biomarkers for diagnosis of breast
131 cancer in Egyptian women. We found, compared with to the expression in normal breast tissue, that beclin-1 and TGF- β
132 were over-expressed in tumor tissue. Moreover, the over-expression of beclin-1 and TGF- β in breast cancer was
133 considerably associated with TNM stage. These findings may suggest that beclin-1 and TGF- β have direct influence on
134 development and progression of breast carcinoma. In the current study, beclin-1 level was significantly increased in breast
135 cancer groups in comparison to healthy subjects (P value < 0.01). Our results are supported with different previous
136 studies presented by **Hamurcu, et al.**, [8] who have studied the relative concentrations of beclin-1 in the tissue of
137 patients with breast cancer, metastatic disease and healthy women and stated that expression levels of beclin-1 was
138 higher in highly aggressive, metastatic and noninvasive cancer cells compared to non-tumorigenic normal human breast
139 cells. The function of autophagy is exist different opinion in solid cancers including breast cancer [8]. In another study
140 **Wang, et al.**, [9] founded a significant increase of beclin-1 expression in eighteen breast carcinoma patients which was
141 associated with tumor progression. Also, **Park, et al.**, [10] revealed that the expression of beclin-1 was increased in
142 tumor cells relative to normal-appearing and adjacent colonic mucosa in all cases. Similarly, **Chen, et al.**, [11] reported an
143 increased expression of beclin-1 in tumor tissue compared with under-expression in normal gallbladder specimen ($p <$
144 0.05). The concept that autophagy represents a mechanism that promotes tumor growth is based on the need of tumor
145 cells to adjust to ischemia in the surroundings that are hypoxic, besides growth factors and supplemental deprivation.
146 Consistent with this aspect, autophagy is activated in hypoxic environment of tumors [12]. Beclin-1 was firstly found to
147 have tumor suppressor role. However, elevated expression of beclin-1 was related with cancer progression in some
148

149 carcinoma [13]. In contrast to our study, decreased beclin-1 expression was observed in seventy percent of the breast
150 tumors, and the protein levels were co-related to the mRNA levels [14]. However, **Claude-Taupin, et al.**, [15] reported
151 that no major difference in beclin-1 mRNA expression between tumor and healthy adjacent tissues. The current study
152 revealed that TGF- β was significantly increased in breast carcinoma patients compared to control group (P value< 0.001)
153 and this finding **are is** affirmed by **Ciftci, et al.**, [16] who have studied the expression level of TGF- β and established that
154 the mean serum TGF- β level of breast cancer patients was considerably higher than controls. There was no considerable
155 difference according to known disease-related clinicopathological parameters. Our finding is supported by **El-Aziz et al.**,
156 [17] who revealed that the level of TGF- β was considerably higher in malignant groups than normal control group with p
157 value (<0.0001). The results of the present study was supported by **Scherer et al.**, [18] who detected **the level of TGF-**
158 **β** in breast cancer patients; **he found** that its level was increased. In the current study, ROC curve of beclin-1 showed
159 95% sensitivity and 98.5% specificity and this is supported by **Harb, et al.**, [19] who reported that the sensitivity of beclin-1
160 as a predictor for advanced stage of IDC was 85.5% and the specificity was 98.5%. Also, ROC curve of TGF- β showed
161 100% sensitivity and 98.5% specificity and this is supported by **El Hussein, et al.**, [20] who reported that 98.2%
162 sensitivity and 100% specificity demonstrating the diagnostic power of this studied marker in differentiating between
163 breast cancer patients and controls. TGF- β is connected with expanded cancer progression, higher cell movement,
164 cancer invasiveness, and metastasis. Furthermore, it is included in cancer surrounding medium alteration and
165 advancement of migration and invasiveness [21]. TGF- β elevated the mRNA levels of beclin-1, and other protein kinase
166 implicated in death process. Moreover, TGF- β evoked autophagy in some mammary carcinoma cell. These findings
167 illustrate that TGF- β signaling pathway activates autophagy in certain human cancer cells and that induction of autophagy
168 is a novel aspect of biological role of TGF- β [22]. Concerning clinicopathological factors, the over-expression of beclin-1
169 was considerably related with TNM stage (p< 0.05) but it showed no noteworthy association with the other factors. The
170 current results were similar to **Chen, et al.**, [12] **work** who declared that over expression of beclin-1 was essentially
171 related with TNM stage but had no vital relationship with age, sex, lymphatic metastasis, or tumor differentiation. Also, the
172 affiliation between hormonal status and TGF- β expression was examined for both Estrogen receptors (ER) and
173 Progesterone receptors (PR) by **El-Aziz, et al.**, [17] concluding that no noteworthy difference of TGF- β expression
174 neither between ER positive versus ER negative tumors, nor between PR positive and PR negative tumors and
175 this is similar to our data. Concerning the relation between tissue tumor subtypes and TGF- β level, **El-Aziz et al.**, [17]
176 reported no critical difference between TGF- β level and tissue tumor subtypes as we found.

177 178 179 **5. CONCLUSION**

180 This study affirmed **overexpression** of beclin-1 in breast carcinoma tissues and demonstrated that beclin-1 and TGF- β
181 **high expression** levels were associated with forceful clinical outcomes of breast cancer patients. **Also, their**
182 **overexpression** contributed to the tumor development. **The present study has a limitation: the sample size was relatively**
183 **small. Consequently, the present study should be done on large number of breast cancer population to emphasize our**
184 **results and clarify the diagnostic importance of beclin1 and TGF- β in the process of tumorigenesis.**

186 187 188 **REFER/ENCES**

- 189
190
191 1. Morsy M M, Raafat A, Mohamed M, Fayed A. (2018). Prognostic Value of Percentage of Positive to Total Excised

Axillary Lymph Nodes in Egypt with Triple Negative Breast Cancer : Multiple-Centers Experience. *Int. J. of Life Sciences*, 6(3), 719–732.

2. Arakawa S, Honda H, Yamaguchi H, Shimizu S. (2017). Atg12 Conventional autophagy ATG5-ATG12 LC3-PE Beclin-1 Alternative autophagy Rab-9. *Proceedings of the Japan Academy, Series B*, 93(6), 378–385.
3. Abraham NM, Kirubel MM, and Abraham DA. (2018). Autophagy as a Possible Target for Cancer Therapy. *Journal of Orthopedic Oncology*, 04(01), 1–9. <https://doi.org/10.4172/2472-016X.1000124>.
4. Morris DH, Yip CK, Shi Y, Chait BT, Wang QJ. (2015). BECLIN 1-VPS34 COMPLEX ARCHITECTURE: UNDERSTANDING THE NUTS AND BOLTS OF THERAPEUTIC TARGETS. *Front. Biol. (Beijing)*, 10(5), 398–426. <https://doi.org/10.1038/s41556-018-0061-z>.
5. Furler RL, Nixon DF, Brantner CA, Popratiloff A, Uittenbogaart CH. (2018a). TGF- β sustains tumor progression through biochemical and mechanical signal transduction. *Cancers*, 10(6), 1–18. <https://doi.org/10.3390/cancers10060199>.
6. Wang ., Gong X, Xu J, Xie R. (2018). The Role of TGF- β in Gastrointestinal Cancers. *Journal of Cancer Science & Therapy*, 10(11), 345–350. <https://doi.org/10.4172/1948-5956.1000566>.
7. Mizushima N. (2017). The exponential growth of autophagy-related research: from the humble yeast to the Nobel Prize. *FEBS Letters*, 591(5), 681–689. <https://doi.org/10.1002/1873-3468.12594>.
8. Hamurcu Z, Delibaşı N, Geçene S, Şener EF, Dönmez-Altuntaş H, Özkul Y, et al, (2018). Targeting LC3 and Beclin-1 autophagy genes suppresses proliferation, survival, migration and invasion by inhibition of Cyclin-D1 and uPAR/Integrin β 1/ Src signaling in triple negative breast cancer cells. *Journal of Cancer Research and Clinical Oncology*, 144(3), 415–430. <https://doi.org/10.1007/s00432-017-2557-5>.
9. Wang MC, Wu AG, Huang Y, Shao, GL, Ji SF, Wang RW, et al, (2015). Autophagic regulation of cell growth by altered expression of Beclin 1 in triple-negative breast cancer. *International Journal of Clinical and Experimental Medicine*, 8(5), 7049–7058.
10. Park JM, Huang S, Wu TT, Foster NR, Sinicrope FA (2013). Prognostic impact of Beclin 1, p62/sequestosome 1 and LC3 protein expression in colon carcinomas from patients receiving 5-fluorouracil as adjuvant chemotherapy. *Cancer Biology and Therapy*, 14(2), 100–107. <https://doi.org/10.4161/cbt.22954>.
11. Chen Y, Yan J, Yu S, Wang X, Zheng Q. (2014). Over-expression of beclin-1 in gallbladder carcinoma and its relationship with prognosis. *Wspolczesna Onkologia*, 18(3), 171–176. <https://doi.org/10.5114/wo.2014.41395>.
12. Ávalos Y, Canales J, Bravo-Sagua R, Criollo A, Lavandero S, Quest AFG, et al, (2014). Tumor Suppression and Promotion by Autophagy. *BioMed Research International*, 2014, 1–15. <https://doi.org/10.1155/2014/603980>.
13. Parkhitko A, Myachina F, Morrison TA, Hindi KM, Auricchio N, Karbowiczek M, Henske EP. (2011). Tumorigenesis in tuberous sclerosis complex is autophagy and p62/sequestosome 1 (SQSTM1)-dependent. *Proceedings of the National Academy of Sciences*, 108(30), 12455–12460. <https://doi.org/10.1073/pnas.1104361108>.
14. Li Z, Chen B, Wu Y, Jin F, Xia Y, Liu, X. (2010). Genetic and epigenetic silencing of the beclin 1 gene in sporadic breast tumors. *BMC Cancer*, 10(98), 1–12. <https://doi.org/10.1186/1471-2407-10-98>.
15. Claude-Taupin A, Fonderflick L, Gauthier T, Mansi L, Pallandre JR, Borg et al, (2018). ATG9A Is Overexpressed in Triple Negative Breast Cancer and Its In Vitro Extinction Leads to the Inhibition of Pro-Cancer Phenotypes. *Cells*, 7(12), 1–17. <https://doi.org/10.3390/cells7120248>.
16. Ciftci R, Tas F, Yasasever CT, Aksit E, Karabulut S, Sen F, et al, (2014). Clinical significance of serum transforming growth factor beta 1 (TGFB1) level in breast cancer. *Journal of Clinical Oncology*, 32(15), e11526. <https://doi.org/10.1200/jco.2014.32.15>.
17. El-Aziz G, Kamel MM, Alkaffas M, Abdelhady EG, Rashed LA. (2018). Can Transforming Growth Factor Beta Affect Breast Cancer by Targeting MicroRNA 195? *J Mol Cell Biochem*, 2(1), 1–4. Retrieved from <http://www.imedpub.com/journal-molecular-cellular-biochemistry>.
18. Scherer SD, Bauer J, Schmaus A, Neumaier C, Herskind C, Veldwijk MR, Sleeman JP. (2016). TGF- β 1 is present at high levels in wound fluid from breast cancer patients immediately post-surgery, and is not increased by intraoperative radiation therapy (IORT). *PLoS ONE*, 11(9), 1–14. <https://doi.org/10.1371/journal.pone.0162221>.
19. Harb OA, Salem AA., Haggag R, El-shorbagy S, Gertallah LM. (2016). Immunohistochemical expressions of SQSTM1 / p62 , Beclin-1 , and SOX4 in infiltrating duct carcinoma of the breast. *Egyptian Journal of Pathology*, 62, 95–103. <https://doi.org/10.1097/01.XEJ.0000484380.44411.44>.
20. EL-Husseini M, Hussein F, Bassily N, Abdelghany B. (2013). Clinical significance of TGF alpha , TGF beta1 and VEGF in Sera of Egyptian patients with breast cancer Clinical Significance of TGF Alpha , TGF Beta1 and VEGF in Sera of Egyptian Patients with Breast Cancer Abstract : The Egyptian Journal of Hospital Medicine July, 52, 555–565. <https://doi.org/10.12816/0000592>.
21. Zarzynska J. (2014). Two faces of TGF-beta1 in breast cancer. *Mediators of Inflammation*, 2014, 141747. <https://doi.org/10.1155/2014/14174>.
22. Kiyono K, Suzuki H, Matsuyama H, Morishita Y, Komuro A, Kano MR, et al, (2009). Autophagy is activated by TGF-beta and potentiates TGF-beta-mediated growth inhibition in human hepatocellular carcinoma cells. *Cancer Research*, 69(23), 8844–8852.

UNDER PEER REVIEW