

GROWTH AND YIELD OF RADISH (*Raphanus sativus* L.) AS INFLUENCED BY DIFFERENT LEVELS OF KALLI ORGANIC FERTILIZER ON THE JOS PLATEAU

Umar, U. M designed the study and wrote the protocol. Ibrahim, Iro, I wrote the first draft of the manuscript, performed the statistical analysis and managed the analyses of the study. Obidola, SM, managed the literature searches. All authors read and approved the final manuscript.

ABSTRACT

Aims: To determine the effect of Kalli organic fertilizer on the growth and yield of radish.

Study Design: The experimental design used for this study was a Randomized Complete Block Design (RCBD) consisting of four (4) treatments (0, 400, 500 and 600kg ha^{-1}) which were replicated four (4) times.

Place and Duration of the Study: The experiment was conducted at Federal College of Forestry Jos, Plateau State located in the North Central part of Nigeria between September – October, 2018.

Methods: Soil samples were collected and analysed. Agronomic practices such as land preparation, planting, fertilizer application, weeding and harvesting were also carried out. The data was collected on plant height, number of leaves, leaf area, number of roots, length of roots, diameter of roots, root weight and root yield. Analysis of Variance (ANOVA) using Minitab 17 statistical package was used to analyse the data and where significance was declared Duncan Multiple Range Test (DMRT) was used to separate the means.

Results: The findings from this research work reveal that there was no significant effect of Kalli organic fertilizer on the plant height, number of leaves. But the leaf area was found to be significant ($p=0.05$) at 8WAP with the application of 600kg ha^{-1} recording the largest (143.30cm 2) leaf. The number of roots, length of roots, weight of root and total yield was found to be significant ($P=0.05$) at the application of different levels of Kalli organic fertilizer. The application of 600kg ha^{-1} gave the highest (35.50) number of radish roots, highest (24.83cm) length of radish roots, highest (7.20kg) weight of radish and highest (18000kg ha^{-1}) total yield of Radish.

Conclusion: Based on this research study it can be concluded that the application of Kalli (600kg ha^{-1}) organic fertilizer significantly increases the yield of Radish. Thus it is recommended that 600kg ha^{-1} of organic fertilizer can be applied for optimum production of Radish.

Keyword: Radish, Kalli, Organic Fertilizer, Sweet potato

1.0 INTRODUCTION

Radish (*Raphanus sativus* L.) belongs to the family Brassicaceae, genus raphanus and species sativus. It is one the most important and popular root vegetable, grown in tropical, sub-tropical and temperate regions of the world. It is grown both as an annual and a biennial vegetable crop depending upon the type for the purpose it is grown. Radish is predominantly a cool season vegetable crop. But, Asiatic types can tolerate higher temperature than European varieties. In the mild climate, radish can be grown almost all the year round except for few months of summer (PCARRD. 2009).

Radish is grown and consumed all over the world and is considered part of the human diet, even though it is not common among some populations. Its young tuberous roots can be eaten raw in salad or cooked as

a vegetable. It has a pungent flavour and considered as an appetizer. The young leaves are also cooked and eaten as vegetables. The preparations of radish are useful in liver and gall bladder troubles, roots, leaves, flowers and pods are active against gram-positive bacteria, urinary complaints, piles and gastrodynia. Also, salt extracted from roots, dried and burnt to white ash is said to be used in stomach troubles (Satish, 2016).

Most of the nutrients absorbed by plants come from organic matter. Therefore, the unique formulation of Kalli 100% organic fertilizer, which consists of animal and chicken wastes and earth based mineral resources like rock phosphate, lime, gypsum bentonite and more makes it very rich in nutrients content. They provide rich sources of organic matter to be added to the soil for the growth of variety of crops. Thus, Kalli fertilizer aims to serve society by increasing income to farmers through organic fertilization which revitalizes the soil and increase farm yields (Kalli, 2017).

Global awareness for the hazard of long-term use of chemical fertilizer is growing because of this, more and more farmers are shifting to organic fertilizer. Among the benefits of using fertilizers are non toxic food, lower cost, better fertility and of course a safer environment (Xicberg and Offcrmanna, 2002), low soil fertility among others is one of the major challenges in crop production. Hence, the cost of these fertilizer (Kummeling *et al.*, 2008) has necessitated the need to search for less expensive and environmentally friendly fertilizer. This research study therefore was aimed at determining the effect of Kalli organic fertilizer on the growth and yield of radish.

2.0 MATERIAL AND METHOD

2.1 Study Area

The experiment was conducted at Federal College of Forestry Jos, Plateau State located in the North Central part of Nigeria which lies at a Latitude of 9.94 and Longitude 8.89 within the Guinea Savannah region with a mean annual rainfall of 1460mm and temperature range between 19°C and 32°C (Iro *et al.*, 2019).

2.2 Materials

The materials used for this experimental study include certified radish seeds, Kalli organic Fertilizer, Measuring tape, Meter rule, rope, Cutlass, Auger bit, Shovel, Rake, Wheelbarrow, watering can, and Hoe.

2.3 Soil Analysis

Soil Samples were collected at random from the field at two different depths (0-15cm and 15-30cm) with the aid of auger bit, hand trowel and polythene bag to store the sample which was later dried under room temperature and taken to Agricultural Service and Training Centre (ASTC) for analysis.

2.4 Experimental Design

The experimental design used for this study was a Randomized Complete Block Design (RCBD) consisting of four (4) treatments (0, 400, 500 and 600kg ha^{-1}) which were replicated four (4) times.

2.5 Agronomic Practices

a. Land Preparation: Land Preparation was done on the 10th of October, 2018. The land was cleared manually using a cutlass and hoe. The soil was thoroughly dogged and the clumps were broken into finely tilt soil particles for ease of sowing (Iro *et al.*, 2019).

b. Planting: Radish seeds were planted at 2 - 5cm depth and at a spacing of 20cm x 30cm. Kalli Organic fertilizer was applied immediately after planting using the side placement method of 5cm away from the planting spot.

c. Weeding: Weeding was carried out twice manually at two (2) weeks interval.

d. Harvesting: This was done when the radish root protrudes onto the soil surface.

2.6 Data Collection

Radish plants were randomly selected and tagged. Data from the tagged plants was collected at 2 weeks interval on;

- Plant height: The plant height was measured from the base to tip of the plant with the aid of meter rule in centimeter (cm).
- Leaf count: Number of leaves was counted per plant.
- Leaf Area: The leaf length and breadth were measured and leaf area calculated using $\text{Leaf Area (LA)} = \text{Length} \times \text{Breadth}$ in centimeter squared (cm^2)
- Number of Roots: This was done by counting the number of roots each plant produces
- Length of Root: This was done using a meter rule in centimeter (cm)
- Diameter of Root: This was measured using a vernier calipa in centimeter (cm)
- Root Weight: This was done by using a weighting balance to determine the weight of the radish root in kilogram (kg)

Data collected was subjected to analysis of variance (ANOVA) at 5% level of significance using Minitab 23 and where significance was declared, Tukey's method was used to separate the means.

3.0 RESULTS AND DISCUSSION

3.1 Results

3.1.1 Physical And Chemical analysis of Soil in the Study Area

The physical and chemical property of the soil in the study area is presented in Table 1. The result showed that the soil pH was 5.9 which is slightly acidic. It is the preferred soil pH range for good growth and development of most crops. Organic matter had an average value of 11.5%, while the respective nutrient constituents of nitrogen, phosphorus, potassium, calcium and magnesium were 0.036%, 6.2, 95.9, 5.3 and 3.6ppm were in average quantities for optimum production of most crops. The soil can be classified as sandy loam. The percentage composition of sand, silt and clay (10.88% clay, 12% silt, and 77.12% sand) confirms that the presences of organic matter, which make the soil good for crop production.

Table 1: Physical and Chemical Properties of Soil in the Study Area

Sample	PH	N (%)	P PPM	K PPM	Ca PM	MgPPM	O.M(%)	H+ mMol/ 100g	Clay (%)	Silt (%)	Sand (%)	Textural Class
--------	----	-------	-------	-------	-------	-------	--------	---------------	----------	----------	----------	----------------

0-30cm	5.9	0.036	6.2	95.9	5.3	3.6	11.5	1.57	10.88	12	77.12	Sandy loam
--------	-----	-------	-----	------	-----	-----	------	------	-------	----	-------	------------

Source: - Agricultural Services and Training Center KASSA/VOM, 2018.

3.1.2 Composition of Kalli Organic Fertilizer

The composition of Kalli organic fertilizer is presented in Table 2. The result reveals that Kalli organic fertilizer composed of 7.55g/kg of N, 2.86g/kg of P, 1.86g/kg of K, 0.95g/kg of Ca, 0.356g/kg of MgO, 0.795g/kg of Fe, organic Carbon had 17.80g/kg while 35.54g/kg was made of organic matter. This implies that Kalli organic fertilizer has a minimum of 4% N and about 45% organic matter.

Table 2: Composition of Kalli Organic Fertilizer

Variable	Composition (gkg ⁻¹)
N	7.55
P	2.86
K	1.80
Ca	0.95
MgO	0.356
Fe	0.795
Organic Carbon	17.80
Organic Matter	35.54

3.1.3 Plant Height

The result from table 3 shows that there were no significant differences at $P \leq 0.05$ level of probability between the treatments statistically. The application of 600kg ha⁻¹ of kalli fertilizer gave the highest plant height of 24.65cm, followed by 500kg ha⁻¹ with 23.55cm, then 400kg ha⁻¹ with 22.98cm and the control which gave the least plant height 21.48cm. This shows that the application of 600kg of kalli gave the highest plant height at 4WAP. At 8WAP no significant difference was observed at $P \leq 0.05$ between the treatments. The application of 600kg ha⁻¹ of kalli gave the highest plant height with 25.93cm followed by 400kg ha⁻¹ with 25.55cm, then 500kg ha⁻¹ with 25.25cm and the control which has the least plant height with 24.40cm. This shows that the application of 600kg ha⁻¹ of kalli fertilizer gave the highest plant height at 8WAP.

3.1.4 Number of Leaves

The result (table 3) shows that there was no significant difference statistically at $P \leq 0.05$ between the treatments. The application of 600kg ha⁻¹ of kalli fertilizer gave the highest mean number of leaves of 13.90 followed by 500kg ha⁻¹ with 13.05, and then 400kg ha⁻¹ with 12.85 and the control which gave the least mean number of leaves with 12.30. This shows that the application of 600kg ha⁻¹ of kalli has the highest mean number of leaves at 4WAP. At 8WAP no significant difference was observed at $P \leq 0.05$ between the treatments. The application of 600kg ha⁻¹ of kalli fertilizer has the highest mean number of leaves of 15.85, followed by 400kg ha⁻¹ with 15.45, then 500kg ha⁻¹ with 14.70 and the control which gave the least mean number of leaves with 14.20. This shows that the application of 600kg ha⁻¹ of kalli has the highest mean number of leaves.

3.1.5 Leaf Area

No significant ($P \leq 0.05$) effect of Kalli organic fertilizer was observed statistically at 4WAP on leaf area of Radish. However, at 8WAP the result (Table 3) reveals that significant ($P \leq 0.05$) effect of Kalli organic fertilizer was measured on leaf area. The application of 600kg ha^{-1} recorded the largest (143.30cm^2) leaf as against the application of 500 (126.07cm^2), 400 (117.75cm^2) and 0 (87.94cm^2) kg ha^{-1} Kalli organic fertilizer.

Table 3: Growth Characteristics of Radish As Influenced By Different Levels of Kalli Organic Fertilizer on the Jos Plateau

Treatment (kg ha^{-1})a	Plant Height (cm)		Number of Leaves		Leaf Area (cm^2)	
	4WAP	8WAP	4WAP	8WAP	4WAP	8WAP
0	21.48 ^a	24.40 ^a	12.30 ^a	14.20 ^a	94.71 ^a	87.94 ^a
400	22.98 ^a	25.55 ^a	12.85 ^a	15.45 ^a	112.11 ^a	117.75 ^{ab}
500	23.55 ^a	25.25 ^a	13.05 ^a	14.70 ^a	116.83 ^a	126.07 ^{ab}
600	24.65 ^a	25.93 ^a	13.90 ^a	15.85 ^a	129.22 ^a	143.30 ^b
SE $\pm$	1.23	0.70	0.71	0.56	17.11	13.56
CV(%)	10.75	5.48	10.74	7.94	29.30	26.85
LS	NS	NS	NS	NS	NS	*

Means that do not share a letter are significantly different, SE = Standard Error, LS = Level of Significance, ns = Not Significant, * = Significant at $p \leq 0.05$ Level of Significance

3.1.6 Number of Root

The number of roots was found (table 4) to be significant ($P \leq 0.05$) at the application of different levels of Kalli organic fertilizer. The highest (35.50) number of radish roots was obtained at the application of 600kg ha^{-1} followed by the application of 500, 400 and 0kg ha^{-1} with 29.75, 28.25 and 22.00 radish roots respectively.

3.1.7 Length of Root

The length of roots was found (table 4) to be highly significant ($P \leq 0.01$) at the application of different levels of Kalli organic fertilizer. The highest (24.83cm) length of radish roots was obtained at the application of 600kg ha^{-1} followed by the application of 500, 400 and 0kg ha^{-1} with 21.33, 19.43 and 15.51cm radish roots length respectively.

3.1.8 Diameter of Root

No significant effect of Kalli organic fertilizer was observed on root diameter of Radish statistically. However, the application of 600kg ha^{-1} gave the highest (4.28cm) root diameter of Radish.

3.1.9 Weight of Root

The result of the effect of Kalli organic fertilizer on weight of root is presented in Table 4. The result indicates that there was significant ($P \leq 0.01$) effect of Kalli organic fertilizer on weight of root. The application of 600kg ha^{-1} measured the highest (7.20kg) weight of radish while the application of 500kg ha^{-1} was at par with 400 and 0kg ha^{-1} with the least weight.

Table 4: Yield Characteristics of Radish As Influenced By Different Levels of Kalli Organic Fertilizer on the Jos Plateau

Treatment (kg/ha)	Number of Root	Length of Root (cm)	Diameter of Root (cm)	Weight of Root (kg)
0	22.00a	15.51 ^a	3.39a	4.85a
400	28.25ab	19.43 ^b	3.71a	5.15a
500	29.75ab	21.33 ^b	3.81a	5.43a
600	35.50b	24.83 ^c	4.28a	7.20b
SE±	2.05	0.62	0.31	0.42
CV(%)	21.38	17.99	17.60	21.39
LS	*	**	NS	**

Means that do not share a letter are significantly different, SE = Standard Error, CV = Coefficient of Variation, LS = Level of Significance, * = Significant at 0.05, ** = Significant at 0.01

3.1.10 Total Yield

The result from Figure 1 reveals that the highest (18000kg ha^{-1}) total yield of Radish was measured at the application of 600kg ha^{-1} followed by the application of 500, 400 and 0kg ha^{-1} with 13575, 12875 and 12125kg ha^{-1} respectively.




Figure 1: Effect of Kalli Organic Fertilizer on Total Yield of Radish

3.2 Discussion

3.2.1 Effect of Kalli organic fertilizer on the growth of radish

The findings from this research work reveal that there was no significant effect of Kalli organic fertilizer on the plant height, number of leaves. But the leaf area was found to be significant at 8WAP. This shows that the growth characteristics of Radish are not affected by the application of Kalli organic fertilizer. This result is contrary to Satish (2016) who observed that the plant height and number of leaves of Radish was significantly increased by various treatments of organic manure at all the growth stages. Result revealed that plant height, number of leaves, tuber length, tuber diameter and pest resistance were not

significantly affected by the applications of varying levels of vermicast (Eric and Politud, 2016). According to the previous both the deficit and excess amount of nutrients in the soil that could be caused by mineral fertilization can be compensated for by the application of organic fertilizers, which is in line with the results obtained in the current study (Sisay *et al.*, 2008). Different organic fertilizers and application rates significantly influenced the number of leaves during the first 8 weeks after thinning in 2005 and 2006 except at 8 weeks after thinning for the organic fertilizer in 2006 (Alice, 2008).

3.2.2 Effect of Kalli organic fertilizer on the yield of radish

The number of roots, length of roots, weight of root and total yield was found to be significant ($P \leq 0.05$) at the application of different levels of Kalli organic fertilizer. The application of 600kg ha^{-1} gave the highest (35.50) number of radish roots, highest (24.83cm) length of radish roots, highest (7.20kg) weight of radish and highest (18000kg ha^{-1}) total yield of Radish. This might be due to the availability of the nutrients in readily available form. Radish total root yield was significantly affected due to effect of organic manure (Satish, 2016). These yield components were significantly affected by the varying levels of vermicast (Eric and Politud, 2016). Yield components of carrot were also increased in response to the increased rate of combined “orga” and urea fertilizer application (Sisay *et al.*, 2008). Alice (2008) stated that the yield (fresh and dry mass) was not significantly influenced during the first year. In the second year, the fresh mass was only significantly influenced by the rate of organic fertiliser and not the type. Dry mass of carrots in the same year was positively influenced by organic fertiliser.

4.0 Conclusion

Based on this research study it can be concluded that the application of Kalli (600kg ha^{-1}) organic fertilizer significantly increases the yield of Radish. Thus it is recommended that 600kg ha^{-1} of organic fertilizer can be applied for optimum production of Radish.

Reference

- Alice NM. Influence of Organic Fertilizers on The Yield And Quality Of Cabbage And Carrots. Degree of Magister Scientiae Agriculturae, Faculty of Natural and Agricultural Sciences Department of Soil, Crop and Climate Sciences University of the Free State Bloemfontein. 2008
- Eric R, Politud R. Growth and Yield Performance of Radish (*Raphanus sativus* L.) ‘cv’ 'SNOW WHITE' in Response to Varying Levels of Vermicast Applications *International Journal of Scientific and Research Publications*, Volume 6, Issue 5, May 2016, 53 ISSN 2250-3153 www.ijsrp.org
- Ibrahim Iro I., Jameela A, Ninani KN. Growth and Yield Components of Groundnut (*Arachis hypogea* L) as Affected by Phosphorous Fertilizer Application on the Jos Plateau. *Asian Journal of Research in Agriculture and Forestry*, 3(3): 1 – 8, 2019; ISSN: 2581 – 7418. DOI:10.9734/AJRAF/2019/v3i330039
- Kalli Organic Fertilizer Manual. Dharul Hijra Fertilizer Company Limited, KM 10 Bwari – Jere Road, Gnam, Kaduna, Nigeria. 2017.
- Kummeling IC, Thijs MH, Vcr BE, SMjders J, Penders f, Stelma R, Yanrcc PA, Van Den Brant Dangrave PC. Consumption of Organic Food and Risk of a Topic Diseasew during the First 2 Years of life in the Motherlands. *B.J. Nutrion*. 2008, 00. 508-605.

- 251 PCARRD. Phillipine Recomends for Radish, 2007 Available; <http://mboard.pcaTd.dost.gov.ph>
252 [/forum.view topic.php/id-11385](http://mboard.pcaTd.dost.gov.ph/forum.view topic.php/id-11385).
253 Satish D. Effect of different organic manure on growth and yield of Radish (*Raphanus sativus* L.). MSc.
254 Thesis, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior R.A.K College of
255 Agriculture Sehore 466001 (M.P.), 2016.
- 256 Sisay H, Tilahun S, Nigussie D. Effect of combined application of organic-P and inorganic-N fertilizers
257 on yield of carrot. *African Journal of Biotechnology* Vol. 7 (1), pp. 027-034, 4 January, 2008
258 Available online at <http://www.academicjournals.org/AJB> ISSN 1684–5315