

The cubital tunnel syndrome due to a synovial cyst : a case report

Abstract:

The cubital tunnel syndrome is a compressive neuropathy of the ulnar nerve around the elbow, is the most common neurocompression syndrome in the upper limb after carpal tunnel syndrome. Ulnar nerve entrapment is usually located under the thickening of the proximal fascia between two heads of the ulnar flexor of the carpus or arcade of Osborne at distal part of the epicondylar curve. However, it could also be caused by occupying masses along the course of the nerve.

This report shows a case of cubital tunnel syndrome caused by ulnar entrapment with medial synovial cyst. Epidemiologic, pathogenesis and diagnosis are discussed and literature was reviewed.

Keywords:

Ulnar nerve, nerve compression, cubital tunnel syndrome, ulnar nerve release, synovial cyst.

INTRODUCTION:

Cubital tunnel syndrome is a symptomatic ulnar nerve dysfunction at the level of the elbow resulting from a combination of compression, traction, and friction. It's one of the most prevalent upper extremity compressive neuropathies. It's the second most common nerve compression syndrome in peripheral nerve compression disease [1]. Although, potential ulnar nerve entrapment can occur multiple points along its course, such as the arcade of struthers, the medial intermuscular septum, the medial epicondyle, the cubital tunnel and the deep flexor pronator aponeurosis, the most common site of entrapment is the cubital tunnel and has been referred to as the tardy ulnar nerve palsy in the past [2,3]. However, it could also be caused by the occupying masses along the course of ulnar nerve. We present a rare case of cubital tunnel syndrome caused by a synovial cyst. The aim of this work is to study the different epidemiological, etiopathogenic, clinical and treatment of this rare disease. A literature review was made.

Case report:

A 58-year-old gentleman who had suffered with progressive paraesthesia on left upper extremity visited our hospital. Tingling sensation on medial two fingers and the ulnar side of the limb developed one year ago. There was no history of previous trauma at the left elbow, neither history of diabetes nor vascular and neuropathic diseases. On direct questioning he reports the worseness of the tingling and the loss of strength 6 months ago. Clinically he had atrophy of the hypothenar and interosseous muscle in the left hand. Froment's sign was positive. Tinel's sign was also positive at the medial side of the elbow. Sensory examination revealed loss of normal discrimination on the entire little finger and the ulnar side of the ring finger. Elbow radiographs didn't show degenerative changes nor proliferative osteophyte nor calcific disposition. The nerve conduction studies with electromyography revealed evidence of ulnar nerve compression at the elbow. Ultrasonography revealed a cystic mass that compressed the ulnar nerve at the level of cubital tunnel.

In order to relieve the nerve compression, surgery was planned. After induction of general anesthesia, interior sweeping 10 cm incision around medial epicondyle was made. The subcutaneous tissues were dissected, identifying the ulnar nerve who was flattened and compressed in the cubital tunnel by a synovial cyst whose diameter was 2 cm. We resected completely the cyst. After further exploration we didn't find another point of compression. The arcade of Struthers, the Osborne's fascia and the deep flexor pronator aponeurosis were divided. After that, an anterior subcutaneous transposition of the nerve was performed to avoid re-compression by the cyst recurrence. Synovial cyst was confirmed by histopathological examination.



Figure1 : clinical photograph demonstrating a claw hand deformity and intrinsic muscle atrophy.



Figure2 :preoperative ultrasonography shows ulnar nerve compressed by anechoic mass.



figure3 : intra operative photograph showing the cyst compressing the ulnar nerve in the cubital tunnel.



figure4 : intra operative photograph showing total removal of the cyst.

Discussion :

Cubital tunnel syndrome is a common and debilitating condition that can be overlooked until after nerve damage has occurred [11].

The elbow cyst with cubital tunnel syndrome is rarely reported in the literature. The prevalence rate is 3~8% [7], and there was occasional case reports in Europe and the United States [8].

Kato [10] thought that the morbidity rate of the disease is up to 8% in a large sample of 472 patients.

Diagnoses of cubital tunnel syndrome is made through a combination of history, physical examination, and an electroneurographic examination .

Paresthesia is anticipated in the little finger and ulnar half of the ring finger. Weakness of the interossei, the adductor pollicis, and the ulnar lumbrical muscles, which occurs with

advanced disease, may cause characteristic posturing in the hand (Wartenberg sign, Froment sign, and claw hand deformity, respectively) [14].

Cubital tunnel syndrome may be categorized as mild, moderate, or severe disease. Common grading systems include modified McGowan and Dellon classifications [1].

The use of imaging, such as sonography and MRI, to detect the cause of compressions, including those, such as substantial or cystic mass that are rare and unexpected [2,12,13].

The ulnar nerve may be compressed at multiple points along its course, the most common location is at the elbow [9].

There are 4 sites where the ulnar nerve is frequently vulnerable to compression. These are (1) the arcade of Struthers (medial intermuscular septum), (2) the ulnar groove, (3) the humeroulnar arcade (or cubital tunnel), and (4) the exit point between the 2 heads of FCU. Among these locations, the most common sites of compression are the ulnar groove and humeroulnar arcade [18].

Cubital tunnel syndrome could be caused by soft-tissue masses such as ganglia [5] synovial chondromatosis [6] which might compress the ulnar nerve as it lies in the condylar groove or within the cubital tunnel.

In previous reports, synovial cysts were related with the post-traumatic osteoarthritis of the elbow and can cause a relatively acute onset of cubital tunnel syndrome [6].

Conservative treatment measures focus on pain relief, inflammation reduction, and rehabilitation. This includes patient education and behavior modification, non-steroidal anti-inflammatories, night splints, elbow pads, physical therapy, ultrasound, pulsed signal therapy, and corticosteroid injections.

If conservative management is not successful in preventing progression of impairment after several months, surgery may be required.

There are still debates on which surgical procedure is optimal for the treatment of cubital tunnel syndrome. Some authors favor anterior transposition [15], whereas others report high success rates with simple decompression [16] or medial epicondylectomy [17].

For the treatment, though in situ decompression was considered as a good choice in recent years, the decompression with subcutaneous transposition was better procedure for avoiding re-compression caused by the cyst recurrence after the cysts were removed.

CONCLUSION:

Cubital tunnel syndrome is a common condition that can be overlooked until after nerve damage has occurred. Timely recognition and treatment is paramount to good clinical outcomes to avoid irreversible muscle atrophy and functional deficit. In our case, a rarely aetiology, synovial cyst,

caused the compression. This soft-tissue masse was susceptible to be missed and ignored during the treatment using in situ decompression with the mini incision. Regular incision is benefit for finding some little mass to ovoid the second operation for the incomplete release or recurrent symptomtology.in our opinion decompressing the nerve with subcutaneous transposition was good procedure for avoiding recompression caused by eventually cyst recurrence.

References

- 1: Boone, Sean, Richard H. Gelberman, and Ryan P. Calfee. "The Management of Cubital Tunnel Syndrome." *The Journal of Hand Surgery* 40, no. 9 (September 2015): 1897–1904.
- 2 : Chang, Wen Kai, Yong Ping Li, Deng Feng Zhang, and Bing Sheng Liang. "The Cubital Tunnel Syndrome Caused by the Intraneural or Extraneural Ganglion Cysts: Case Report and Review of the Literature." *Journal of Plastic, Reconstructive & Aesthetic Surgery* 70, no. 10 (October 2017): 1404–8.
- 3: Wojewnik, Bart, and Randy Bindra. "Cubital Tunnel Syndrome – Review of Current Literature on Causes, Diagnosis and Treatment." *Journal of Hand and Microsurgery* 01, no. 02 (September 5, 2016): 76–81.
- 4: Palmer BA, Hughes TB. Cubital tunnel syndrome. *J Hand Surg Am* 2010;35:153e63.
5. Kato H, Hirayama T, Minami A, et al. Cubital tunnel syndrome associated with medial elbow ganglia and osteoarthritis of the elbow. *J Bone Jt Surg Am* 2002;84-A:1413e9.
6. Kim CH, Kim SH, Kim MS, et al. Cubital tunnel syndrome, associated with synovial chondromatosis. *J Korean Neurosurg Soc* 2008;43:109e10.
7. Özgür Taşkapılıoğlu , Ahmet Bekar , M.erim Kürşat , Ulviye Yalçinkaya. Ulnar nerve entrapment neuropathy due to extraneural ganglia at the elbow: A case report. *Journal of Neurological Sciences (Turkish)*. 2005;22(1)079-083.
8. H Kato, T Hirayama, A Minami, N Iwasaki, K Hirachi. Cubital tunnel syndrome associated with medial elbow ganglia and osteoarthritis of the elbow. *J Bone and Joint Surg*, 2002; 84(8): 1413-1419.
9. Karatas A, Apaydin N, Uz A, Tubbs R, Loukas M, Gezen F. Regional anatomic structures of the elbow that may potentially compress the ulnar nerve. *J Shoulder Elbow Surg*. 2009;18(4):627–631.

10. Ming jie Gao, Jingxian Pei, Jian feng Bai. Ulnar nerve compression caused by forearm ganglion cyst: clinical analysis of 12 cases. *China drug and clinical*. 2010;10(1):83-84.
- 11- Robertson C, Saratsiotis J. A review of compressive ulnar neuropathy at the elbow. *J Manip Physiol Ther*. 2005;28(5):345.
12. B Gregoli, C Bortolotto, F Draghi. Elbow nerves: normal sonographic anatomy and identification of the structures potentially associated with nerve compression. A short pictorial-video article. *J Ultrasound*. 2013;16:119-121.
13. HJ Chiou, YH Chou, SP Cheng , et al. Cubital tunnel syndrome:Diagnosis by high-resolution ultrasonography. *J Ultrasound Med*. 1998;17(10):643-648.
- 14- Root CG, London DA, Schroeder NS, Calfee RP. Anatomical relationships and branching patterns of the dorsal cutaneous branch of the ulnar nerve. *J Hand Surg Am*. 2013;38(6):1131e1136.
- 15-Kleinman WB : Cubital tunnel syndrome: anterior transposition as a logical approach to complete nerve decompression. *J Hand Surg Am* 24 : 886-897, 1999.
- 16-Pavelka M, Rhomberg M, Estermann D, Loscher WN, Piza-Katzer H : Decompression without anterior transposition : an effective minimally invasive technique for cubital tunnel syndrome. *Minim Invasive Neurosurg* 47 : 119-123, 2004.
- 17- Muermans S, De Smet L : Partial medial epicondylectomy for cubital tunnel syndrome : Outcome and complications. *J Shoulder Elbow Surg* 11 : 248-252, 2002.
- 18- Robertson C, Saratsiotis J : A review of compressive ulnar neuropathy at the elbow. *J Manipulative Physiol Ther* 28 : 345, 2005.