

PHYLOGENETIC RELATIONSHIP OF BACTERIAL SPECIES INVOLVED IN BIOREMEDIATION OF HYDROCARBON POLLUTED SOILS

Abstract

The bio-stimulation of hydrocarbon degrading microbial population in soil using agricultural wastes was carried out. Top soil (0-25cm depth) from three points were bulked to form compost soil samples, Six kilogramme each of the composite soil sample was weighed and transferred into 150 plastic bucket with drainage holes at the base. The soil in each plastic bucket was spiked with 300ml crude oil and amended with different concentrations of agro-wastes and allowed for duration of 30, 60 and 90days. The soil samples were then collected and analysed for both total heterotrophic bacterial and fungal counts and crude oil utilizing bacterial and fungal counts. Data collected were subjected to a three way analysis of variance and significant means were separated using Least Significant Difference Test at 5% probability level. The result showed that the application of the amendments increased the bacterial and fungal counts in the soil at different treatment levels. However, a higher proliferation rate was observed with bacteria and fungi species exposed to higher waste concentrations compared to their counterparts exposed to lower waste concentrations. The phylogenetic relationship of the hydrocarbon degrading bacterial species shows that the identified bacteria were in two clusters: cluster 1 consist of *Bacillus cereus*, *Bacillus thuringiensis*, *Bacillus altitudinis*, *Pseudomonas aeruginosa*, *Proteus mirabilis* and *Proteus penneri*, while cluster 2 consist of *Serratia marcescens*, *Providencia rettgeri* and *Enterobacter asburiae*. The bacterial species obtained shows a greater relationship, this imply that the similarity of the bacterial species could be the reason for their high proficiency in degrading the hydrocarbons in the soil.

Key-words: Phylogenetic, Soil, Bio-stimulation, Hydrocarbon, microbial, agro-waste, degradation

Introduction

Oil exploration activities in producing communities have inadvertently affected the natural land occupied by a rich biodiversity of plants and animal species. The lost in biodiversity of plants and animals species have resulted in abandonment of waste land, with an alarming outcry by the occupancy of those community. At present the waste land at Ogoni is receiving remediation attention by the federal government. Bioremediation involves the application of

microorganism in contaminated site to reducing or removing the contaminant from the soil, water or air. Bioremediation is a mechanism for removing pollutants from polluted environment thus restoring the original natural environment (Sasikuma and Papmazath, 2003). The treatment of polluted environment is necessary because it helps to protect the soil, water and the people in the ecosystem. Bioremediation by means of stimulating the growth of microorganisms have achieved a wide success, in that many waste land has been restored. Biostimulation over the years have exploited the potential use of agricultural wastes in the enhancement of microbial growth in the soil. Agbor *et al.* (2013) in their study reported the potentials of plantain peels and cocoa pod husks in increasing the microbial population of the soil which culminated in the reduction of hydrocarbon content of the soil. Biostimulation using inorganic fertilizer has been extensively employed worldwide in reclaiming oil polluted soil (Jobson *et al.* 1974; Dibble and Bartha 1979; and Song *et al.* 1990 and USEPA 1990; Ekpo *et al.*, 2012). This current research examined the biostimulation potentials of various agro-wastes in enhancing microbial growth in the soil.

Materials and Methods

Microbial analysis was done in the Department of Microbiology (University of Calabar) and Nigerian Institute of Medical Research (Lagos State), while sequencing analysis was done at Inqaba Biotechnology Pty South Africa. Nigerian light crude oil was obtained from NAOC (Nigerian Agip Oil Company), located in Port Harcourt, Rivers State, Nigeria. Agro-wastes such as groundnut husks (GH), maize cobs (MC), empty fruit bunch of oil palm (EFBOP) and cassava peels (CP) were collected from local farmers and processing industries in Cross River State, Nigeria. The collected agro-wastes (GH, MC, EFBOP, CP) were sun-dried for 10 days and blended to powder using electric blender (Model 4250, Braun, Germany). The dispersants were sieved to pass through 2 mm sieve. They were labelled and stored in containers. Top soils (0-25cm depth) were evenly obtained from three (3) points, using a Dutch auger, then homogenized to form composite soil sample. Six kilograms (6kg) of the soil samples was weighed and transferred into each of the hundred and fifty (150) labelled plastic buckets (PB) with drainage holes at the sides and based. The arrangement of the PB was in triplicate using completely randomized design (CRD). Artificial pollution was done by introducing 300ml (0.3 liters) of crude oil into PB containing the soils, except the pristine soils samples that served as the positive control. The PB containing the polluted soils

were mixed thoroughly and allowed to stand for 14 days (these was to allow indigenous microorganisms to become acclimatized with the new soil condition). The amendments were applied in single and combined forms using the following concentrations: pristine control (unpolluted (positive), 0%), crude oil control (polluted (negative), 0%), 3.33% and 6.67% and 10% of the amendments (5 levels). Soil samples for microbial analysis were collected at 30, 60 and 90days.s

Calculation of treatment in percentage

$$PT = \frac{QOW}{QS} \times 100 \text{ where}$$

PT = Percentage of treatment

QOW = Quantity of organic wastes

QS = Quantity of soil

Substitute in the equation

$$0\% \text{ treatment} = \frac{0g}{6000g} \times 100$$

$$3.33\% \text{ treatment} = \frac{200g}{6000g} \times 100$$

$$6.67\% \text{ treatment} = \frac{400g}{6000g} \times 100$$

$$10\% \text{ treatment} = \frac{600g}{6000g} \times 100$$

Soil microbial analysis

Soil samples were taken to the Laboratory of Department of Microbiology for determination of bacterial and fungal populations in the soil using surface plating method.

Determination of total heterotrophic bacteria

The spread plate method using nutrient agar (NA) was used in determining the total heterotrophic bacteria count in the soil according to APHA (1998). Serial dilutions were prepared with one gram of soil on a ten-fold. From the test tube of 10^{-6} and 10^{-7} 0.1ml of the dilutions were spread on the plates in duplicate. Discrete bacterial colonies were counted after incubation at 28°C for 24 hours and the counts were calculated based on colony forming unit per gram of soil.

Determination of total fungi

The spread plate method using Sabouraud dextrose agar (SDA) was adopted according to the procedure of APHA (1998). Serial dilutions were prepared with one gram of soil on a ten-fold. From the test tube of 10^{-6} and 10^{-7} 0.1ml of the dilutions were spread on the plates in duplicate. Discrete fungal colonies were counted after incubation at 28°C for 48 hours and calculated based on colony forming unit per gram of soil sample used.

Determination of crude oil-utilizing bacteria (CUB)

The procedure of Hamamura *et al.* (2006) was adopted. The viable count method using the surface spreading technique was used. About 1.5% agar was added to the mineral salts medium to solidify. A ten (10) fold serial dilutions with 1g of soil was prepared and 0.1ml of 10^{-5} and 10^{-6} dilution were spread on the plates in duplicates. After agar plates inoculation, a sterile Whatman No.1 filter paper was dipped into crude oil, allowed to dry and was placed on the inside of the lid (cover) of the Petri dishes. The saturated filter paper containing crude oil acted as a source of carbon and energy for growth of the bacterial through vapour phase transfer (Okpokwasili & Amanchukwu, 1988). The plates were inverted and incubated at normal temperature for 5 days, during which the CUB were counted from the duplicate plates and expressed as colony forming unit per gram of soil samples.

Determination of crude oil-utilizing fungi

The procedure of Hamamura *et al.* (2006) was adopted. The viable count method using the surface spreading technique was used. About 1.5% agar was added to the mineral salts medium to solidify. A ten (10) fold serial dilutions with 1g of soil was prepared and 0.1ml of 10^{-4} and 10^{-6} dilution were spread on the plates in duplicates. After agar plates inoculation, a sterile Whatman No.1 filter paper was dipped in crude oil, allowed to dry and was placed on the inside of the lid (cover) of the Petri dishes. The saturated filter paper containing crude oil acted as a source of carbon and energy for growth of the fungi through vapour phase transfer (Okpokwasili & Amanchukwu, 1988). The plates were inverted and incubated at normal temperature for 7 days, during which the CUF were counted from the duplicate plates and expressed as colony forming unit per gram of soil samples (CFU/g). Isolate purification was done on a freshly prepared Sabourand dextrose agar (SDA) plate.

Purification and maintenance of microbial isolates

The bacterial isolates obtained from mineral salt medium were purified by repeated sub-culturing. The isolates were subjected to series of transfers unto fresh medium. The bacterial isolates were transferred onto fresh nutrient agar medium and incubated at 28°C for 24 hours. Pure colonies of bacterial were maintained on slopes of nutrient agar (NA) and stored in a refrigerator at 4°C till needed for studies.

Characterization and identification of the isolates using traditional methods

Standard inocula was prepared from the preserved stock culture by taking a loopful of the isolates and aseptically inoculating onto sterile nutrient agar (NA) plates. The plates were incubated at 28°C for 24 hours. The characterization of the isolates was performed, by employing Gram staining reaction, oxidase test, catalase, citrate test, urease test, coagulase test, TSI (triple sugar iron agar) test, MIO (motility indole ornithine) test and methyl red and Voges Proskauer test as described in Bergey's Manual of Determinative Bacteriology, 9th edition (1994). The fungal isolates were identified using the wet mount method and

lactophenol in cotton-blue. Fungal identification was followed the scheme of Hunter & Bennett (1973).

DNA extraction

DNA Extraction was performed at the Anaerobe Laboratory, Molecular Biology and biotechnology Division, Nigerian Institute of Medical Research Yaba Lagos. Methodology was based on PCR and metagenomics analysis. While sequencing analysis was done at a Inqaba Biotechnology Pty South Africa.

DNA extraction was from a 24 hours' growth of microbial isolates in Nutrient broth harvested by centrifugation at 14, 000 x g for 10 minutes. The cells were washed three times in 1 ml of Ultra-pure water by centrifuging at 12,000 rpm for 5 min. DNA extraction and purification was done using ZR Fungal/Bacterial DNA MiniPrep™50 Preps. Model D6005 (Zymo Research, California, USA). 50-100 mg of bacterial cells was re-suspended in 200 µl of sterile water. This was transferred into a ZR Bashing Bead™ Lysis Tube. Exactly 750 µl Lysis Solution was added to the tube. The bead containing the solution was secured in a bead beater fitted with a 2 ml tube holder assembly and process at maximum speed for 5 minutes. The ZR Bashing Bead™ Lysis Tube was Centrifuged in a microcentrifuge at 10,000 x g for 1 minute. 400 µl of the supernatant was pipeted into a Zymo-Spin™ IV Spin Filter in a Collection Tube and centrifuged at 7,000 x g for 1 minute. This was followed by the addition of 1,200 µl of Fungal/Bacterial DNA Binding Buffer into the filtrate in the Collection Tube. After this 800 µl of the mixture was transferred into a Zymo-Spin™ IIC Column in a Collection Tube and centrifuge at 10,000 x g for 1 minute.

The flow through was discarded from the Collection Tube and the process was repeated to obtain the remaining products. The 200 µl DNA Pre-Wash Buffer was added into the Zymo-Spin™ IIC Column in a new Collection Tube and centrifuge at 10,000 x g for 1 minute. This was followed by the addition of 500 µl Fungal/Bacterial DNA Wash Buffer into the Zymo-Spin™ IIC Column and centrifuged at 10,000 x g for 1 minute. The Zymo-Spin™ IIC Column was transferred into a clean 1.5 ml microcentrifuge tube and 100 µl of DNA Elution Buffer was then added directly to the column matrix. This was centrifuged at 10,000 x g for 30 seconds to elute the DNA. The Ultra-pure resulting filtrate (DNA) obtained was used as a template during the assay. This was transported in ice to the laboratory for sequencing.

DNA Sequencing

DNA sequencing was performed by Sanger (dideoxy) Sequencing Technique to determine the nucleotide sequence of the specific microorganism isolated using automated PCR cycle-Sanger Sequencer™ 3730/3730XL DNA Analyzers from Applied Biosystems (Russell, 2002; Metzenberg 2003). This results were obtained as nucleotides when amplified using sequencing primer -16S: 27F: 5'-GAGTTTGATCCTGGCTCAG-3' and 518R: 5'-ATTACCGCGGCTGCTGG-3' (Weisburg *et al.*, 1991). Sequence data from resultant nucleotides base pairs was downloaded and read using FinchTv software followed by direct nucleotide blasting on <http://blast.ncbi.nlm.nih.gov>. For every set of isolate, a read was BLASTED and the resultant top hits for every BLAST result showing species name was used to name the specific organism. Corresponding gene back accession number and query length of sequences blasted was also recorded.

Statistical analysis

Data collected were subjected to a three-way analysis of variance and significant means were separated to least significant difference test at 5% probability level.

RESULTS

Total heterotrophic bacterial (THB) counts of the soil enhanced with agro-wastes

Microbial population in any polluted soil environment is an important tool in determining the degradation process of the petroleum hydrocarbon in the soils. The results obtained for the bacterial population of the soil showed that soil treated with 3%, 6% and 10% CasP₁₄P + EFBOP₁₄P and polluted soils treated with 6% and 10% EFBOP₁₄P had significantly high ($P < 0.05$) bacterial counts with insignificant difference ($P > 0.05$) in the average THB counts as compared to the positive and negative control and also the other treatment groups (Table 1). These were followed by polluted soil treated with 3%, 6%, 10% GnH₁₄P + MaC₁₄P, and GnH₁₄P + EFBOP₁₄P, 10% CasP₁₄P, 6% EFBOP₁₄P, 10% GnH₁₄P + CasP₁₄P and 6%, 10% CasP₁₄P + MaC₁₄P which had insignificant difference ($P > 0.05$) in the average THB counts of the soil. The next were the counts obtained from soil treated with 10% GnH₁₄P, 6% CasP₁₄P, 3% CasP₁₄P + MaC₁₄P and 6% GnH₁₄P + CasP₁₄P which had no variation in the mean THB counts of the soils but the counts were greater than the counts obtained from soil amended with 3% CasP₁₄P, 3% GnH₁₄P + CasP₁₄P, 10% MaC₁₄P and 6% MaC₁₄P + EFBOP₁₄P which had no variation

Table 1: Effect of treatment levels on the microbial population of crude oil polluted-soils

Parameters	Treatment levels	THBC (CFU/g)	TFC (CFU/g)	CUBC (CFU/g)	CUFC (CFU/g)
GnH ₁₄ P	PC	6.71 ^h ±1.97 x10 ⁶	3.92 ^d ±1.49 x10 ⁶	5.60 ^h ±1.56 x10 ⁶	4.21 ^f ±0.99x10 ⁶
	COC	4.48 ⁱ ±1.64 x10 ⁶	3.43 ^d ±1.23 x10 ⁶	4.07 ⁱ ±2.27x10 ⁶	3.43 ^f ±1.07x10 ⁶
	3%	1.01 ^g ±4.9 x10 ⁷	4.48 ^d ±1.49 x10 ⁶	9.81 ^f ±4.79x10 ⁶	5.16 ^f ±3.91x10 ⁶
	6%	1.38 ^e ±5.32x10 ⁷	5.57 ^c ±1.88 x10 ⁶	1.11 ^e ±3.75x10 ⁷	6.06 ^e ±3.39x10 ⁶
MaC ₁₄ P	10%	1.84 ^c ±8.67x10 ⁷	5.79 ^c ±1.83 x10 ⁶	1.32 ^d ±5.31 x10 ⁷	6.81 ^e ±3.36x10 ⁶
	3%	9.69 ^g ±5.83 x10 ⁶	5.17 ^c ±1.24 x10 ⁶	8.16 ^g ±4.13 x10 ⁶	4.80 ^f ±1.61x10 ⁶
	6%	1.27 ^f ±10.46x10 ⁷	6.78 ^c ±2.33 x10 ⁶	1.11 ^e ±8.78 x10 ⁷	6.69 ^e ±1.58x10 ⁶
GnH ₁₄ P+MaC ₁₄ P	10%	1.58 ^d ±6.5x10 ⁷	7.72 ^c ±1.40 x10 ⁶	1.26 ^d ±8.64 x10 ⁷	7.37 ^e ±2.17x10 ⁶
	3%	1.97 ^{bc} ±4.10x10 ⁷	6.29 ^c ±2.23x10 ⁶	1.40 ^d ±6.25 x10 ⁷	7.10 ^e ±4.40 x10 ⁶
	6%	2.12 ^b ±4.21 x10 ⁷	7.03 ^c ±1.26 x10 ⁶	1.63 ^b ±5.96 x10 ⁷	8.26 ^d ±5.37 x10 ⁶
CasP ₁₄ P	10%	2.27 ^b ±8.2 x10 ⁷	8.18 ^c ±2.33 x10 ⁶	1.77 ^a ±6.95x10 ⁷	9.71 ^c ±4.9 x10 ⁶
	3%	1.72 ^d ±7.77 x10 ⁷	5.01 ^c ±3.01 x10 ⁶	5.91 ^h ±3.53 x10 ⁶	3.74 ^f ±1.85 x10 ⁶
	6%	1.94 ^c ±11.66 x10 ⁷	5.89 ^c ±2.89 x10 ⁶	7.56 ^g ±5.35 x10 ⁶	4.41 ^f ±1.62 x10 ⁶
EFBOP ₁₄ P	10%	2.24 ^b ±3.16 x10 ⁷	6.39 ^c ±1.94 x10 ⁶	8.84 ^g ±5.18 x10 ⁶	4.98 ^f ±1.78 x10 ⁶
	3%	2.18 ^b ±6.44 x10 ⁷	6.0 ^c ±1.29 x10 ⁶	6.29 ^h ±2.13 x10 ⁶	4.41 ^f ±1.92 x10 ⁶
	6%	2.57 ^a ±10.3 x10 ⁷	9.38 ^a ±2.41 x10 ⁶	8.64 ^g ±2.13 x10 ⁶	5.04 ^f ±2.30 x10 ⁶
	10%	2.64 ^a ±8.77x10 ⁷	1.04 ^a ±3.37 x10 ⁷	1.01 ^f ±8.34 x10 ⁷	5.93 ^e ±1.64 x10 ⁶

Table 1: Continues

Parameters	Treatment levels	THBC (CFU/g)	TFC (CFU/g)	CUBC (CFU/g)	CUFC (CFU/g)
CasP ₁₄ P+ EFBOP ₁₄ P	3%	2.57 ^a ±5.63x10 ⁷	7.14 ^c ±2.24 x10 ⁶	8.78 ^g ±7.89 x10 ⁶	5.12 ^f ±2.28 x10 ⁶
	6%	2.66 ^a ±5.77 x10 ⁷	8.43 ^c ±6.05 x10 ⁶	1.03 ^f ±8.49 x10 ⁷	5.84 ^e ±2.8 x10 ⁶
	10%	2.73 ^a ±6.19 x10 ⁷	1.09 ^a ±5.69 x10 ⁷	1.18 ^e ±8.35 x10 ⁷	6.30 ^e ±3.25 x10 ⁶
GnH ₁₄ P+ EFBOP ₁₄ P	3%	1.97 ^{bc} ±6.73 x10 ⁷	5.79 ^c ±1.72 x10 ⁶	8.87 ^g ±3.13 x10 ⁶	4.84 ^f ±1.56 x10 ⁶
	6%	2.21 ^b ±6.59 x10 ⁷	8.08 ^c ±2.37 x10 ⁶	1.12 ^e ±5.07 x10 ⁷	5.86 ^e ±2.01 x10 ⁶
	10%	2.36 ^b ±6.43 x10 ⁷	9.78 ^a ±2.44 x10 ⁶	1.29 ^d ±7.16 x10 ⁷	6.31 ^e ±1.89 x10 ⁶
CasP ₁₄ P+MaC ₁₄ P	3%	1.88 ^c ±7.61 x10 ⁷	5.01 ^c ±3.01 x10 ⁶	1.32 ^d ±9.22 x10 ⁷	8.38 ^d ±2.87 x10 ⁶
	6%	2.19 ^b ±11.87 x10 ⁷	5.89 ^c ±2.37 x10 ⁶	1.52 ^c ±10.95 x10 ⁷	1.08 ^b ±4.6 x10 ⁷
	10%	2.37 ^b ±13.1x10 ⁷	6.39 ^c ±1.94 x10 ⁶	1.74 ^a ±14.18 x10 ⁷	1.32 ^a ±6.69 x10 ⁶
MaC ₁₄ P+ EFBOP ₁₄ P	3%	1.43 ^e ±5.57 x10 ⁷	5.2 ^c ±2.92 x10 ⁶	1.10 ^e ±9.4 x10 ⁷	6.40 ^e ±2.59 x10 ⁶
	6%	1.66 ^d ±5.32 x10 ⁷	6.53 ^c ±2.85 x10 ⁶	1.35 ^d ±10.18 x10 ⁷	7.90 ^e ±3.76 x10 ⁶
	10%	1.84 ^c ±5.72 x10 ⁷	7.12 ^c ±2.89 x10 ⁶	1.65 ^b ±9.83 x10 ⁷	1.01 ^c ±6.19 x10 ⁶
GnH ₁₄ P+CasP ₁₄ P	3%	1.70 ^d ±6.31 x10 ⁷	7.97 ^c ±3.04x10 ⁶	6.41 ^h ±2.81 x10 ⁶	3.97 ^f ±1.30 x10 ⁶
	6%	1.88 ^c ±6.65 x10 ⁶	9.02 ^a ±5.50 x10 ⁶	8.39 ^g ±4.33 x10 ⁶	4.69 ^f ±1.72 x10 ⁶
	10%	2.07 ^b ±7.02 x10 ⁶	1.04 ^a ±4.61 x10 ⁷	1.00 ^f ±4.85 x10 ⁷	5.08 ^f ±1.60 x10 ⁶

Mean with the same superscript along the vertical arrays indicate no variation

Legend:

GnH₁₄P Groundnut husks 2014 powder
MaC₁₄P Maize cob 2014 powder
EFBOP₁₄P Empty fruit bunch of oil palm 2014 powder
CasP₁₄P Cassava peels 2014 powder
PC Pristine control
COC Crude oil control

in the mean THB counts of the soils (Table 1). The soils treated with 3% GnH_{14}P and 3% $\text{MaC}_{14}\text{P} + \text{EFBOP}_{14}\text{P}$ had significantly reduced ($P < 0.05$) mean THB counts

in the soils but significantly increase ($P < 0.05$) than the average THB counts obtained in the pristine control and the crude oil control soils. However, the crude-oil control soil was observed to produce the lowest bacterial population compared to the pristine control and the treated groups (Table 1). Figure 1 present the results of the THB counts obtained at different duration of study and the results showed that the average THB counts of the soil treated with $\text{EFBOP}_{14}\text{P}$ and $\text{CasP}_{14}\text{P} + \text{EFBOP}_{14}\text{P}$ at 30days and 60days were significantly increase ($P < 0.05$) than other amended groups with no variation in the mean values. These were followed by soil treated with CasP_{14}P at 60 days, $\text{CasP}_{14}\text{P} + \text{EFBOP}_{14}\text{P}$ at 90 days, $\text{GnH}_{14}\text{P} + \text{MaC}_{14}\text{P}$ at 60 days, $\text{GnH}_{14}\text{P} + \text{EFBOP}_{14}\text{P}$ and $\text{CasP}_{14}\text{P} + \text{MaC}_{14}\text{P}$ at 30th and 60th day with insignificant difference ($P > 0.05$) in the average THB counts of the soils, these counts were more than the counts obtained from the soil treated with $\text{EFBOP}_{14}\text{P}$ and $\text{GnH}_{14}\text{P} + \text{MaC}_{14}\text{P}$ at 90 days with no variation in the mean values obtained. These were also more than the mean THB counts obtained from soils treated with $\text{GnH}_{14}\text{P} + \text{MaC}_{14}\text{P}$, CasP_{14}P at 30 days, and $\text{GnH}_{14}\text{P} + \text{EFBOP}_{14}\text{P}$ at 90 days with no variation the mean values obtained, more than the counts obtained from the soil treated with GnH_{14}P , MaC_{14}P , $\text{MaC}_{14}\text{P} + \text{EFBOP}_{14}\text{P}$ and $\text{GnH}_{14}\text{P} + \text{CasP}_{14}\text{P}$ at 30th, 60th and 90th day of treatment duration. Figure 1 showed that the total heterotrophic bacterial counts at 60 days were higher than the count obtained at 30 days, while the counts obtained for 30 days were more than the THB counts at 90 days. However, from the result obtained it was observed that the soil amended with $\text{CasP}_{14}\text{P} + \text{EFBOP}_{14}\text{P}$ had higher ($P < 0.05$) mean THB counts in the soil than other treatment groups but followed by soil treated with $\text{EFBOP}_{14}\text{P}$ (Figure 1). It was also observed that the combined treatment produced higher ($P < 0.05$) THB counts than the single amended groups. These imply that polluted soil amelioration is best achieved with the combined treatments.




FIG.1: Total heterotrophic bacterial counts at different durations of soil treatment

Legend:

- MaC₁₄P Maize cob 2014 powder
- EFBOP₁₄P Empty fruit bunch of oil palm 2014 powder
- CasP₁₄P Cassava peels 2014 powder
- DAST Days after soil treatment

Total fungal (TF) counts in soil enhanced with agro-wastes

The result as presented on Table 1 showed that the TF in soil treated with 6% and 10% EFBOP₁₄P, 10% CasP₁₄P + EFBOP₁₄P and soil treated with 6% and 10% GnH₁₄P + CasP₁₄P had more counts than the pristine soil (PS +ve), crude oil polluted soil (COPS -ve) and other treatment groups. These were followed by soils treated with 6%, 10% GnH₁₄P, 3%, 6%, 10% MaC₁₄P, GnH₁₄P + MaC₁₄P, CasP₁₄P + MaC₁₄P, MaC₁₄P + EFBOP₁₄P, 3% and 6% GnH₁₄P + EFBOP₁₄P and 3% GnH₁₄P + CasP₁₄P with no variation in the mean TF counts of the soils, these counts were also more than the counts obtained in the PS and COPS soil. These results imply that the treatment of the polluted soil with agro-wastes increase the TF counts of the soils than the counts obtained in the controls. The PS and COPS had no significant variation in the mean TF counts of the soil. Figure 2 showed the results of the TF counts of the soil due to duration of the study, it was observed that the average TF counts of the soil treated with CasP₁₄P + MaC₁₄P at 60 days were more than other amended groups and durations. These were also followed by soil ameliorated with CasP₁₄P + MaC₁₄P at 30days of soil treatment. The soil treated with CasP₁₄P + EFBOP₁₄P was next, higher ($P < 0.05$) than the counts obtained from soil enhanced with EFBOP₁₄P at 60 days of soil treatment, these were followed by other treated soils that showed no variation in the mean TF counts of the soils. The increased in fungal counts were observed to be significantly higher at 60 days of the study than other study period. These results could imply that the complete acclimatization of the fungal population exposed to the agro-wastes was achieved at 60 days of soil study. The fungal population in soil treated with CasP₁₄P + MaC₁₄P were higher than other treatment groups, while the combined form of the different agro-wastes resulted in high fungal counts than the single treatments.




FIG. 2: Total fungal counts in polluted soils treated with agro-wastes

Legend:

MaC₁₄P Maize cob 2014 powder

EFBOP₁₄P Empty fruit bunch of oil palm 2014 powder

CasP₁₄P Cassava peels 2014 powder

DAST Days after soil treatment

Crude oil-utilizing bacterial (CUB) counts in soils amended with agro-wastes

The determination of crude oil utilizing bacteria was also assessed, to ascertain the populations of the bacteria that are purely hydrocarbon degraders in the soils. The results as presented in Table 1 showed that the soil treated with 10% GnH₁₄P + MaC₁₄P and CasP₁₄P + MaC₁₄P had more CUB counts, with no variation in mean values obtained. These were followed by CUB counts soils treated with 6% GnH₁₄P + MaC₁₄P and 10% MaC₁₄P + EFBOP₁₄P with no variation in the average values but higher than the count obtained from soil treated with 6% CasP₁₄P + MaC₁₄P. These were also followed by soil treated with 10% GnH₁₄P, 10% MaC₁₄P, 3% GnH₁₄P + MaC₁₄P, 10% GnH₁₄P + EFBOP₁₄P and 3% CasP₁₄P + MaC₁₄P, with no variation in the average values but higher than the counts obtained from soil treated with 6% GnH₁₄P, 6% MaC₁₄P 10% CasP₁₄P + EFBOP₁₄P, 6% GnH₁₄P + EFBOP₁₄P and 3% MaC₁₄P + EFBOP₁₄P with insignificant difference in the average CUB counts of the soils. The CUB counts obtained from the agro-wastes treated soils were observed to be higher than the count obtained from the pristine and crude oil control soils. However, the pristine soil had higher CUB counts than the crude oil controls.

The result as presented in Figure 3 showed that the soil treated with GnH₁₄P + MaC₁₄P and CasP₁₄P + MaC₁₄P at 60 days had significantly increase ($P < 0.05$) in CUB counts, with no variation in the average values obtained. These were higher than the counts obtained in soil amended with CasP₁₄P + MaC₁₄P at 30 days, MaC₁₄P + EFBOP₁₄P and GnH₁₄P + CasP₁₄P at 60 days with insignificant differences ($P > 0.05$) in the average CUB counts obtained in the soils. The counts obtained from GnH₁₄P + MaC₁₄P at 90 days was next, higher than the counts obtained from the soil treated with GnH₁₄P + MaC₁₄P at 30 days of soil treatment. It was also observed that the counts obtained from the soils treated with GnH₁₄P, MaC₁₄P, CasP₁₄P + GnH₁₄P, GnH₁₄P + GnH₁₄P, CasP₁₄P + MaC₁₄P at 60 days and MaC₁₄P + EFBOP₁₄P at 30 days soil treatment had insignificant difference ($P > 0.05$) in mean CUB counts of the soils. The results for CUB counts in the soil at 90 days of soil remediation were decreased as compared to the high counts recorded at 60 days, followed by the count obtained at 30 days of soil remediation. The results as presented on Fig. 3 also showed that the CUB counts of the soil treated with GnH₁₄P + MaC₁₄P and CasP₁₄P + MaC₁₄P were significantly increased than other amended soils.

Crude oil-utilizing fungal counts in soils treated with agro-wastes

Crude-oil contamination reduces the population of non hydrocarbon utilizing fungal species in soil and survival of the hydrocarbon utilizing fungal is reduced. The degradation process would take a longer time to complete, as a mean of accelerating the rapid growth of hydrocarbon-utilizing fungi left in soil, agro-wastes at varying levels were applied in the soils. The results obtained showed that the soils treated with 10% CasP₁₄P + MaC₁₄P had increased CUF counts, these were followed by the counts obtained in soils treated with 6% CasP₁₄P + MaC₁₄P, also followed by soils treated with 10% GnH₁₄P + MaC₁₄P and 10% MaC₁₄P + EFBOP₁₄P with insignificant difference ($P > 0.05$) in the mean values obtained.

These were more than the count obtained in soils treated with 3% CasP₁₄P + MaC₁₄P and 6% GnH₁₄P + MaC₁₄P which had insignificant differences (P>0.05) in the average CUF counts obtained in the soils. The soils treated with 3%, 6% GnH₁₄P, 3%, 6% MaC₁₄P, 3% GnH₁₄P + MaC₁₄P, 10% EFBOP₁₄P, 6%, 10% CasP₁₄P + EFBOP₁₄P, 6% and 10% GnH₁₄P + EFBOP₁₄P, and 3%, 6% MaC₁₄P + EFBOP₁₄P had insignificant differences (P>0.05) in the average fungal counts obtained in soils but significantly higher than the CUF counts obtained in soils treated with 3%, 6%, 10% CasP₁₄P, and 3%, 6%, 10% EFBOP₁₄P and the pristine and crude oil polluted soils (Table 1). The results as presented on Table 1 showed that no variation existed between the CUF counts obtained in the pristine and the crude oil polluted soils.




FIG.3 : Crude oil utilizing bacterial counts in polluted soils amended with agro-wastes

Legend:

MaC ₁₄ P	Maize cob 2014 powder
EFBOP ₁₄ P	Empty fruit bunch of oil palm 2014 powder
CasP ₁₄ P	Cassava peels 2014 powder
DAST	Days after soil treatment

The results obtained for the crude oil-utilizing fungi at different duration of study indicated that there were differences in the mean values obtained. The soils treated with CasP₁₄P + MaC₁₄P at 60 days of study showed significantly higher ($P < 0.05$) mean fungal counts. These were followed by the fungal counts obtained in soil treated with GnH₁₄P + MaC₁₄P at 60 days, CasP₁₄P + MaC₁₄P at 30 days and 90 days and MaC₁₄P + EFBOP₁₄P at 60 days with no variation in the mean CUF counts obtained in the soil but more than the mean fungal counts obtained in soils treated with GnH₁₄P + MaC₁₄P at 30 days, MaC₁₄P + EFBOP₁₄P and GnH₁₄P + CasP₁₄P at 30 days of soils treatment. The results as presented in Figure 4 indicated that the crude oil utilizing fungal population obtained from soils treated with the single amendment was reduced as compared with some of the amendments in combined form. Figure 4 also showed that the CUF counts obtained at 60 days were more than the counts obtained at 30 days and 90 days.

Identification of microorganisms in the soils

The bacterial species identified using the traditional methods of identification (biochemical test) were as follows: *Bacillus* spp, *Proteus* spp, *Chromobacterium* sp, *Serratia* sp, *Pseudomonas* sp, *Streptococcus* sp, *Escherichia coli*, *Micrococcus* sp, *Achromobacter* sp, and *Providencia* sp (Table 1). The fungal genera that were identified in the soils (Wet-mount and Lacto-phenol in cotton blue) were as follows: *Mucor* sp, *Aspergillus* sp, *Rhizopus* sp, *Penicillium* sp, *Trichoderma* sp, *Yeast* sp and *Fusarium* sp.

TABLE 2

Probable microorganisms identified in soil using traditional methods

Agro-wastes	Bacteria	Fungi
PS	<i>Bacillus</i> sp <i>Proteus</i> <i>Chromobacterium</i> sp	<i>Mucor</i> sp <i>Aspergillus</i> sp
COPS	<i>Bacillus</i> sp <i>Serratia</i> sp	<i>Rhizopus</i> sp <i>Mucor</i> sp
GnH ₁₄ P	<i>Pseudomonas</i> sp <i>Bacillus</i> sp <i>Streptococcus</i> sp	<i>Aspergillus</i> sp <i>Mucor</i> sp <i>Penicillium</i> sp
MaC ₁₄ P	<i>Escherichia coli</i> <i>Proteus</i> spp <i>Pseudomonas</i> sp <i>Enterobacter</i> sp	<i>Mucor</i> sp <i>Trichoderma</i> sp
GnH ₁₄ P+MaC ₁₄ P	<i>Bacillus</i> spp <i>Micrococcus</i> sp <i>Achromobacter</i> sp <i>Pseudomonas</i> sp	<i>Rhizopus</i> sp <i>Penicillium</i> sp
CasP ₁₄ P	<i>Bacillus</i> sp <i>Serratia</i> sp	<i>Aspergillus</i> sp <i>Mucor</i> sp
EFBOP ₁₄ P	<i>Pseudomonas</i> sp <i>Proteus</i> sp	<i>Aspergillus</i> sp <i>Mucor</i> sp
CasP ₁₄ P+ EFBOP ₁₄ P	<i>Providennia</i> sp <i>Pseudomonas</i> sp	<i>Penicillium</i> sp <i>Aspergillus</i> sp
GnH ₁₄ P+ EFBOP ₁₄ P	<i>Bacillus</i> sp <i>Proteus</i> sp	<i>Trichoderma</i> sp <i>Rhizopus</i> <i>Yeast</i> sp
CasP ₁₄ P+MaC ₁₄ P	<i>Bacillus</i> sp <i>Serratia</i> sp	<i>Mucor</i> sp <i>Rhizopus</i>
MaC ₁₄ P+ EFBOP ₁₄ P	<i>Proteus</i> sp <i>Enterobacter</i> sp <i>Bacillus</i> sp	<i>Penicillium</i> sp <i>Aspergillus</i> sp <i>Fusarium</i> sp
GnH ₁₄ P+CasP ₁₄ P	<i>Pseudomonas</i> sp <i>Bacillus</i> sp	<i>Aspergillus</i> sp <i>Rhizopus</i> sp




FIG. 4: Crude oil-utilizing fungal counts in polluted soils amended with agro-wastes

Legend:

- MaC₁₄P Maize cob 2014 powder
- EFBOP₁₄P Empty fruit bunch of oil palm 2014 powder
- CasP₁₄P Cassava peels 2014 powder
- DAST Days after soil treatment




FIG. 5: Microbial population in soils enhanced with different agro-wastes

Legend:

MaC₁₄P Maize cob 2014 powder

EFBOP₁₄P Empty fruit bunch of oil palm 2014 powder

CasP₁₄P Cassava peels 2014 powder

Molecular identification of Microbial isolates

The molecular analyses method of identifying bacterial to species level have been found to be more reliable than the traditional approaches because the technique depends on the

examination of genetic diversity of isolates (Table 3). The maximum parsimony method was used; otherwise BIONJ method with MCL distance matrix was used. The scale bar of the trees represents a 0.1% difference in nucleotide sequences. The phylogenetic relationship of the hydrocarbon degrading bacterial species shows that the identified bacteria were in two clusters: cluster 1 consist of *Bacillus cereus*, *Bacillus thuringiensis*, *Bacillus altitudinis*, *Pseudomonas aeruginosa*, *Proteus mirabilis* and *Proteus penneri*, while cluster 2 consist of *Serratia marcescens*, *Providencia rettgeri* and *Enterobacter asburiae*. Between cluster 1 and 2 the results imply that there is genetic variability between the hydrocarbon degrading bacteria isolated from the soil environment. The *Bacillus cereus*, *Bacillus thuringiensis* and *Bacillus altitudinis* are closely related by 99% similarity. *Pseudomonas aeruginosa*, *Proteus mirabilis* and *Proteus penneri* by are related to each other by 94% similarity. While *Proteus mirabilis* and *Proteus penneri* are 100% similar. *Serratia marcescens* in cluster 2 are related by 100% with *Providencia rettgeri* and *Enterobacter asburiae*. While *Enterobacter asburiae* and *Providencia rettgeri* are closely related by 36% (Fig. 6). The high percentage similarities that the bacteria isolated have with the blast revealed that they are closely related to each other.

Table 3: Sequence identification of fungi species detected in agro-wastes amended soil samples

Sample No.	Query Length	Gene bank accession No	Identity of Isolate obtained
1	1064		No- significant similarity found
2	1122	JQ308547.1	<i>Bacillus cereus</i>
3	1022	EF633995.1	<i>Bacillus cereus</i>
4	993	EF434507.1	<i>Pseudomonas aeruginosa</i>
5	696	FR717839.1	<i>Proteus mirabilis</i>
6	1136	KC150144.1	<i>Proteus mirabilis</i>
7	1144	KC344360.1	<i>Proteus mirabilis</i>
8	1117	KC150144.1	<i>Proteus mirabilis</i>
9	1130	LC002198.1	<i>Proteus mirabilis</i>

10	1054		No- significant similarity found
11	1137	KF933674.1	<i>Bacillus pumilus</i>
		Same as	<i>Same as</i>
		KF933678.1	<i>Bacillus altitudinis</i>
12	1104	HQ259936.1	<i>Proteus penneri</i>
13	1141	KF938667.1	<i>Serratia marcescens</i>
14	1148	KC172019.1	<i>Providencia rettgeri</i>
15	1116	JX941572.1	<i>Bacillus thuringiensis</i>
16	Failed PCR Amp.		<i>DNA may not be present</i>
17	1138	KJ877656.1	<i>Enterobacter asburiae</i>
18	1152	KJ398213.1	<i>Proteus penneri</i>
		Same as	<i>Same as</i>
		JN384144.1	<i>Proteus vulgaris</i>
19	1076	EU161995.1	<i>Bacillus thuringiensis</i>
			<i>Same as</i>
		GU329917.1	<i>Bacillus cereus</i>
20	1130	KC150144.1	<i>Proteus mirabilis</i>




Figure 6: Phylogenetic relationship among hydrocarbon degrading bacterial species

Discussion

Petroleum hydrocarbon pollution has been known and reported by various scientists to produce adverse effect on biodiversity and abundance of soil microbes (Atlas and Bartha 1973). Microbial degradation is an important aspect in bioremediation of polluted soil. The degradation potentials of microorganisms depend solely on their abundance in the soils. The reduction or removal of petroleum hydrocarbons from soils depend on the ability and the metabolic capability of the microorganisms to break down the hydrocarbon components in the polluted soils (Not all microorganisms can degrade hydrocarbons in soil but all hydrocarbon utilizing organisms possessed specific abilities to degrade hydrocarbon components in polluted soils). The results of the baseline analysis showed that the abundance

of the bacteria and fungi in soil after contamination were significantly higher than the population of the microorganisms in soils before pollution with crude oil.

The bacterial genera present in the pristine soil were as follows: *Bacillus* sp, *Chromobacterium* sp and *Proteus* sp. While the fungal genera present in the soil were: *Aspergillus* sp and *Mucor* sp. Soil contamination produces higher bacterial and fungal counts than the natural soil without crude oil. The abundance of soil microbes in the contaminated soil compared to the low counts obtained in the ordinary soil may be due to the presence of crude oil that may served as energy and carbon source to microbes. Atlas (1981) in a similar study observed that the availability of petroleum product in the soil could cause a high proliferation of microorganisms. Amending the contaminated soil with agricultural wastes bio-stimulated the growth of the soil microbes to breakdown hydrocarbons in soil. Evans *et al.* (2004) reported the diversity of bacteria community present in soil microcosms through an effective application of bio-stimulation principle. The most populated microbial species in the soils were Gram positive bacteria during the study. Chikere *et al.* (2011) reported that gram positive bacteria contribute largely to the bioremediation of polluted soils. The findings of Olabisi *et al.* (2009) also showed that the most frequently occurring bacterial genera in hydrocarbon polluted soils treated with melon shell were; *Bacillus*, *Micrococcus*, *Pseudomonas* and *Acinetobacter*. The Gram positive bacteria, due to their metabolic capabilities in degrading environmental pollutant could be considered as best organisms in bioremediation studies. The high advantage attributed to the gram positive bacteria could be due to their metabolic abilities, resilience in highly polluted environments and ability to produce bio-surfactants (Hamamura *et al.*, 2006).

The high content of nitrogen and phosphorus in the agro-wastes used in the study could be one of the major factors that contributed to the high bacterial and fungal population in the soil. Agbor *et al.* (2012) reported that availability of adequate quantity of nitrogen and phosphorus in the soil, bio-stimulated microbial degradation of hydrocarbon products. The compatibility of the soil due to the presence of hydrocarbons could be reduced through the application of adequate levels of agro-wastes which could serve as bulking agent. The loosening of the compacted soil may produce soil aeration, thus providing suitable environmental condition for aerobic microorganisms to attack the hydrocarbons in the soil. Stephen *et al.* (2013) observed significant difference ($P > 0.05$) in microbial counts in diesel-polluted soil amended with cowpea chaff. Akpe *et al.* (2015) reported that hydrocarbon degrading bacteria were more in soils with amended plantain peels and guinea corn shaft than the un-amended soil samples. Bossert & Bartha (1994) reported a significant reduction in microbial counts in polluted soils and attributed the reduction to toxic compounds and certain harmful metabolites present in the pollutant. The success of bio-stimulation of microbial population in the degradation of contaminants in soil enhanced with organic manure such as NPK, urea fertilizer and chicken droppings was reported by Chikere *et al.*, (2009). The high levels of the treatment applications resulted in an increased proliferation of microbial counts in the amended soil compared to those with lower treatment levels. However, among the single amendment used, groundnut husks powder showed a higher microbial population counts compared to other single amendment used, but the combined form of cassava peel

powder + maize cob produced higher microbial counts than the single amendment. This revealed that the combined form of the agro-wastes used possessed high bio-utilization capabilities in the degradation of pollutants in soil. However, the efficiency of the combined wastes could be attributed to the acclimated microorganisms or nutrients found in the single amendments. The report by Okolo *et al.* (2005) and Ebere *et al.* (2011) supports that the combined forms of agricultural wastes in the appropriate measures could result in an effective biodegradation of hydrocarbon products in the terrestrial environment. The soil treated with the agricultural wastes at different levels showed significant increases ($p < 0.05$) in the hydrocarbon-utilizing bacterial and fungal population in the soil. The soil treated with 6% of CasP₁₄P + MaC₁₄P produced higher microbial population in the soil compared to the single amendment at higher treatment application levels. This result is an indication that high treatment levels of the agro-wastes influence higher microbial proliferation in the soil than the single lower treatment levels. The results from this study suggest that microbial degradation is achievable at a faster rate when agro-wastes are applied at the right proportion, especially in a combined form.

Microbial degradation of crude oil polluted soil

The molecular techniques used in the identification of microorganisms have been found to be more appropriate than the traditional approaches because these involve culture-independent analyses such as genotyping the 16S rRNA genes, fluorescence *in situ* hybridization (FISH) (Zwirgmaier *et al.*, 2005), the use of genetic probes, polymerase chain reaction (PCR) and metagenomics (Josephine *et al.*, 1991). Molecular identification involving rRNA enables the identity of novel sequences and diversity. Maslow *et al.* (1993) reported that 16S rRNA analysis for characterization of bacteria is excellent because the technique depends on the examination of genetic diversity of isolates. Olukunle and Boboye (2012) reported that molecular methods for rapid detection of diverse strains of oil-degrading bacteria for the bioremediation process is of great relevance in promoting a sustainable development of our environment with low environmental impact. Although a wide phylogenetic diversity of microorganisms is capable of aerobic degradation of contaminants, *Pseudomonas* species and closely related organisms have been the most extensively studied owing to their ability to degrade many different contaminants. Walker *et al.* (1976) reported that *Pseudomonas* possessed some more competent and active hydrocarbon-degradative enzymes than other biodegraders.

The high percentage similarities that the bacterial and fungal isolates have with the BLAST means that they are closely related to each other. Some of the bacteria obtained in this study have been isolated by other researchers such as Ojo, (2006) and Boboye *et al.*, (2010) by means of traditional techniques. The 16S rRNA comparison reveals that *Bacillus cereus* and *B. thuringiensis* are closely related to each other, Delvecchio *et al.*, (2006). Annweiller *et al.* (2000) reported that *Bacillus cereus* is more tolerant to high levels of hydrocarbon in soil due to their resistance endospore. Muthuswamy *et al.* (2008) reported that *Bacillus* genera is capable of degrading short carbon chain length in hydrocarbon-contaminated soils. The bacterial species identified in the soils during the bioremediation study using molecular identification methods were identified as *Bacillus cereus*, *Pseudomonas aeruginosa*, *Proteus mirabilis*, *Bacillus pumilus* or *Bacillus altitudinis*, *Proteus penneri*, *Serratia marcescens*, *Providencia rettgeri*, *Bacillus thuringiensis*, *Enterobacteria asburiae* and *Proteus penneri* same as *Proteus vulgaris*.

Conclusion

It is therefore concluded that the agricultural wastes used during the study were effective in enhancing bacterial population in the polluted soils. The phylogenetic analysis of the identified bacteria shows that the bacterial species are closely related with no much evolutionary divergence.

Reference

- Agbor, R.B., Ekpo I.A., Udofia U.U., Okpako E.C. & Ekanem B.E (2012). Potentials of cocoa pod husks and plantain peels in the degradation of total petroleum hydrocarbon content of crude oil polluted soil. *Archives of Applied Science Research*, 4(3): 1372-1375.
- Akpe A. R., Esumeh F. I., Aigere S. P., Umanu G. & Obiazi H. (2015). Efficiency of Plantain Peels and Guinea Corn Shaft for Bioremediation of Crude Oil Polluted Soil. *Journal of Microbiology Research*, 5(1): 31-40
- Atlas, R. & Bartha, R. (1973): Fate and effects of polluting petroleum in the environment. *Research Review* 49: 49 – 75.
- Atlas, R. (1981). *Microbial Ecology*. New York Press; 5(7): 54 – 90.
- Annweiler, E., Richnow, H. H., Antranikian, G., Hebenbrock, S., Garms, C., Franke, S., Franke, W. & Michaelis, W. (2000). Naphthalene degradation and incorporation of naphthalene derived carbon into biomass by the thermophile *Bacillus thermoleovorans*. *Applied Environmental Microbiology*, 66: 518-523
- Boboye, B., Olukunle, O. F., & Adetuyi, F. C. (2010). Degradative activity of bacteria isolated from hydrocarbon-polluted site in Ilaje, Ondo State, Nigeria, *African Journal of Microbiology Research*, 4(23): 2484-2491.
- Bossert, I. & Bartha, R. (1994). The fate of petroleum in soil ecosystem. In: *Petroleum Microbiology*, Macmillan, New York, U.S.A, 435-473.
- Chikere, B. O. & Okpokwasili, G. C. (2009). Enhancement of biodegradation of petrochemicals by nutrient supplementation. *Nigerian Journal of Microbiology*. 17:130-135.
- Chikere, C. B., Okpokwasili & Chikere, B.O. (2011). Monitoring of microbial hydrocarbon remediation in the soil. *Biotechnology*, 1(3): 117-138.
- Delvecchio, V., Connolly, J., Alefantis, T., Walz, A., Quan, M., Patra, G., Ashton, J., Whittington, J., Chafin, R., Liang, X., Grewal, P., Khan, A., & Mujir C. (2006). “Proteomic Profiling and Identification of Immunodominant Spore Antigens of *Bacillus anthracis*, *Bacillus cereus*, and *Bacillus thuringiensis*.” *Applied Environmental Microbiology*, 72(9): 6355–6363.
- Dibble, J.T.; and Bartha, R. 1979. Effects of environmental parameters on biodegradation of oil sludge. *Applied Environmental Microbiology*. 37: 729-739.

- Ebere, J. U., Wokoma, E. C. & Wokocha, C. C. (2011). Enhanced remediation of a hydrocarbon-polluted soil. *Journal of Environmental and Earth Sciences* 3(2): 70 – 74.
- Evans, F. F., Rosado, A. S., Sebastian, G. V., Casella, R., Machado, P. L. O. A. & Holmstrom, C. (2004). Impact Of Oil Contamination And Biostimulation On The Diversity Of Indigenous Bacterial Communities In Soil Microcosms. *FEMS Microbiology Ecology*, 49: 295-305.
- Hamamura, N., Olson, S. H., Ward, D. M. & Inskeep, W. P. (2006). Microbial population dynamics associated crude oil biodegradation in diverse soil. *Applied and Environmental Microbiology*. 72: 6316-6324.
- Jobson, A.; McLaughlin, M.; Cool, F.D.; and Westlake, D.W.S. 1974. Effects of amendments on the microbial utilization of oil applied to soil. *Applied Environmental Microbiology*. 27: 166-71.
- Josephine, K. L., Pillai, S. D., Way, J., Gerba, C. P & Pepper, I. L. (1991). *Soil Science Society of America Journal*. 55: 1326-1332.
- Maslow, J. M., Arimi, S.M & Arbeit, R.D. (1993). Molecular epidemiology: application of contemporary techniques to the typing of microorganisms. *Clinical Infectious Disease*. 17: 153-164.
- Muthusamy, K., Gopalakrishnan S., Ravi T. K & Sivachidambaram, P. (2008). Biosurfactants: properties, commercial production and application, *Current Science*, 94: 736-747
- Ojo, O. A. (2006). Petroleum- utilization by native bacterial population from a wastewater carnal Southwest Nigeria. *African Journal of Biotechnology*., 5(4): 333- 337.
- Olukunle, O. F. & Boboye, B. (2012). Phylogenetic analysis of Oil-degrading Bacteria Associated with Polluted Sites in River State, Nigeria. *Archives of Applied Science Research*, 4 (4): 1600-1608
- Okolo, J. C., Amadi, E. N & Odu, C. T. I. (2005). Effects of soil treatments containing poultry manure on crude oil degradation in a sandy loam soil. *Applied Ecology and Environmental Research* 3(1): 47 – 53.
- Olabisi, L. S., Reich, P. B., Johnson, K. A., Kapuscinski, A. R., Suh, S. & Wilson, E. J. (2009). Reducing greenhouse gas emissions for climate stabilization: Framing regional options. *Environmental Science Technology*. 43: 1696 – 1703.
- Sasikumar, C.S. and T. Papinazeth (2003). Environmental management: bioremediation of polluted environment. *Proceedings of the Third International Conference on Environmental and health*, Chennai, India. pp. 456-469.
- Song, H.; Wang, X.; and Bartha, R. 1990. Bioremediation potential of terrestrial fuel spills. *Applied Environmental Microbiology*. 56: 652-6.
- Stephen, E., Job, O.S. & Abioye, O.P. (2013). Study On Biodegradation Of Diesel Contaminated Soil Amended With Cowpea Chaff. *Journal of Science & Multidisciplinary Research*, 2(1), 14-18

USEPA. 1990. Alaskan Bioremediation Project. Publication EPA/600/889/073, 1-21.

Walker, J. D. & Colwell, R. R. (1976). Enumeration of petroleum degrading microorganism. *Applied and Environmental Microbiology*. 31: 198 – 207.

Zwirgmaier, K., Fichtl, K. & Ludwig, W. (2005). *In Situ* Functional Gene Analysis: Recognition of Individual Genes by Fluorescence *In Situ* Hybridization. *Methods in Enzymology*, 397: 338-351.

UNDER PEER REVIEW