1	<u>Original Research Article</u>
2	
3	Impact of Climate Change on Production of Major Food and Commercial
4	Crops of India: a five decadal study
5	Running Title: Agroecosystems and Climate Change
6	
7	Abstract
8	Climate change is posing a great threat to agriculture and food security, especially in the
9	agriculture oriented and developing countries like India. The present study was carried out to
10	critically study the impact of climate change on productivity of major cereal and commercial
11	crops by statistically analyzing the time series data. The analysis inferred that crop production of
12	both food and commercial crops in India has increased since 1960-61. It was observed that major
13	food crops (rice & wheat) were adversely affected by increase in maximum temperature and
14	decrease in rainfall. The alternative measures such as area under cultivation, irrigation, fertilizer
15	and pesticide consumption were observed to be nullifying that negative impact of climate change
16	by enhancing the overall production. However, the commercial crops were observed to be
17	positively affected by the increasing temperature. The study suggested that although the
18	agriculture sector is able to withstand the adverse impact of climate change till now, but in near
19	future this situation can become reversed. This necessitates the implementation of appropriate
20	adaptation and mitigation measures to deal with the problems of climate change and to ensure

Original Research Article

the food security and food safety along in long run. 21

Keywords: Agriculture, crop production, climate change, adaptation, mitigation 22

23 Introduction

Agriculture production of any country is directly dependent on its climate and weather 24 conditions since minor changes in temperature, precipitation and CO₂ concentration can 25 drastically impact its crop growth. Higher levels of CO₂ generally increase productivity of plants 26 through enhancement in plant photosynthesis due to CO₂ fertilization effect but the long-term 27 28 effects are uncertain and might involve negative effects on plant food web, decreased plant

nutritional values, reduced N content of plant etc (Nogia et al., 2016). Hence, to achieve the
optimum plant productivity, a balance in atmospheric carbon level is primarily needed.

Since agriculture relies greatly on adequate water supply, temperature, and a balance of gases in the atmosphere, farming is most vulnerable to the effects of climate change. Also 80% of the world's arable land is progressively being planted with a handful of crop commodities (corn, soybean, wheat, rice, and others) and that too are grown under "modern monoculture systems" which due to their ecological homogeneity are particularly vulnerable to climate change as well as biotic stresses (Heinemann et al. 2013).

Climate change will have variable impacts across regions and cropping systems. There are concerns that climate change will hamper the world's ability to provide sufficient food for the global population (Hatfield et al., 2011). The impacts on agriculture and food security are more prominent especially in the agriculture oriented and developing countries like India. These countries have limited arable land but heavy dependence on agriculture (Mendelsohn *et al.*, 2006; Stern, 2006; Nelson *et al.*, 2009) and also have poor technological and financial capabilities for mitigation and adaptation to climate change.

India is facing major challenges to increase its food production to the tune of 300 mt by 2020 to 44 45 feed its ever growing population by producing 50% more grain by 2020 (Kumar and Gautam, 46 2014). Climate change also affects other factors of production agriculture, such as water availability, soil fertility, and pests (Porter, 2014). It will aggravate problems with soil loss 47 through wind and water erosion in addition to environmental externalities which are associated 48 49 with current land use practices. The population of India as on March 1st, 2011 stood at 1,210.7 50 million (623.2 million males and 587.5 million females) with a population density of 382 per sq. km. Cropland is the main occupation of the major population. The economy of the region is 51 52 predominantly agrarian as constitutes a measure of livelihood of a large portion of population.

53 The present study was carried out to critically study the impacts of climate change on 54 productivity of major cereal and commercial crops by statistically analyzing the time series data. 55 This research has been conducted taking India as a case study.

56 Materials and Methods

India is the seventh largest country with 2.4% of total area of the world with great physical 57 diversity. The major land use/land cover of the country can be categorized as cropland, built-up, 58 59 forest, open forest, pine forest, scrub land, barren land and water. Mainly rabi and kharif crops are grown with paddy (rice) as kharif crop and whereas Wheat is cultivated during rabi season. 60 The study is based on the secondary data about crop production of major food crops (rice, wheat, 61 coarse cereals, pulses), major commercial crops (ground nut, rapeseed & mustard, sugarcane, 62 cotton (lint), raw jute & mesta), cultivation area and inputs use (Fertilizer, Pesticides & irrigation 63 inputs) since 1960-61 to 2015-16 that has been collected from the Handbook of Statistics on the 64 Indian Economy (2015-16, 2016-17 and 2017-18) being published by Reserve Bank of India, 65 and records of Ministry of Agriculture & Farmers Welfare, Government of India (2016-17). The 66 temperature and rainfall data of the country was retrieved from web portal of Indian 67 Meteorological Department, Ministry of Earth Sciences for the selected study period. The 68 analysis of data was made through descriptive statistics for better interpretation and description 69 of various conditions or scenarios. 70

Multivariate regression analysis was performed to confirm the percentage of the response
variable variation from the predictor variable that is explained by a linear model in Equation:

$$Y = \alpha_0 + \beta 1X1 + \beta 2X2 + \beta 3X3 + \mu$$

74 Where,

- 75 Y= Crop Production
- 76 α_0 = Constant
- 77 X1= Temperature variations
- 78 X2= Rainfall variations
- 79 X3= variations in Cropping Area
- 80

73

Y is the observed Production due to temperature, precipitation and cropping area and β1, β2, and
β3 are coefficients of the temperature, precipitation and cropping area respectively. Similarly,
X1, X2 and X3 are the observed changes in the temperature, precipitation (rainfall) and cropping
area respectively, during the study period.

85 **Results and Discussion**

The analysis of meteorological data revealed an average increment of 0.3°C in annual maximum temperature and an average increment of 35.17 mm in annual rainfall of the country since 1960. Rainfall expressed more fluctuations than temperature during the selected study period. These fluctuations in main climatic variables in countries like India can be an alarming sign for agricultural activities (Fig 1).






Fig 1. Climatograph for India since 1960 to 2015

The overall crop production of both food and commercial crops in India has increased since 1960-61 with some fluctuations in between (Fig 2). Similar observations were inferred from the data on area under cultivation of different crops (Fig 3). Also, it was found that use of irrigation; consumption of fertilizers (N+P+K) and pesticide depicted an increasing trend towards 2015-16 (Fig 4).






Fig 2. Crop production (million tonnes) of (a) major food crops and (b) major commercial
 crops over 55 years



102 Fig 3: Area under cultivation (million hectares) of (a) major food crops and (b & c) major

103 commercial crops over 55 years

104



105

Fig 4. Net irrigated area (million hectares), Fertilizer consumption (lakh tonnes) and
 pesticide consumption ('000 tonnes) over 55 years.

108 **Regression analysis for major food crops**

109

110 Multivariate regression analysis method was applied to identify the impact of various factors on

111 Crop production of different crops in India over the past 55 years. The findings revealed that

different variables affected the production of different crops differently (Table 1).

101

113 The effects of the climatic parameters, i.e. of temperature & rainfall were observed to be 114 detrimental for rice production as the increase in temperature & decrease in rainfall negatively 115 affected Rice crop production. But the increase in cropping area was observed to be combating 116 this negative effect since it contributed positively with a significant increase in Rice crop 117 production. The adjusted R^2 value expressed that 88% variability in Rice production is explained 118 by these variables.

The regression analysis for wheat indicated that increase in maximum temperature and decrease in rainfall affected negatively the production upto some extent while increase in cropping area had positive impact on the Wheat production. The adjusted R^2 value expressed that all these three variables, i.e. temperature, rainfall and cropping area were contributing to 94% variability in Wheat crop production. The increase in minimum temperature, however, was found to be positively affecting the production of both cereal crops.

The increase in maximum temperature as well as decrease in rainfall did not have significant effect on the production of coarse cereals and pulses. These crops are not much dependent on rainfall pattern for their growth. The area under of coarse cereals crops was observed to be decreasing and thereby negatively affecting the production of crop. Adjusted R^2 value expressed that the studied variables were explaining only 57% of the variability of the cereal crop production and 52% in case of production of pulses.

1	3	1
-		-

 Table 1 Regression results for major food crops

	Rice	Wheat	Coarse Cereals	Pulses
	2.045	0.007	2.510	
Max. Temp.	-3.045	-0.695	2.713	3.632
Min. Temp.	16.905	11.691	-2.942	-0.184
Rainfall	-0.015	-0.011	0.014	0.002
Cropping Area	6.278	4.359	-0.607	1.087
Intercept	-408.68	-243.51	13.03	-118.385
Model R ²	0.89	0.94	0.60	0.55
Adjusted R ²	0.88	0.94	0.57	0.52
Observations	56	56	56	56

132

134 **Regression analysis for major commercial crops**

Changes in studied variables did not have significant effect on Groundnut production as they 135 only determined 33% (adjusted R^2) of the variability in crop production. Temperature, rainfall 136 variability and increase in cropping area were observed to have some positive impact on the 137 production of mustard according to the regression results with high adjusted R^2 value (91%). 138 Similar results were obtained in case of Sugarcane and Cotton crop production with high R² 139 values of 98% and 86% respectively. Although, temperature, rainfall and cropping area seemed 140 141 to positively affecting the production of Raw Jute & Mesta production but their determination potential towards production was only 47%. Here, the increase in minimum temperature was 142 143 observed to be negatively affecting the production of rapeseed and mustard, sugarcane and cotton (Table 2). 144



Table 2: Regression Results for major commercial crops

	Groundnut	Rapeseed & Mustard	Sugarcane	Cotton (Lint)	Raw Jute & Mesta
Max. Temp.	1.435	1.332	7.240	7.549	2.094
Min. Temp.	0.717	-0.053	-2.417	-3.004	2.417
Rainfall	0.006	9.52E-05	0.023	0.010	0.001
Cropping Area	0.415	1.360	91.827	5.345	1.954
Intercept	-60.36	-40.85	-300.41	-209.17	-103.29
Model R ²	0.38	0.93	0.98	0.87	0.51
Adjusted R ²	0.33	0.92	0.98	0.86	0.47
Observations	56	56	56	56	56

146

The results of the study inferred that impact of the climatic factors is different in context of 147 different crops in the last 55 years. In case of Rice, variation in climatic factors affected 148 negatively the crop production as indicated by the regression model result. It was observed 1°C 149 unit rise in temperature can affect Rice production by 3% decrease and one-unit decrease in 150 rainfall will affect it with slight decrease in production by 0.01%. The reason behind it is the 151 availability of other sources of irrigation to the cultivars as indicated by the Irrigation pattern 152 results (Fig. 4) and therefore is not so much dependence on rainfall directly for Rice production. 153 For Wheat, temperature has negative significance as 1°C increase in temperature will lead to 154 around 0.64% decrease in production as Wheat is a winter season (when temperature remains 155

around 10°C) crop. Through the analysis of the impact of climatic factors on commercial crops it was observed that, changes in climatic factor do not have significant impact in case of groundnut while increase of 1°C in temperature may lead to 7.24% and 7.54% increase in sugarcane and cotton production respectively.

But here, the important thing to note is that some other factors also affect the overall production of crops. In case of India, by the analysis of crop production of various crops over the past 55 years with considering all the possible factors, it was estimated that overall production of many crops has increased despite of negative impacts of climatic factors on certain crops as these are combated by other adaptation measures such as increase of irrigation, fertilizer, pesticides inputs and increase in the area under the cultivation of crops (Fig 4 and 2).

166 **Conclusion and Recommendations**

167 The technological advances along with investments in irrigation, infrastructure and institutions in 168 last five decades have supported India to come out of the food security syndrome and promoted 169 its level in the International agricultural market. However, curbing the problem of feeding ever 170 increasing growing population still remains a challenge in terms of producing more and more 171 food. Moreover, the projections of climate change impacts towards 2100 have suggested 172 significant changes in temperature and rainfall will lower the rice yield 15% and wheat yield by 173 22% (Birthal et al, 2014). For India with limited arable land, the situation can become much worse if proper adaptation and mitigation measures in agriculture sector are not taken into 174 consideration. 175

176 It has been suggested that local weather conditions (rain, temperature, sunshine and wind) in 177 combination with locally adapted plant varieties, cropping systems, and soil conditions can 178 maximize food production if plant diseases can effectively be controlled (Kumar and Gautam, 179 2014). Adaptations to experienced and projected climate changes are already occurring (Moser 180 and Ekstrom 2010) and but will still need to continue even if mitigation efforts are widely 181 implemented (IPCC 2014).

182 The overall findings of this study indicated that climate variables have differential impact on the 183 production of different crops. It was observed that major food crops (rice & wheat) were adversely affected by increase in maximum temperature and decrease in rainfall. However, the commercial crops were observed to be positively affected by the increasing temperature. These conditions may also lead to increased weed & pest proliferations. Moreover, increased temperature quickly ripens the crop which results in malnourished crop and less nutrient food. Further, the study concluded that area under the cultivation of different crops, use of selective inputs such as irrigation, fertilizer, pesticides etc. has also increased during the last 55 years.

190 Thus, it can be concluded that although climatic variables have significant impacts on various food and commercial crops but the alternative measures are nullifying that negative impact by 191 192 enhancing the overall production. Further, there is an urgent need to take coordinated steps in 193 direction of adaptation and mitigation towards climate change to ensure the food security and food safety in long run. India has already pledged to address the global climate challenge as a 194 195 responsible nation, both at domestic and international level. The Government of India has launched its National Mission for Sustainable agriculture with a focus on soil and water 196 197 conversation, water use efficiency, soil health management, and rain-fed area development. Also, 198 two other programs viz., agro-meteorology advisory service and farmers' awareness program 199 have been launched to scale up sustainable agriculture program (Tripathi and Mishra, 2017).

From the results of the study, it is concluded that although the agriculture sector is able to 200 combat the adverse impacts of climate change till now, however, in near future this situation can 201 202 become reversed. This necessitates the implementation of appropriate measures to deal with the problems of climate change. The first and foremost need is to provide incentives to promote 203 networks and/or to form clubs that bring likeminded farmers together on same platform for 204 communication and adaptation strategies as a response to climate change (Tripathi and Mishra, 205 206 2017). The knowledge about changes in climate especially the fluctuations in temperature and rainfall patterns should be spread at farmer level. Farmer decision making ability can have a 207 208 significant effect on reducing on-farm vulnerability by addressing problems of soil loss and 209 degradation and adoption of soil and water conservation practices (Lehman et al. (2015). Adoption of these practices can improve agroecosystem resilience (Kremen and Miles 2013) by 210 increasing the production of a more diverse range of ecosystem services. 211

212 Farmers should also be provided with crop specific incentives and insurance against climate risks. There is a limited scope left for the expansion of cropping area in context of enormously 213 214 increasing population in the country. Also, the agricultural land is facing various types of degradation. So there is an utmost urgency to give a boost to research and development in the 215 fields for development of high temperature and drought resistant new crop varieties along with 216 the promotion of sustainable agricultural practices. Adoption of some other important measures 217 such as mixed/intercropping, change in planting dates, water harvesting, micro-irrigation, 218 agroforestry etc. should be emphasized. It is further recommended that any programs that are 219 working to minimize the adverse impact of climate change on food crops production should first 220 consider the important cereal crops such as rice and wheat that are the staple food diet for major 221 Indian population and are being most affected by the higher temperatures relative to the other 222 food crops. 223

224 **References**

Birthal PS, Khan MT, Negi DS, Agarwal S. Impact of Climate Change on Yields of Major Food
Crops in India: Implications for Food Security. Agricultural Economics Research Review. 2014;
27 (2):145-155. DOI: 10.5958/0974-0279.2014.00019.6

Hatfield JL, Boote KJ, Kimball BA, Ziska LH, Izaurralde RC, Ort D, Thomson AM, Wolfe D.
Climate impacts on agriculture: Implications for crop production. Agronomy Journal.
2011;103(2):351-70.

Heinemann JA, Massaro M, Coray DS, Agapito-Tenfen SZ, Wen JD. Sustainability and
innovation in staple crop production in the US Midwest. Int J Agric Sustain. 2013;
doi:10.1080/14735903.2013. 806408

IPCC. Summary for Policymakers. In: C.B. Field et al., editors. Climate Change 2014: Impacts,
Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working
Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.
Cambridge, UK and New York, NY: Cambridge University Press. Pp 1-32. 2014.

- Kremen C, Miles A. Ecosystem services in biologically diversified versus conventional farming
 systems: Benefits, externalities, and trade-offs. Ecology and Society. 2012;17(4):40-65.
- Kumar R, Gautam HR. Climate Change and its Impact on Agricultural Productivity in India. J
 Climatol Weather Forecasting. 2014; 2:109. doi:10.4172/2332-2594.1000109

- Lehman RM, Cambardella CA, Stott DE, Acosta-Martinez V, Manter DK, Buyer JS, Maul JE,
- 243 Smith JL, Collins HP, Halvorson JJ, Kremer RJ, Lundgren JG, Ducey TF, Jin VL, Karlen DL.
- 244 Understanding and enhancing soil biological health: The solution for reversing soil degradation.
- 245 Sustainability. 2015;7:988-1027.
- Mendelsohn R, Dinar A, Williams L. The distributional impact of climate change on rich and
 poor countries. Environment and Development Economics. 2006;11:159-178.
- Moser SC, Ekstrom JA. A framework to diagnose barriers to climate change adaptation. PNAS.
 2010;107(51):22026-31.
- 250 Nelson GC, Rosegrant MW, Koo J, Robertson R, Sulser T, Zhu T, Ringler C, Msangi S, Palazzo
- A, Batka M, Magalhaes M, Valmonte-Santos R, Ewing M, Lee D. Climate Change: Impact on
- 252 Agriculture and Costs of Adaptation. International Food Policy Research Institute. Washington,
- 253 D.C. 2009.
- Nogia P, Sidhu GK, Mehrotra R, Mehrotra S. Capturing atmospheric carbon: biological and nonbiological methods. International Journal of Low-Carbon Technologies. 2016;11: 266–274.
- 256 Porter JR, Xie L, Challinor AJ, Cochrane K, Howden SM, Iqbal MM, Lobell DB, Travasso MI.
- Food security and food production systems. In: Field CB et al. editors. Climate Change 2014:
- 258 Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of
- 259 Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate
- 260 Change. Cambridge, UK and New York, NY: Cambridge University Press. pp. 485–533. 2014
- 261 Stern N. The Economics of Climate Change: The Stern Review. H.M. Treasury, London. 2006.
- Tripathi A, Mishra AK. Knowledge and passive adaptation to climate change: An example from
 Indian farmers. Climate Risk Management. 2017;16: 195–207.
- 264