

Doctors' knowledge on radiation exposure during CT scan prescriptions in Hassan II Hospital of Agadir-Morocco

Abstract

The purpose of this study was to assess physicians' knowledge of radiation doses and potential health risks of radiation exposure from CT.

A standardized questionnaire was distributed to physicians. The questionnaire covered the demographic data of the prescriber, the frequency of referrals for CT scan examinations, the physicians' knowledge of radiation doses, the potential health risks of radiation exposure from CT scan and training on patients' radiation protection.

A total of 72 physicians (55%) completed the questionnaire. 99% of the practitioners' prescribe CT examinations for patients during their exercises but only 10% of physicians use the guideline during CT prescriptions. 38% of prescribers took into account the ratio benefit/risk related to x-rays during radiological exam prescription. While 4% of prescribers' explained the risk related to x-rays to the patients during radiological exam prescription, 14% of physicians have correctly estimated the effective dose received during an abdomen pelvic scan compared to the dose of a standard chest x-ray radiograph in an adult. 54% of doctors underestimated the lifetime risk of fatal cancer attributable to a single computed tomography scan of the abdomen pelvic and 8% of practitioners have received formal training on risks to patients from radiation exposure.

Recurrent training in advanced radiation protection of patients could lead to significant improvements in knowledge and practice of CT prescribers.

Key words: CT Scans, Physicians' knowledge, Patient' Radiation Exposure.

1. Introduction

Every day in each hospital, Physicians use various X-rays technologies to screen diagnose, stage and treat cancers with the aim of saving thousands of lives. The use of CT in medical diagnosis delivers radiation doses to patients which are higher than those from other radiological procedures.

The biological effects of low doses received during medical diagnostic imaging can cause harm. The cancer radiogenic is well documented^{1,2}, indeed the lifetime attribute to the risk of cancer is 1 for every 82 in high-use groups³ and 1 in every 1000 CT abdomen pelvic examination⁴. For example, in the United Kingdom, it has been estimated that 100 to 250 death cases occur each year because of the radiological exposures^{5,6}.

In any diagnostic procedure the dose of radiation delivered should be^{7, 8} enough to answer the relevant clinical question. Moreover, it should be and as low as reasonably achievable to minimise the risk to the patient.

41 It is very important that physicians who prescribe radiological imaging should be well
42 trained in deciding whether diagnostic imaging is necessary and have an accurate knowledge
43 of the associated risks.

44 The absence of studies on doctors' knowledge in Moroccan Hospitals and the lack of
45 knowledge on the medical exposure per inhabitant in Morocco ^{9,10} initiated us to undertake
46 the current study.

47 The aim of this study is to assess knowledge of patient radiation exposure from CT
48 examinations prescribed in Hassan II Hospital.

49

50 **2. Materials and Methods**

51

52 The concerned population included the prescribers of CT scans in Hassan II hospital. In total,
53 the entire population studied comprised 130 practitioners. The participants in this study have
54 received a standardized questionnaire.

55 The 16 sections of the questionnaire were designed to evaluate the current practice regarding
56 the prescriptions of CT examinations. The questionnaire covered five main areas:

57 The first requested demographic data of prescriber (department, gender, qualification, years of
58 experiences).

59 The second part included questions and aimed at:

- 60 - Prescription frequency of CT scans,
- 61 - The use of medical imaging examinations guide before prescription.
- 62 - Knowledge of benefit /risk ratio of the use of x-rays,
- 63 - Routine patient's information about possible health risks.

64 The third tackled doctors' knowledge on radiation doses which can be assessed into two
65 approaches:

- 66 - Compare the average of effective dose received during Abdomen pelvic CT scan
67 ($D_{AP}^{CT} \sim 11$ mSv) and Radiography Skull ($D_{SC}^R = 0,07$ mSv) examinations ⁴.
- 68 - Evaluate the effective dose received during Abdomen pelvic CT scan examination.

69

70 The fourth dealt with prescribers' knowledge of the risk of cancer induction after one CT scan
71 Abdomen pelvic examination.

72 Finally, we asked doctors if they had already received training with regards to radiation
73 protection.

74 **3. Results**

75 **3.1. The study population**

76 Out of the 130 physicians' practitioners in our hospital, 72 participated in the
77 questionnaire giving a response rate of 55%. There were 42 men (sex ratio 1.4). The study
78 group contained the General practitioners, Interns, Surgeons and Medical specialists. The
79 percentage of each specialty was respectively 10%, 19%, 36% and 37%. The average
80 professional experience for all participants was $10,29 \pm 6,83$ years with 58% of them having
81 more than 10 years of experience.

82 **3.2. Current Prescribers Practice Regarding CT Examinations**

83 99% of respondents to the survey were prescribers' of CT examinations. The
84 physicians' non prescribers were Medical specialists in dermatology.

85 Only 8% of our study group used a guideline for prescribing the less irradiating exam.
86 It was constituted by 33% of Interns and 20% of General practitioners.


87 38% of them said that they always take into account the benefit /risk ratio of X-rays
88 when prescribing a scanner, while 54% sometimes use it and 8% never. The benefit/risk of X-
89 rays is still considered by 42% of senior doctors and only 17% of juniors.

90 Only 4% of practitioners have always informed patients about the probable risks due
91 to their exposure to X-radiation, while 68% did so occasionally and 28% never. Thus only 5%
92 of Physicians seniors have always passed such information to the patient, while Interns
93 represent 0%.

94
95 **3.3. Knowledge of Doses and Health Risks Related to Radiations by Doctors**


96
97 On the assessment of the effective dose received during an abdomen-pelvic CT compared to
98 chest X-ray front, 14% of our practitioners had correctly assessed that dose. 11% had
99 overestimated it while 54% of practitioners had underestimated it and 21% have no answers,
100 regardless of the different specialties (Figure N°1).

101



102 **Figure N°1:** Assessment of knowledge of the effective dose received during an abdomen-pelvic CT
 103 comparatively to an adult chest radiography by physicians per speciality.


104
 105 During the absolute evaluation of the dose delivered during a standard abdomen-pelvic CT,
 106 with reference to natural radiation in Morocco estimated to average 2.5 mSv per year, 8% of
 107 practitioners had correctly assessed the dose. 10% of prescribers had overestimated it, 58%
 108 underestimated it while 21% had expressed no opinion (Figure N°2).



110 **Figure N°2:** Assessment of the knowledge of effective dose received during an abdomen-pelvic CT
 111 comparatively to annual background exposure in Morocco by physicians per speciality.

112
 113 Estimations of doses delivered were misjudged, and the risk of radiation-induced cancer were greatly
 114 underestimated since a large majority of practitioners (58%) had replied that there was no risk of
 115 radiation-induced cancer due to the realization of one abdomen-pelvic CT (Figure N°3).

116 .



117 **Figure N°3:** Assessment of knowledge on radiation-induced cancer after one abdomen-pelvic CT by physicians
 118 per speciality.

119 **3.4. Further Education and Training:**

120 Only 8% of clinicians had already benefited from training in radiation protection of patients. The more
 121 detailed analysis showed that neither Intern, General Doctor nor Surgeon had received training of this
 122 type while only 20% of Medical Specialists have received such training.

123 **4. Discussion**

124 Our study group showed that Physician' knowledge of radiation exposure from medical
 125 imaging is insufficient, and that is due to the fact that they don't inform their patients of the
 126 risks of radiation exposure, and they underestimate radiation exposure of frequently used
 127 diagnostic imaging and the associated risks.

128 • Only 8% of physicians of this study used a guideline during prescriptions of CT exam.
 129 Yet the European directive on the radiation protection for medical purposes requires
 130 justification of the radiological procedure which is one of the necessary steps to obtain
 131 the radiation protection of patients as part of a quality assurance process ¹¹.The lack of
 132 use of referral guidelines could be explained by the Moroccan radiologists by the
 133 absence of national protocols ¹².

134
 135 • 38% of prescribers in our study group took into account the ratio benefit/ risk. This
 136 result is much lower than 70% reported by Gervaise et al. in a similar study for a
 137 population of French hospital doctors ⁴.

138

- 139
- 140
- 141
- 142
- 143
- 144
- 145
- 146
- 147
- 148
- 149
- 150
- 151
- 152
- 153
- 154
- 155
- 156
- 157
- 158
- 159
- 160
- 161
- 162
- 163
- 164
- 165
- 166
- 167
- Only 4 % of our physicians group have explained the x ray risk to the patients during prescription. This result is much lower than 22% reported by Lee et al. in a similar study for a population of emergency physicians in USA ¹³ and than 25% reported by Gervaise et al. in a similar study for a population of French hospital doctors ⁴.
 - The knowledge on radiation doses in our study group is limited. In detail, we asked to compare the average effective dose received during an abdomen pelvic CT scan in adults to a standard chest radiograph. Only 14 % of the study participants answered correctly. This result is lower than 30% reported by Lee et al. in a similar study for a population of emergency physicians in USA ¹³. And it is also less than 32,5 % obtained by Merzenich et al. in a similar study in Germany ¹⁴. It is more than 13 % reported by Gervaise et al. in a similar study for a population of French hospital doctors ⁴.
 - The physicians' knowledge on the lifetime risk for the development of cancer after one abdomen pelvic CT examination was answered correctly by only 42% (approx.1 cancer death per 1,000 deaths) of responds in our study group^{15,16}. This result is higher than 12,5 % reported by Jacob et al. for a population of hospital doctors ¹⁷. It is approximately the same as the 31% obtained by Rice et al. for a population of paediatrics surgeons ¹⁸. It is higher than 39% reported by Gervaise et al. in a similar study for a population of French hospital doctors ⁴.
 - The poor knowledge results achieved in this study could be explained by many factors: About 92% of the questioned doctors reported that they have never undergone formal training on patients' radioprotection. This reflects a poor knowledge of the principles of radiation protection by our clinicians. This result is higher than 75 % reported by Gerben et al. for a physician population of the Australian emergency departments ¹⁹, and higher than 34% reported by Gervaise et al. in a similar study for a population of French hospital doctors ⁴.

168 **5. Conclusion**

169 The objective of this study was to explore the physicians' knowledge on patients' radiation
170 protection during their prescriptions of CT scan examinations. The obtained results showed
171 that the physicians' knowledge on patient's radioprotection is characterized by:

- 172 • 8 % of physicians used a guideline during prescriptions of CT exam.
- 173 • 38% of prescribers took into account the benefit/risk ratio.
- 174 • Only 4 % of our doctors have been explained the x ray risk to the patients during
- 175 prescriptions.
- 176 • 14 % of the physicians have correctly approximated the radiation doses received
- 177 during an abdomen pelvic CT scan.
- 178 • 42% of physicians' have estimated the lifetime risk for the development of cancer
- 179 after one abdomen pelvic CT examination in a correct way.
- 180 • 92% of doctors have never undergone formal training on patients' radioprotection.

181 We recommend training during the university curriculum of interns and also the periodic
 182 ongoing training of all doctors from all specialties with the aim of improving their
 183 understanding of medical radiation exposure.

184 6. References

- 185 1. Ron E. Cancer risks from medical radiation. *Health Phys* 2003; 85: 47-59.
- 186 2. Mather R. The physics of CT dose. Toshiba America Medical Systems Inc.
- 187 3. Griffey RT, Sodickson A. Cumulative radiation exposure and cancer risk estimates in
- 188 emergency department patients undergoing repeat or multiple CT. *AJR Am J Roentgenol*
- 189 2009; 192: 887-892.
- 190 4. Gervaise. A, Esparbe-Vigneau. E, Pernin. M, Naulet. T, Porton. Y, Lampierre- Combe. M:
- 191 Evaluation of knowledge prescribing CT examinations on the radiation protection of patients.
- 192 Elsevier Masson. *Radiology Journal* (2011), 92, 671-678.
- 193 5. Einstein AJ, Henzlova MJ, Rajagopalan S. Estimated risk of cancer with radiation
- 194 exposure from 64-slice computed tomography coronary angiography. *JAMA* 2007; 298:
- 195 317-323.
- 196 6. Shiralkar S, Rennie A, Snow M, et al. Doctors' knowledge of radiation exposure:
- 197 questionnaire study. *BMJ* 2003; 327: 371-372.
- 198 7. Health Physics Society. ALARA. [http://www.hps.org](http://www.hps.org/public) /public
- 199 information/radterms/radfact1.html (accessed Mar 2009).
- 200 8. Cember H. Introduction to health physics. 3rd edition. McGraw-Hill, 1996.
- 201 9. Semghouli S, Amaoui. B, Maamri: Estimated radiation exposure from medical imaging for
- 202 patients of radiology service of Al Faraby Hospital, Oujda Morocco. *International Journal of*
- 203 *Cancer Therapy and Oncology*.2015; 3(3):3325
- 204 10. Semghouli, S., Amaoui, B., El Kharras, A., Shaim, A., Choukri, A., Hakam, 2017. Physicians
- 205 knowledge of radiation risk in prescribing CT imaging in Moroccan hospitals. *Br.J. Appl. Sci.*
- 206 *Technol.* 20(3),1
- 207 11. Council Directive 97/43/EURATOM/ of 30 June 1997 on Health protection of individuals
- 208 against the dangers of ionizing radiation in relation to medical exposure and repealing
- 209 Directive 84/466/Euratom. (OJ L-180 of 9 July 1997).
- 210 12. Semghouli. S, Amaoui. B, El Fahssi. M, Choukri A., Hakam O. K: Practitioners' Knowledge
- 211 on Patients' Radioprotection in Emergency and Radiology Services of Hassan II Hospital
- 212 Agadir Morocco. *International Journal of Modern Physics and Applications* ISSN: 2381-6945
- 213 Vol. 1, No. 5, (2015), pp. 205-209.
- 214 13. Lee CI, Haims AH, Monico EP, Brink JA, Forman HP: Diagnostic CT scans: assessment of
- 215 patient, physician, and radiologist awareness of radiation dose and possible risks.
- 216 *Radiology*. 2004 May;231(2):393-8.

- 217 14. Merzenich., H, Krille. L, Hammer. G, Kaiser. M, Yamashita. S, Zeeb. H: Paediatric CT scan
218 usage and referrals of children to computed tomography in Germany-a cross-sectional survey
219 of medical practice and awareness of radiation related health risks among physicians. BMC
220 Health Services Research 2012, 12:47.
- 221 15. Board on Radiation Effects Research Division on Earth and Life Studies, Committee to Assess
222 Health Risks from Exposure to Low Levels of Ionizing Radiation. Health Risks From
223 Exposure To Low Levels Of Ionizing Radiation Beir Vii Phase 2.National Research
224 Council Of The National Academies The National Academies Press Washington, D.C. 2006.
- 225 16. BrennerDJ , Doll R, Goodhead DT, salle EJ, Terre CE, Little JB, Lubin JH, Preston DL, Prest
226 on RJ, Puskin J, Ron E, Sachs RK,Samet JM, Setlow RB, Zaider M. Cancer risks attributable
227 to low doses of ionizing radiation: assessing what we really know. Proc Natl Acad Sci
228 USA. November 2003 25; 100 (24): 13761-6.
- 229 17. Jacob K, Vivian G, Stell JR. X ray dose training: are we exposed to enough? Clin
230 Radiol.2004;59:928–934.
- 231 18. Rice H E, Frush D. P, Harker M J, Farmer D, Waldhausen J H , Peer assessment of pediatric
232 surgeons for potential risks of radiation exposure from computed tomography scans. Journal
233 of Paediatric Surgeons Volume 42, Issue 7, Page 1157-1164.
- 234 19. Gerben B Keijzers and Charles J Britton: Doctors’ knowledge of patient radiation exposure
235 from diagnostic imaging requested in the emergency department. MJA• Volume 183 N
236
237