

1

2

3 **The Comparative Daily Activity Patterns of Dog**

4 **Faced Baboon (*Papio anubis*) in Captivity: A**

5 **Case Study of the University Zoo and Kano**

6 **Zoological Garden**

7

8

9

10

11

ABSTRACT

Aim: This study compares the daily activity patterns of dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Wudil and Kano Zoological Garden.

Materials and methods: This study of the activities of dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Wudil Zoo and Kano Zoological Garden was carried out from 6:00am to 6:00pm between December 2016 to January 2017. The observation in the activity patterns of dog face baboon (*Papio anubis*) were recorded in the recording sheet, observation ~~is~~ was done three times a week at 20 minutes interval in each of the cages under study.

Results: The findings on activity pattern of dog faced baboon (*Papio anubis*) in captivity shows that the day time activities decrease from morning to evening. 47.5% of the activities which include resting, movement and feeding were carried out in the morning, followed by afternoon and evening with 33.3% and 19.1% activities. The result of the activities of dog faced baboon in Kano zoological garden, indicated that 42.7% of the activities performed by dog faced baboon in captivity ~~is~~ are resting, this is followed by movement which accounted for 34.9% of the activities, while feeding activities account for the least with 22.4%. It shows that about 43.2% of the activities carried out by dog faced baboon in Kano University of Science and Technology Wudil, Zoo is Resting, followed by the Movement with 34.8% of the activities and the feeding activity account for 22%.

Conclusion: Due to the fact that majority of the baboons activities usually take place between Morning and afternoon, it is recommended that visitors should pay visitation to the

Zoo pen during that time. Feeding and chasing of animals by the visitors should be discouraged.

12

13 *Keywords: Papio Anubis, feeding, movement, resting, Kano University of Science and*
14 *Technology Wudil and Kano Zoological Garden*

15

16

17

18

19 **1. INTRODUCTION**

20

21 Activity patterns have been studied in several primate taxa including hominoids [1,2]
22 cercopithecines [3, 4, 5] colobines, [6]. Time is limited for most animals [7, 8]. Thus, animals
23 are faced with the challenge of allocating the limited time to different activities. According to
24 the optimality theory, “the amount of time that an organism spends engaged in various
25 activities depends on the cost of the activity relative to the derived benefits in that organism's
26 habitat” [9].

27 The amount of time spent on foraging activities therefore relates to the energy content of the
28 food relative to the costs of obtaining the food plus the cost of all other activities (resting,
29 moving or socialising). Thus, specifically, food availability and energy content are critical
30 determinants of an animals' daily activity pattern. Therefore, factors that influence the
31 availability of food have a strong bearing on time allocation profiles in baboons.

32 Due to the different costs and benefits of specific activities animals have varying time
33 allocation profiles based on age and sex for certain activities [10]. Furthermore, since these
34 activities cannot be performed simultaneously some individuals may allocate time between
35 various behaviours better than others [7, 8]. The costs and benefits of these activities
36 change with changes in the ecological and social state of the environment as well as the
37 physiological state of the animal. This gives rise to temporal and spatial variation in
38 individual activity budgets of the animal. Baboons allocate the greater proportion of their time
39 to foraging activities [11, 12, 13, 14, 15]. De Hoop and Mkuzi baboon troops spent 69.8 %
40 and 66.5 % of their time foraging respectively [14]. In a study of Alto, Hook and Lodge
41 baboon groups in [12] report them to spend 69.8 %, 75.2 % and 43 % of their time foraging,
42 respectively. The Lodge troop spent relatively less time foraging than Alto and Hook groups.

43 Weather patterns have both direct and indirect influences on the activity pattern of primates.
44 Rainfall and temperature have pervasive effects on animals [16] and so influence time
45 allocation patterns both temporally and spatially.
46 This study seeks to identify different types of activities carryout by dog faced baboon (*Papio*
47 *anubis*) in captivity.

48

49 **2. MATERIAL AND METHODS**

50

51 **2.1 Study area**

52 The study was carried out in Kano University of Science and Technology Zoo and Kano
53 Zoological Garden. **Wudil** has a total area of 362km² is located within Sudan savannah
54 region of Nigeria. The experimental site is located between the latitude 11^o 37'N and
55 longitude 8^o 58'E at an altitude of 403m above the sea level. The annual maximum rainfall is
56 850mm-870mm with a minimum and maximum temperature of 26^oc - 30^oc. The relative
57 humidity of the region is always low and ranges between 40% - 51%.-

58 **2.2 MATERIALS**

59 Field notebooks, stop clock, Recording sheet, Biro and Digital camera

60 **2.3 DATA COLLECTION**

61 Sampling method was used to study the activities of dog faced baboon (*Papio anubis*) in
62 Kano University of Science and Technology Wudil Zoo and Kano Zoological Garden from
63 6:00am to 6:00pm between December to January 2016. The observation in the activity
64 patterns of dog face baboon (*Papio anubis*) are recorded in the recording sheet, observation
65 is done three times a week at 20 minutes interval in each of the cages under study. Note:
66 this research is limited to period when the temperature is extremely low (Hammattan period).
67 The activity parameters recorded include: Feeding, Moving, and Resting and are described
68 as follows:

69 **Feeding:** the feeding began when the animal first made contact with any part of food or
70 other food substances, feeding bout terminated when the either moved more than one full
71 stride, even if it was carrying some food material on its hand and mouth or stopped looking
72 at the food material, by this definition, a switch to a new food type in the absence of either of
73 these condition was not for bout to be consider terminated, thus a single feed bout could
74 include more than one food type [17, 9].

75 **Resting:** this includes behavior during which an animal was neither feeding, moving or
76 engaged in other social behavior that include sleeping auto-grooming, looking around etc [9,
77 10].

78 **Moving:** this includes all locomotion activities like walking, running, climbing, jumping and
79 leaping but excluding short movements during feeding and locomotion during social behavior
80 e.g when primates chased one another [9, 10].

81 **Other activities:** other social behavior including all other activities which an animal's
82 attention and behavior where clearly directed toward another individual. These include allo-
83 grooming, mounting, mating, chasing, playing, aggressive or agnostic behaviours [9, 10].

84 All the activities are carried out in the **morning, afternoon and evening.**

85 **2.4 DATA ANALYSIS**

86 The data collected are subjected to descriptive statistics which includes frequency
87 distribution and percentage. The analysis of variance will be use to study the degree of
88 variation among the activities and also between two different animals.

91 **3. RESULTS AND DISCUSSION**

92
93 The result of the day time activities of dog faced baboon (*Papio anubis*) in captivity is
94 presented in Table 1. The result shows that the day time activities decrease from morning to
95 evening. 47.5% of the activities which include resting, movement and feeding were carried
96 out in the morning, followed by afternoon and evening with 33.3% and 19.1% activities.

97 **Table 1 Variation in day time activities of dog faced baboon (*Papio anubis*) in**
98 **captivity.**

100 DAY TIME	FREQUENCY (ACTIVITIES)	PERCENTAGE
101 Morning	67	47.5
102 Afternoon	47	33.3
103 Evening	27	19.1
104 Total	141	100

105
106 The result of the activities of dog faced baboon in Kano zoological garden showed in Tables
107 2 and 3, indicated that 42.7% of the activities performed by dog faced baboon in captivity ~~are~~
108 is resting, this is followed by movement which accounted for 34.9% of the activities, while
109 feeding activities account for the least with 22.4%. The result of this study is in variance with
110 the finding of [9] who reported 50.00% for feeding and 8.50% for resting for the kwano forest
111 baboons. In his study kwano forest baboon spent relatively higher proportion of time feeding
112 and lesser proportion of time resting and movement, this is probably due to the level of
113 availability and distribution of food resources at the site compared to captive environment.

114

115 **Table 2: Variation in the activities of dog faced baboon (*Papio anubis*) in Kano**
 116 **Zoological garden/day.**

117

118	NUMBER OF DAYS	FREQUENCY (ACTIVITIES)	PERCENTAGE (%)
119			
120	DAY 1		
121	Feeding	32	22.7
122	Moving	50	35.5
123	Resting	59	41.8
124	DAY 2		
125	Feeding	32	23.0
126	Moving	49	35.2
127	Resting	58	41.7
128	DAY 3		
129	Feeding	30	21.4
130	Moving	52	37.1
131	Resting	58	41.4
132	DAY 4		
133	Feeding	31	22.5
134	Moving	46	33.3
135	Resting	61	44.2
136	DAY 5		
137	Feeding	30	21.3
138	Moving	53	37.6
139	Resting	58	41.1
140	DAY 6		
141	Feeding	31	22.5
142	Moving	47	34.0
143	Resting	60	43.5
144	DAY 7		
145	Feeding	32	23.3
146	Moving	43	31.4
147	Resting	62	45.2
148	Total	974	100

149

151 **Table 3: variation in the activities of dog faced baboon (*Papio anubis*) in Kano**
 152 **Zoological garden/week**

153	ACTIVITIES	FREQUENCY (ACTIVITIES)	PERCENTAGE
154	Feeding	218	22.4
155	Movement	340	34.9
156	Resting	416	42.7
157	Total	974	100

158

159

The result of the dog faced baboon activities is indicated in Table 4 and 5, # shows that about 43.2% of the activities carried out by dog faced baboon in Kano University of Science

160 and Technology Wudil, Zoo is Resting, followed by ~~the~~ Movement with 34.8% of the
 161 activities and the feeding activity accounts for 22%. The low frequency of feeding and
 162 movement in the first and second days may be as a result that the animal ~~was~~
 163 menstruating on the first and second days of my research which leads to the resultant of
 164 high resting activity in the period. ~~However,~~ This result of the dog faced baboon activities in
 165 captivity indicated in the table above ~~is however~~, agrees with the finding of who reported [1]
 166 highest resting period than feeding and movement period

167 The Dog faced Baboon activities are significantly related to day time period. The baboon
 168 were observed to be very active in the morning followed by afternoon and evening. These
 169 activities which is made up of mostly movement and feeding may be due to the presence of
 170 visitors in the morning and afternoon. This activity pattern ~~during~~ morning, ~~a~~Afternoon and
 171 evening ~~has~~ve been commonly reported among arboreal species [17]. The daily activities of
 172 dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Zoo and
 173 Kano Zoological Gerden ranged between 6:00am in the morning to 6:00pm in the evening in
 174 which most of the visitors usually pay their visit. However, [17] was of the opinion that
 175 adaptive significances of diurnal variability in primate's activities budget are poorly
 176 understood. With regards to individual activities, resting, which includes sleeping, looking
 177 about etc. was the most frequent activity carried out by the dog faced baboon in captivity.
 178 This may be due to the confinement in which the baboons were kept. Most of the baboon's
 179 time was spent in sitting postures, standing or playing posture. In this position, the hind limb
 180 may be placed in variety of positions and the fore limbs of the baboon often at rest on the
 181 knees or between hind limb. However, despite the confinement, movement also constitute
 182 the substantial percentage of the dog faced baboon activities in Kano University of Science
 183 and Technology and Kano Zoological Garden. The movement which accounted for 34.9% in
 184 Kano Zoological Garden and 34.8 in the University Zoo include walking, running, climbing,
 185 leaping and riding. This significance percentage might be due to the fact that dog faced
 186 baboon are usually regarded as one of the most entertaining animal within the Zoo.
 187 Meanwhile, of both the animals studied, the Baboon in the Kano Zoological Garden has the
 188 slightly higher frequency of activities. This might ~~be~~ also ~~be~~ due to its ability to communicate
 189 or play with more number of visitors.

190

191 **Table 4: Variation in the activities of dog faced baboon in Kano University of Science**
 192 **and Technology Wudil, Zoo/day.**

193	NUMBER OF DAYS	FREQUENCY (ACTIVITIES)	PERCENTAGE
194			
195	DAY 1		

196	Feeding	29	21.0
197	Moving	37	26.8
198	Resting	72	52.1
199	DAY 2		
200	Feeding	27	19.8
201	Moving	35	25.7
202	Resting	74	54.4
203	DAY 3		
204	Feeding	30	21.9
205	Moving	42	30.6
206	Resting	65	47.4
207	DAY 4		
208	Feeding	30	22.2
209	Moving	45	33.3
210	Resting	60	44.4
211	DAY 5		
212	Feeding	31	22.3
213	Moving	56	40.3
214	Resting	52	37.4
215	DAY 6		
216	Feeding	32	23.3
217	Moving	60	43.8
218	Resting	45	32.8
219	DAY 7		
220	Feeding	32	23.7
221	Moving	58	43.0
222	Resting	45	33.3
223	Total	957	100

224

225

226 **Table 5: Variation in the activities of dog faced baboon (Papio anubis) in Kano**

227 **University of Science and Technology Zoo/week**

228	ACTIVITIES	FREQUENCY (ACTIVITIES)	PERCENTAGE
229	Feeding	211	22.0
230	Movement	333	34.8
231	Resting	413	43.2
232	Total	957	100

233

234 **CONCLUSION**

235 This study was designed to gather information on the daily activity pattern of Dog faced

236 Baboon in Kano University of Science and Technology and Kano Zoological Garden. From

237 the study, the following conclusion can be made. The dog faced baboons are most active in

238 the morning. ~~Also-Furthermore, r~~Resting constitute the most frequent activity of dog faced

239 baboon in captivity. Most of the baboon activities have short duration.

240

241

242 **COMPETING INTERESTS**

243 Authors have declared that no competing interests exist.

244

245

246 **REFERENCES**

247 1. Goodall, J. The chimpanzees of Gombe: patterns of behaviour, Belknap Press,
248 Cambridge, 1986.

249 2. Clutton-Brock, T.H.C. and Harvey P. Species differences in feeding and ranging
250 behavior in primates. In: primate Ecology (Ed. T.H.C. Clutton-Brock), New York:
251 Academic press, 1977. pp. 557-584

252 3. Post, D.G. Activity Patterns of Yellow Baboons (*Papio cynocephalus*) In the
253 Amboseli National Park, Kenya. *Animal Behaviour* 1981, 29: 357-374.

254 4. Van Schaik C.P. Why are diurnal primates living in groups? *Behaviour* 1983,
255 87:120-144.

256 5. Isbell, L.A., Preutz J.D. and Young T.P. Movements of vervet (*Cercopithecus*
257 *aethiops*) and patas monkeys (*Erythrcebus patas*) as estimators of food resource
258 size, density and distribution. *Behaviour Ecology Sociobiology* 1993, 42:123-133.

259 6. Strushaker, T.T. *The red colobus monkey*, University of Chicago Press, Chicago,
260 1975.

261 7. Dunbar, R.I.M. *Primate Social Systems*, Cornell University Press, Ithaca, 1988.

262 8. Dunbar, R. I. M. Time: A Hidden Constraint on the Behavioural Ecology of
263 Baboons, *Behaviour Ecology Sociobiology*, 1992;31: 35-49.

264 9. Gessaman, J.A. Ecological energetics of homeotherms: a view compatible with
265 ecological modelling, Utah University Press *Monograph series*, 1973 20: 1-155.

266 10. Melle, E.M. socio-ecology of Olive baboons thesis Department of wildlife and
267 fisheries management, university of Ibadan, 2004.

268 11. Altmann, J and Altmann, S *Baboon Ecology: African Field Research*. Karger, Basel,
269 1970.

270 12. Amboseli, Kenya, Bronikowski and Altmann Mate Guarding Constrains Foraging Activity of
271 Male Baboons, *Animal Behaviour*, 1996; 51: 1269–1277.

272 13. Barton, R.A, Whitten A, Strum S.C, Byrne, R. W and Simpson, A.J Habitat and
273 Resource Availability in Baboons, *Animal Behaviour*, 1992; 43: 831-844.

- 274 14. Gaynor, D Foraging and feeding behaviour of chacma baboons in a woodland
275 habitat, PhD thesis, University of Natal, 1994.
- 276 15. Johnson, S.E. & Bock, J. (tradeoffs in skill acquisition and time allocation among
277 juvenile chacma baboons, *Human Nature* Volume, 2004; 15 (1): 45-62.
- 278 16. Bronikowski, A.M and Altmann, J Foraging in variable environment: weather
279 patterns and the behavioural ecology of baboons, *Behaviour Ecology Sociobiology*,
280 1996; 39: 11-25
- 281 17. Post, D.G Feeding Behaviour of Yellow Baboons (*Papio Cynocephalus*) In
282 Amboseli National Park, Kenya, *International Journal of Primatology*, 1982; 3(4): 403-
283 430.
- 284 18. Clutton-Brock, T.H.C. Some aspect of intra-specific variation in feeding and ranging
285 behavior in primates. In: primate Ecology (Ed. By T.H.C. Clutton-Brock), pp. 529-
286 536. New York: Academic press, 1977.
- 287

UNDER PEER REVIEW