

**Floristic analysis and phytosociology in an area of Caatinga, **country****

**name**

**ABSTRACT-** Studies on the composition and structure of vegetation can provide important information for decision-making and the application of forest-management techniques. The aim of this research was to analyse an area of Caatinga using vegetation characterisation and forest inventory in the Rocha Eterna Community, in the district of São João do Piauí, **country name**. Simple Random Sampling was used, installing 17 sampling units of 20 x 20 m. The phytosociological parameters of the horizontal and vertical structure, the floristic diversity of the species and the timber production in the area were evaluated. Fabaceae was the most representative family. The three most representative species in the area were *Senna acuruensis* (Benth.) H.S. Irwin & Barneby, *Sideroxylon obtusifolium* (Roem. & Schult.) T.D.Penn. and *Mimosa tenuiflora* (Willd.) Poir. The Shannon-Weaver diversity index (H') for the area was 1.74 nats.ind<sup>-1</sup>. The estimated basal area for the area was 8.68 m<sup>2</sup>.ha<sup>-1</sup>. The estimated actual volume was 36.56 m<sup>3</sup>.ha<sup>-1</sup>. Species diversity in the fragment under study was considered low.

**Keywords:** Floristic diversity. Vegetation structure. Survey of the Caatinga.

**1. INTRODUCTION**

Caatinga vegetation covers most of the area of semi-arid climate in the Northeast of Brazil (Giulietti et al., 2004). The vegetation is composed of herbaceous and woody

25 species, including cacti and bromeliads, with an emphasis on the xerophytic and  
26 deciduous aspect, a typical characteristic of the species, which display great  
27 heterogeneity in relation to their phyto-physiognomies and horizontal and vertical  
28 structures (Prado, 2003).

29 The degradation of the Caatinga through anthropogenic action, deforestation and  
30 the burning of forested areas for agricultural activities is increasingly compromising the  
31 natural resources and sustainability of the biome. Further, there is little understanding of  
32 the biome, especially of how to use the resources it offers, as there are particular aspects  
33 of the Caatinga that must be considered if it is to regenerate and again be exploited  
34 (Tabarelli et al., 2000; Vasconcelos et al., 2017).

35 Research into floristic and phytosociological composition in forests of the  
36 Caatinga Biome is an important tool for detailing plant species diversity and verifying  
37 how they are distributed in a given environment using structural analysis, which allows  
38 an estimation of the distribution of individuals by species and consequently by family  
39 (Souza, 2009).

40 Phytosociological studies also aid in forest management planning, environmental  
41 impact studies and forest restoration and reclamation projects in degraded areas, and  
42 may also indicate the potential use for a given species through information on richness  
43 and abundance, volumetric potential and diameter-class distribution that will influence  
44 its use, whether for wood, charcoal, cuttings or posts, among others (Santos et al.,  
45 2017a).

46 Studies on the composition and structure of vegetation can provide basic  
47 information for decision-making in the application of forest-management techniques  
48 (conservation and maintenance), so that any intervention in the forest should be planned  
49 and preceded by a detailed inventory, which would provide information such as

50 estimates of the floristic composition, and of the horizontal, vertical and parametric  
51 structures (Souza, 2003).

52 It should be noted that forest inventories are still scarce in the southern region of  
53 the State of Piauí, so that research which evaluates the conditions under which the  
54 natural vegetation is found is extremely important. This study therefore aimed to  
55 analyse an area of caatinga vegetation using vegetation characterisation and forest  
56 inventory in the Rocha Eterna Community, in the district of São João do Piauí.

57

## 58 **2. MATERIAL AND METHOD**

### 59 **2.1 Study Area**

60 The **research** work was carried out with the support of the Non-Governmental  
61 Organisation SOS Sertão in a fragment of Caatinga of approximately 200 ha in the  
62 Rocha Eterna Community of the district of São João do Piauí, in the microregion of the  
63 Upper Middle Canindé. The area is located at 08°21'29 S and 42°14'48" W, and is about  
64 482 km from the state capital, Teresina.

65 The city of São João do Piauí has an average altitude of 222 m above sea level,  
66 with minimum temperatures of 22°C and maximum temperatures of 39°C, and a climate  
67 that is considered semi-arid, hot and dry. The average annual rainfall is 500 mm, with  
68 two periods a year: a rainy season lasting from 3 to 4 months and a dry season during  
69 the remaining months (Aguiar; Gomes, 2004).

70 The soils of the region are shallow, considered young, sometimes stony, still  
71 influenced by the underlying material, with a predominant vegetation of tree and shrub  
72 Caatinga, where the soils are of sedimentary origin and extremely sandy (Aguiar;  
73 Gomes, 2004; Nascimento et al., 2008).

74 The terrain in the region shows a flattened surface with areas of depression in  
 75 which temporary lagoons form, surfaces of low plates, flat relief with smoothly  
 76 undulating areas and altitudes varying between 150 and 300 metres, undulating surfaces  
 77 and moving relief, corresponding to slopes with the residual extensions of plates,  
 78 gradients and steep slopes, with valleys, elevations and tabular summit surfaces of flat  
 79 relief (Aguiar; Gomes, 2004).

## 80 **2.2 Data collection and processing**

81 For the survey, an area of approximately 200 ha of native forest was selected,  
 82 where 17 plots of 20 x 20 m (400 m<sup>2</sup>) were randomly allocated, resulting in an  
 83 inventory area of 6,800 m<sup>2</sup>.

84 Inside the 17 plots, measurements were taken of the tree and shrub forest species  
 85 having a circumference at breast height (CBH) equal to or greater than 10 cm, where  
 86 they were identified *in loco* by means of their dendrological aspects; when possible,  
 87 botanical material was collected for later identification and/or confirmation of the  
 88 scientific names by consulting the literature. The APG II system (2003) was used to  
 89 classify the plants up to species level.

90 After acquiring the data, the phytosociological parameters of the horizontal and  
 91 vertical structure of the sampled species were evaluated (Felfili; Rezende, 2003)  
 92 together with the floristic diversity, using the Shannon-Weaver Index (H'), the Simpson  
 93 Dominance Index (C) and the Pielou Evenness Index (J'), as per Table 1.

94

95 **Table 1.** Shannon-Weaver diversity index (H'), Simpson dominance index (C) and  
 96 Pielou evenness index (J').

Index	Designation	Formula
Diversity	Shannon-Weaver (H')	$H' = \sum_{i=1}^s \left( \frac{n_j}{N} \cdot \ln \frac{N}{n_j} \right)$

Dominance	Simpson (C')	$c' = \sum_{i=1}^s \frac{(n_j \cdot (n_j - 1))}{N \cdot (N - 1)}$
Evenness	Pielou (J')	$J' = \frac{H'}{H'_{max}}$

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

Where: S = total number of sampled species; N = total number of sampled individuals; ni = number of sampled individuals of the ith species; Ln = neperian logarithm; Hmax = Ln (S) = total number of sampled species.

To calculate the volume, the DBH at 1.30 m including bark of adult trees measured in the sampling units was considered, applying the following formulae:

$$VCc/c = (\pi * (DBH^2) * Ht)/40000$$

(Equation 1)

where: VCc/c = cylindrical volume of the tree with bark;  $\pi$  = "pi" (3.1416...); DBH<sup>2</sup> = diameter at breast height, squared (in centimetres); Ht = total height of the tree (in metres); 40.000 = quadratic conversion factor from centimetres to metres (from the DBH);

$$VA = VCC * ff$$

(Equation 2)

where: VA = actual volume (in cubic meters - m<sup>3</sup>); ff = form factor (0.9 dimensionless) (ZAKIA *et al.*, 1988).

To estimate the stacked volumes, the following formula was applied:

$$VS = VR * fe$$

(Equation

3)

where: VS = stacked volume (in steres); VA = actual volume (cubic metres); fe = stacking factor (3.41 adimensional) (ZAKIA *et al.*, 1988).

118 To describe the community structure, the following phytosociological parameters  
119 were estimated: absolute density (DA), relative density (DR), absolute frequency (FA),  
120 relative frequency (FR), absolute dominance (DoA), relative dominance (DoR),  
121 importance value index (IVI) and coverage value index (CVI).

122 Processing or computation of the field data was carried out using an Excel 2010  
123 spreadsheet and the MATA NATIVA v3.01 software, used in floristic and  
124 phytosociological analysis.

125

### 126 3. RESULTS AND DISCUSSION

127

128 In the survey, 792 individuals were sampled, belonging to 6 families, 17 species  
129 and 15 genera. Of these species, two were identified by their common name only.  
130 Family Fabaceae was the most representative in number of species, together with  
131 Euphorbiaceae; only one species was registered for each of the remaining families, as  
132 shown in Table 2.

133

134 **Table 2.** List of identified species found in the sampling plots in the Rocha Eterna  
135 Community, **country name.**

Family/Species	Common Name	Habit
<b>Annonaceae</b>		
<i>Annona leptopetala</i> (R.E.Fr.) H.Rainer	Bananinha	Arboreal
<b>Bignoniaceae</b>		
<i>Jacaranda jasminoides</i> (Thunb.) Sandwith	Chifre de carneiro	Arboreal
<b>Combretaceae</b>		
<i>Terminalia fagifolia</i> Mart.	Chapada	Arboreal
<b>Euphorbiaceae</b>		
<i>Croton blanchetianus</i> Baill.	Marmeleiro	Arboreal

<i>Manihot glaziovii</i> Müll.Arg.	Maniçoba	Arboreal
<b>Fabaceae</b>		
<i>Machaerium acutifolium</i> Vogel	Violete	Arboreal
<i>Poincianella pyramidalis</i> (Tul.) L.P.Queiroz	Catingueira	Arboreal
<i>Bauhinia cheilantha</i> (Bong.) Steud.	Mororó	Arboreal
<i>Pityrocarpa moniliformis</i> (Benth.) Luckow & R.W.Jobson	Catandúva	Arboreal
<i>Senna acuruensis</i> (Benth.) H.S.Irwin & Barneby	Canela de Velho	Arboreal
<i>Machaerium</i> Pers.	Jacarandá de Sangue	Arboreal
<i>Mimosa tenuiflora</i> (Willd.) Poir.	Jurema preta	Arboreal
<i>Piptadenia stipulacea</i> (Benth.) Ducke.	Jurema Branca	Arboreal
<i>Dimorphandra mollis</i> Benth	Folha Miúda	Arboreal
<b>Sapotaceae</b>		
<i>Sideroxylon obtusifolium</i> (Roem. & Schult.) T.D.Penn.	Espinheiro	Arboreal
<b>Undetermined</b>		
Undetermined 1	Birro	Arboreal
Undetermined 2	Violete Preto	Arboreal

136

137 It should be pointed out that in other surveys already carried out in the Caatinga  
138 region, the families Fabaceae and Euphorbiaceae are commonly recorded as having the  
139 largest number of species. (Calixto Júnior; Drumond, 2014; Dias et al., 2014; Leite et  
140 al., 2015; Santos et al., 2017b).

141 Family Fabaceae presented 9 species, these with around 8 genera; the genus  
142 *Machaerium* was repeated and represented by two species; the genus has around 140  
143 species, among which, *Jacaranda* stands out with numerous variations.

144 It is common in floristic and phytosociological surveys for the most representative  
145 species to be Fabaceae and Euphorbiaceae, considered as one of the most numerous  
146 families among plants groups.

147 Vasconcelos et al. (2017) showed similar results for the number of species of  
148 family Euphorbiaceae in studies carried out in the district of São Francisco, Piauí;  
149 however, being an ecotone of the Cerrado and Caatinga biomes, this was an area with a  
150 higher floristic diversity index as well as having a greater number of species and  
151 individuals per unit area (absolute density).

152 In a survey carried out in an area of Caatinga in the State of Paraíba, Oliveira et al.  
153 (2009) found that the families Fabaceae and Euphorbiaceae excelled in the number of  
154 species and also had a large total number of individuals. Rodal (2002), found that the  
155 families Fabaceae and Euphorbiaceae were the most representative in phytosociological  
156 studies of the woody component in an area of Caatinga in the district of São Raimundo  
157 Nonato, Piauí.

158 The aim of this study was to register species of the tree and shrub stratum,  
159 however, the only component found was arboreal, although some shrub species were  
160 seen outside the plots. What demonstrates the development of species in the area is their  
161 being better established in the forest; management should be seen as a positive way of  
162 maintaining biodiversity, compared to intense extraction of the vegetation.

163 The most abundant species in the inventoried region were *Senna acuruensis* (339),  
164 *Sideroxylon obtusifolium* (178) and *Mimosa tenuiflora* (Willd.) Poir. (130) (Table 3).

**Table 3.** List of inventoried species, in the Rocha Eterna Community, in the district of São João do Piauí. The values are in descending order of IVI. Where N = number of individuals, FA = Absolute Frequency (%), FR = Relative Frequency (%), DA = Absolute Density (ind.ha<sup>-1</sup>), DR = Relative Density (%), DoA= Absolute Dominance (m<sup>2</sup>.ha<sup>-1</sup>), DoR= Relative Dominance (%), IVI = Importance Value Index (%), CVI

---



= Coverage Value Index (%).

Scientific Name	N	DA	DR(%)	FA	FR(%)	DoA	DoR	CVI (%)	IVI (%)
<i>Senna acuruensis</i> (Benth.) H.S.Irwin & Barneby	339	498.529	42.8	100	14.78	10.228	80.08	61.44	45.89
<i>Sideroxylon obtusifolium</i> (Roem. & Schult.) T.D.Penn.	178	261.765	22.47	88.24	13.04	0.562	4.40	13.44	13.30
<i>Mimosa tenuiflora</i> (Willd.) Poir.	130	191.176	16.41	94.12	13.91	0.425	3.32	9.87	11.22
<i>Pityrocarpa moniliformis</i> (Benth.)	31	45.288	3.92	94.12	13.90	0.163	1.28	2.60	6.39
<i>Bauhinia cheilantha</i> (Bong.) Steud.	15	22.059	1.89	52.94	7.83	0.029	0.22	1.06	3.31
<i>Croton blanchetianus</i> Baill.	31	45.588	3.91	35.29	5.22	0.065	0.51	2.21	3.21
<i>Terminalia fagifolia</i> Mart.	7	10.294	0.88	23.53	3.48	0.574	4.5	2.69	2.95
<i>Manihot glaziovii</i> Müll.Arg.	13	19.118	1.64	35.29	5.22	0.105	0.82	1.23	2.56
Undetermined 2	11	16.176	1.39	35.29	5.22	0.104	0.82	1.10	2.47
<i>Dimorphandra mollis</i> Benth.	9	13.235	1.14	29.41	4.35	0.062	0.48	0.81	1.99
<i>Jacaranda jasminoides</i> (Thunb.) Sandwith	4	5.882	0.51	23.53	3.48	0.087	0.68	0.59	1.55
Undetermined 1	7	10.294	0.88	17.65	2.61	0.149	1.17	1.03	1.55
<i>Machaerium acutifolium</i> Vogel	6	8.824	0.76	11.76	1.74	0.093	0.73	0.74	1.07
<i>Piptadenia stipulacea</i> (Benth.) Ducke	3	4.412	0.38	17.65	2.61	0.005	0.04	0.21	1.01
<i>Annona leptopetala</i> (R.E.Fr.) H.Rainer	5	7.353	0.63	5.88	0.87	0.025	0.2	0.41	0.57
<i>Poincianella pyramidalis</i> (Tul.) L.P.Queiroz	1	1.471	0.13	5.88	0.87	0.06	0.47	0.30	0.49
<i>Machaerium</i> Pers.	2	2.941	0.25	5.88	0.87	0.038	0.3	0.27	0.47
<b>Total</b>	<b>792</b>	<b>1164.706</b>	<b>100</b>	<b>676.47</b>	<b>100</b>	<b>12.772</b>	<b>100</b>	<b>100</b>	<b>100</b>

165

166

167 For the importance value index (IVI), which estimates the ecological importance  
 168 of a given species in the community, and comprises the sum of each value of the  
 169 relative parameters, the species *Senna acuruensis* had the highest value, of 45.89%,  
 170 followed by *Sideroxylon obtusifolium* with 13.30%, *Mimosa tenuiflora* with 11.22% and  
 171 *Pityrocarpa moniliformis* with 6.39%.

172 The species *Senna acuruensis*, *Sideroxylon obtusifolium* and *Mimosa tenuiflora*  
 173 also presented the highest values for coverage in the area; the species are important, as  
 174 together they account for 84.75% of the individuals sampled in the area.

175 Due to species distribution, these values are related, with *Senna acuruensis* and  
 176 *Mimosa tenuiflora* present in all the inventoried plots, and *Sideroxylon obtusifolium*  
 177 occurring in nearly all the plots. The species *Croton blanchetianus* Baill., although

178 presenting considerable values for density, was not among the five species with the  
179 greatest IVI, since it did not stand out as to frequency. All the parameters can be seen in  
180 table 3.

181 The Shannon-Weaver diversity index ( $H'$ ) for the area was  $1.74 \text{ nats.ind}^{-1}$ , the  
182 Simpson Dominance index ( $C$ ) was 0.74 and the Pielou Evenness index ( $J'$ ) was 0.60.

183 The Simpson Dominance Index ( $C$ ) seen in this study also presented lower results  
184 than in other studies in the area of Caatinga, such as that of Sousa, et al. (2017) of 0.88  
185 in Bom Jesus, Piauí, and 0.93 seen by Vasconcelos et al. (2017) in São Francisco do  
186 Piauí. The Shannon index ( $H'$ ),  $1.74 \text{ nats.ind}^{-1}$ , was lower than those found in other  
187 studies in the Caatinga, such as Rodal et al. (1998), Rodal et al. (2006) and Ferras et al.  
188 (2006).

189 This value is considered low, a fact related to the climate conditions and  
190 associated with the types of soils predominant in these regions (Trovão, 2004). When  
191 evaluating the values of these two indices, it is correct to say that the study area has less  
192 diversity when compared to other areas located in the same region.

193 However, the study area presented greater diversity when compared to Caatinga  
194 vegetation in other regions; as for example, an area studied by Holanda et al. (2015) in  
195 Cajaseirinhas, Paraíba, where the vegetation structure in remnants of caatinga with a  
196 history of disturbance showed a value of  $1.5 \text{ nats.ind}^{-1}$ .

197 The value for the Pielou Evenness Index ( $J'$ ) found in this study ( $J = 0.60$ ) was  
198 similar to that found by Pereira Júnior et al. (2012) of  $J = 0.63$ . These values for species  
199 distribution are considered low when compared to other areas of Caatinga; in Monteiro,  
200 Paraíba, a greater value, corresponding to  $J = 0.73$  was found by Pegado et al. (2006),  
201 indicating the more even distribution of plant species throughout the area.

202 The vegetation had a mean height of 4 m for the 792 individuals inventoried,  
 203 reflecting well the characteristics of the vegetation for height and population density.  
 204 Distribution of the basal area and of the estimated actual volume by diameter class  
 205 shows that the highest values were found for a diameter class of between 6.5 and 11.5  
 206 cm, followed by the class between 1.5 and 6.5 cm (Table 4).  
 207

**Table 4.** Distribution of Basal Area (BA), Actual Volume (VA) and Stacked Volume (VS) for diameter class, on the Rocha Eterna property, **country name** .

Class	N	BA (m <sup>2</sup> .ha <sup>-1</sup> )	Act. Vol (m <sup>3</sup> .ha <sup>-1</sup> )	Stacked Vol (st.ha <sup>-1</sup> )
1.5   6.5	436	0.682	2.4405	11.92
6.5   11.5	169	0.951	3.5398	17.28
11.5   16.5	71	1.055	4.186	20.44
16.5   21.5	48	1.338	5.4146	26.44
21.5   26.5	27	1.214	5.0391	24.60
26.5   31.5	20	1.317	5.9434	29.02
31.5   36.5	16	1.424	6.5183	31.82
36.5   41.5	3	0.336	1.4607	7.13
41.5   46.5	1	0.139	1.0001	4.88
46.5   51.5	0	0	0	0.00
51.5   56.5	1	0.228	1.0259	5.01
<b>Total</b>	<b>792</b>	<b>8.684</b>	<b>36.568</b>	<b>178.54</b>

208  
 209 The three species of greatest abundance in the vegetation represent 84.75% of the  
 210 sampled population. It was found that the basal area in this region was very low, this is  
 211 related to the high importance value index found in the primary-stage species. It may  
 212 also be related to disturbances in the area (the history of land use), which may have  
 213 influenced vegetation dynamics. The sampling error was found to be 17.46%, which  
 214 was acceptable and gave satisfactory results.

215 Considering the sum of the basal area of the individuals per hectare, the value  
 216 found in this study was greater than that found by Santos et al. (2017b), (7.6 m<sup>2</sup>.ha<sup>-1</sup>),

217 who evaluated the woody component of a fragment of Caatinga located in another area  
218 in the district of Upanema, in the State of Rio Grande do Norte. Amorim et al. (2005)  
219 evaluated the structure of tree and shrub vegetation in an area of Caatinga in the district  
220 of Serra Negra do Norte, also in the state of Rio Grande do Norte, in the microregion of  
221 Western Seridó, where a lower value ( $6.1 \text{ m}^2.\text{ha}^{-1}$ ) was also found.

222 This difference can be explained by the fact that these areas are located in a  
223 different region of the Caatinga, in another State, and also because in both the above  
224 studies the basal area was calculated from data obtained from the woody stratum (trees  
225 + shrubs), whereas in the present study, only tree-like individuals were considered (see  
226 Table 2).

227 The Actual Volume in the area was greater than that reported by Santos et al.  
228 (2017b), who found a value of  $30.03 \text{ m}^3.\text{ha}^{-1}$  in the town of Desterro, in the district of  
229 Teixeira, Paraíba, and lower than those found by Leite et al. (2015), of  $40.40 \text{ m}^3.\text{ha}^{-1}$   
230 also in the district of Teixeira, Paraíba, in the microregion of Serra do Teixeira.  
231 However, this value was greater than the values found by Xavier et al. (2005) when  
232 assessing the timber potential of the microregions of Sousa and Itaporanga, and who  
233 calculated a volume of  $17 \text{ m}^3.\text{ha}^{-1}$  and  $20.51 \text{ m}^3. \text{ha}^{-1}$  respectively. It would seem that the  
234 habit of the components under study contributed to this significant difference between  
235 the areas, in addition to environmental factors, that have a different effect in each  
236 locality.

237 **4. CONCLUSION**

238 Family Fabaceae had the largest number of individuals. The diversity found in the  
239 area was considered low, also displaying little uniformity in the distribution of  
240 vegetation when compared to other studies in areas of Caatinga in the State of Piauí,  
241 highlighting the regional variations, especially in population density.

242 The place where the Caatinga vegetation is found has an effect on the diversity  
243 indices.

244 The inclusion of only tree species in the study increased the values for basal area  
245 and volume compared to other areas of Caatinga, where these values were calculated  
246 from the complete woody stratum (shrub and tree).

247 The species that stood out for IVI in the fragment under study were *Senna*  
248 *acuruensis*, *Sideroxylon obtusifolium*, *Mimosa tenuiflora*, *Pityrocarpa moniliformis* and  
249 *Bauhinia cheilantha*.

250 **REFERENCES- Can include more recent year references**

251 Aguiar RB, Gomes, JRC. Groundwater Source Registration Project, Piauí: Diagnosis of  
252 the Municipality of São João do Piauí-PI-CPRM-Fortaleza, 2004.

253 Amorim LL, Sampaio EVSB, Araújo EL (2005). Flora and structured shrub-tree  
254 vegetation of an area of Caatinga do Seridó, RN. Brazil. Brazilian Botanical Act, Porto  
255 Alegre, v. 19, no. 3, p. 615-623. <https://doi.org/10.1590/S0102-33062005000300023>.

256 Calixto Junior J.T, Drumond MA (2014). Comparative study of the phytosociological  
257 structure of two Caatinga fragments at different conservation levels. Brazilian Forest  
258 Research, Colombo, v. 34, no. 80, p. 345-355

259 Days PMS, Diodate MA, Grigio AM (2014). Phytosociological survey of forest  
260 remnants in the municipality of Mossoró-RN. *Caatinga Magazine*, Mossoró, v. 27, no.  
261 4, p. 183-190.

262 Felfili J.M, Rezende RP. *Concepts and Methods in Phytosociology*. Brasília: UnB,  
263 Department of Forest Engineering, 2003. 68 p. (Forest Technical Communications).

264 Ferraz JSF, Albuquerque UP, Meunier IMJ (2006). Use value and structure of woody  
265 vegetation on the banks of Riacho do Navio, Floresta, PE, Brazil. *Brazilian Botanical*  
266 *Act*, v.19, n.1, p.125-134. <http://dx.doi.org/10.1590/S0102-33062006000100012>.

267 Giulietti AM, Bocage Neta AL, Castro AAJ, Rojas CFLG, Sampaio EVSB, Virgínio J,  
268 Harley, RM. Diagnosis of native vegetation of the Caatinga biome. In: SILVA, J. D.,  
269 TABARELLI, M., FONSECA, M. D., & LINS, L. V. *Caatinga Biodiversity: Priority*  
270 *Conservation Areas and Actions*. Brasilia: Ministry of the Environment, 2004, p. 48-90.

271 Netherlands AC, Lima FTD, Silva BM, Dourado RG, Alves AR (2015). Vegetation  
272 structure in caatinga remnants with different disturbance histories in Cajazeirinhas (PB).  
273 *Caatinga Magazine*, Mossoró, v. 28, no. 4, p. 142 - 150. [http://dx.doi.org/10.1590/1983-](http://dx.doi.org/10.1590/1983-21252015v28n416rc)  
274 [21252015v28n416rc](http://dx.doi.org/10.1590/1983-21252015v28n416rc).

275 We read JR, Rodal MJN (2002). Phytosociology of the woody component of a stretch  
276 of Caatinga vegetation in the Serra da Capivara Piauí National Park, Brazil. *Minutes*  
277 *Bot Bras*. v. 16, no. P. 23-42. <http://dx.doi.org/10.1590/S0102-33062002000100005>.

278 Birth HTS, Light KS, Wolff LF, Birth MPSCB, Oliveira ME. *Semi-Arid Plants:*  
279 *Knowledge and Uses in Nesting Marrecas*. Teresina: Embrapa Mid-North, 2008.

280 Oliveira PTB, DMBM Thunder, Carvalho ECD, Souza BC, Ferreira LMR (2009).  
281 Floristic and phytosociology of four vegetation remnants in mountainous areas in  
282 Paraíba cariri. *Caatinga Magazine*, Mossoró, v. 22, no. 4, p.169-178.

283 Taken from MAC, Andrade LA, Felix LP, Israel MP (2006). Effect of biological  
284 invasion of mesquite - *Prosopis juliflora* (Sw.) DC. About the composition and structure  
285 of the caatinga shrubby tree stratum in Monteiro, PB, Brazil. *Brazilian Botanical Act*, v.  
286 20, no. 4. p.887-898. <http://dx.doi.org/10.1590/S0102-33062006000400013>.

287 Pereira Junior LR, Andrade AP, Araújo KD (2012). Floristic and phytosociological  
288 composition of a caatinga fragment in Monteiro, PB. *Holos*, v. 6, no. 2, p. 73-87.  
289 <https://doi.org/10.15628/holos.2012.1188>.

290 Rodal MJN, Andrade KVA, Sales MF, Gomes APS (1998). Phytosociology of the  
291 woody component of a vegetation refuge in the municipality of Buíque, Pernambuco.  
292 *Brazilian Journal of Biology*, v.58, n.3, p. 517-526.

293 Rodal MJN, Birth LM (2006). The arboreal component of a dry forest in northeastern  
294 Brazil. *Brazilian Journal of Biology*, v.66, no.2A, p.479-491.

295 Santos WS, Henriques IGN, Santos WS, Ramos GG, Vasconcelos GS, Vasconcelos  
296 ADM (2017b) Floristic-phytosociological analysis and logging potential in a caatinga  
297 area under forest management. *Scientific Farming in the Semiarid, Ducks*, v.13, n.3,  
298 p.213-211, 2017b.

299 Santos WS, Souza MP, Nobrega GFQ, Medeiros FS, Alves AR, Hokanda AC (2017a)  
300 Floristic-phytosociological characterization of the woody component In a caatinga  
301 fragment in the municipality of Upanema-RN. *Native, Sinop*, v.5, no.2, p.85-91.  
302 [10.5935 / 2318-7670.v05n02a02](https://doi.org/10.5935/2318-7670.v05n02a02).

303 Souza JS, Espírito Santo FDB, Fontes AL AL, Oliveira-Filho AT, Botezelli L (2003).  
304 Analysis of floristic and structural variations of the tree community of a semideciduous  
305 forest fragment on the banks of the Capivari river, Lavras - MG. *Tree Magazine*,  
306 Viçosa, v.27, n.2, p.185-206.

307 Souza, P. F. Analysis of the vegetation of a caatinga fragment in the jatobá reservoir  
308 microbasin. Patos-PB, 2009. 51 p. Course conclusion work (Forest Engineering) -  
309 Federal University of Campina Grande, Patos, PB, 2009.

310 DMBM thunder. Phytosociology and Ecophysiological Aspects of the Woody  
311 Component in Caatinga Fragments in the Bodocongó River Basin - Paraíba. 2004. 108f.  
312 Thesis (Doctorate in Natural Resources) - UFCG, Campina Grande.

313 Vasconcelos ADM, Henriques IGN, Souza MP, Santos WS, Santos WS, Ramos GG  
314 (2017). Floristic and phytosociological characterization in Caatinga area for forest  
315 management purposes in São Francisco-PI. Scientific Farming in the Semiarid, Ducks,  
316 v.13, n.4, p.329-337.

317 Xavier EPL, Paes JB, Lira Filho JA (2005). Wood potential of two microregions of  
318 Paraíba State. Biomass and Energy, v.2, n.2, p. 103-112.

319 Zakia MJB, Pareyn FGC, Riegelhaupt E. Weight and Volume Equations for Eight  
320 Native Woody Species of Seridó-RN. In: Forest Management Plan for the Seridó-RN  
321 Region. ch. 4, v. 1, Christmas: Project UNDP / FAO / IBAMA / BRA / 87/007, 1988.