

ANTIBIOTIC SUSCEPTIBILITY PATTERN OF BACTERIA IN SACHET WATER, SOLD UYO METROPOLIS, AKWA IBOM STATE

ABSTRACT

Aims: This study was carried out to examine the microbiological quality of sachet drinking water sold in Uyo metropolis, Akwa Ibom State.

Study design: Sachet water was randomly sampled in Uyo Metropolis.

Place and Duration of Study: Department of Microbiology, Akwa Ibom State University, Nigeria, between June 2018 and November 2018.

Methodology: Six Different brands of sachets water sold and consumed were studied for their physical and microbiological qualities. Thirty (30) sachets water from six (6) different brands were serially diluted and cultured on Nutrient agar, Eosin Methylene Blue agar, MacConkey agar and Salmonella Shigella agar, while Muller Hinton agar was used for sensitivity test. Pure isolates suspension were standardized with 0.5 McFarland turbidity standard and were subjected to antibiotics susceptibility test using Agar Diffusion method.

Results: The microbial counts ranged from 9.20×10^1 CfU/ml to 1.77×10^2 CfU/ml. Bacterial isolates include; *Klebsiella* sp., *Escherichia* sp., *Staphylococcus* sp., *Salmonella* sp., *Pseudomonas* sp., *Citrobacter* sp., *Bacillus* sp. All the isolates were completely susceptible to Chloramphenicol, Ampiclox, Tarvid and Peflacine. Low percentage of these isolates were resistant to Erythromycin, Gentamycin, Septrin, Ciprofloxacin, Norfloxacin, Chloramphenicol, Levofloxacin and Rifampicin but were highly resistant to Amoxil, Ceporex, Augmentin, Ampicillin, Nalidixic acid and Stretomycin. Some of the sachet water brands from bacteriological standpoints did not meet the World Health Organization Standard.

Conclusion: This study indicted sub-standard packaged waters as a vehicle of spread of antibiotic resistant pathogens, and this poses a high risk to public health. Hence, routine monitoring of producers of sachet water should be enforced.

Keywords: Sachet water, Antibiotics resistance, Public health, water standards.

1. INTRODUCTION

The safety and quality of drinking water has become a public health concern in the world. In Nigeria, high demand of safe drinking water cannot be overemphasized considering the inability of the government to provide adequate pipeborne water to the populace. Water is known to be the dwelling place for most bacteria and other microorganisms which cause a

24 variety of waterborne infections [1] and the World Health Organization (WHO) estimated that
25 1.1 billion of the world's population does not have access to safe water. In addition to this,
26 80% of diseases and one-third of deaths in developing countries are due to consumption or
27 drinking of contaminated water [2]. The associated health risks from the consumption of
28 unsafe drinking water vary throughout the world depending on the chemical or
29 microbiological contaminants present in the environment [3]. Many of the bacteria isolated in
30 water distribution systems are opportunistic pathogens. The presence of high numbers of
31 opportunistic pathogens in drinking water is of concern because these microorganisms can
32 lead to infection of certain segments of the population (newborn babies, the sick, and the
33 elderly) [4]. According to the guideline set by the World Health Organisation, quality drinking
34 water must not contain *Escherichia coli* or thermotolerant coliform bacteria, giardia worms,
35 viruses, *Cryptosporidium* spp, *Legionella pneumophila*, *Entamoeba histolytica* and other
36 opportunistic pathogens such as *Clostridium* species, *Klebsiella* species and *Pseudomonas*
37 [2]. The guideline further stated that the water should be tested against the presence of
38 highly virulent pathogens such as *Salmonella typhi*, *Shigella dysenteriae* and *Vibrio cholera*
39 that are responsible for typhoid, bacillary dysentery and cholera diseases respectively which
40 arises due to high level of organic decay and fermentation on tropical waters. All these
41 bacteria must not exist in water that are meant for drinking, hence, sources of water for
42 packaged water are subjected to laboratory test by public analyst in which any of the
43 bacteria must not be found or detected in any 100 ml water sample. "Sachet water is not
44 sterile" according to Linda [3]. Although, sachet water is assumed to be free from certain
45 pathogen during treatment processes, but certain organisms are used to confirm the sterility
46 of the water such as coliforms which act as indicator organisms used to assess the safety of
47 water and thus give an idea of the degree of contamination associated with intake of such
48 sachet water [4,5]. Antibiotics is been revolutionized in medicine diversely, saving many lives
49 because it had a major impact on the rate of survival of pathogens from infection. But with
50 this great and remarkable benefit, it is sad to know it is also the bedrock of many other
51 diseases due to their resistance strains. Recently, almost all important bacterial infections
52 are becoming resistant to antibiotics, and these changing patterns caused a demand for new
53 antibacterial agents. Antimicrobial resistance occurs when bacteria adjust or adapt in ways
54 that permits them to stay alive in the presence of antibiotics designed to kill them, bacteria
55 evolve resistance to these drugs, typically by acquiring chromosomal mutations and
56 multidrug resistant plasmid, which has become a public health concern [6,7,8]. Antibiotics
57 were formally defined to distinguish their biochemicals which are produced by
58 microorganism from the organic chemicals synthesized in the laboratory. But due to recent
59 development, the distinction between both is no longer meaningful due to the fact that the
60 biochemical structures of many naturally occurring antibiotics are now being synthesized by
61 organic chemist and currently, many antibiotics used in medicals are in the chemically
62 modified forms of microbial biosynthetic form [9].

63 Antibiotic resistance occurs when the sensitivity of an organism decreases against an
64 antibiotic when compared to officially available breakpoints, usually measured as a decrease
65 in "inhibition zone diameter". The increased use of antibiotics is often associated with
66 increased resistance of bacteria to these chemicals, especially in the hospital setting [10]. A
67 lot of transmissible diseases are waterborne. Many harmful microbial contaminants have
68 been confirmed to be associated with potable water sources. Many people have resorted to
69 patronizing sachet water with the belief that it is pure-hence, fondly called 'pure water'. It is
70 possible that this so called pure water is not pure after all; hence it may harbour harmful
71 microorganisms as producers of such water may not pay adequate attention to microbial
72 quality. Identification of the major harmful microbial contaminants (*Escherichia coli*,
73 *Salmonella*, *Shigella*, etc.) present in the sachet water is important in assessing its safety.
74 Free from contamination with faecal matter is the most important parameter of water quality
75 because human faecal matter is generally considered to be a greater risk to human health
76 as it is more likely to contain human enteric pathogens [11]. There is need to constantly

77 access the water quality of water sources available to members of any community at
78 intervals. This will help monitor or track and prevent the sudden outbreak of waterborne
79 infections. It is also important to know the antibiotics susceptibility pattern of microorganism
80 common in our environment in case of any outbreak. This research was borne as a result of
81 the widespread use of sachet water in Nigeria especially in Akwa Ibom State, conflicting
82 results on the safety conducted at different locations in the country and lack of data on safety
83 of sachet water locally available. This research is aimed at determining the antibiotic
84 resistant pattern of bacterial isolates obtained from sachet water by testing them against
85 some of the commonly used antibiotics; taking notes of the antibiotic resistant strains.

88 2. MATERIALS AND METHODS

89 2.1 Study area

91 Three major areas in the major city of Uyo in Akwa Ibom State where strategically selected
92 for this study. The areas comprised of towns where sachet-packaged drinking water is sold
93 by hawkers. They included: Abak road, Aka road and Oron road.

95 2.2 Sample collections

96 A total of Thirty (30) sachet water of six different brands was collected randomly from various
97 part of Uyo metropolis in Akwa Ibom state and taken to the laboratory for analysis. The
98 samples were coded as; BC, GO, FD, RS, ML, and CV to reflect the respective brands. They
99 were collected and transported in clean ice containers and stored at 4.0°C for 30-60 minutes
100 so as to maintain the properties of the sample before commencement of analysis. Hygienic
101 and aseptic methods were also observed during sampling of the sachet water.

103 2.3 Sterilization

104 Microbiologically, sterilization is simply any process that eliminates, removes, kills, all forms
105 of life and any other biological agents (such as bacteria, fungi etc) present in a specific
106 region. This was achieved through the process of autoclaving (steam under pressure) and
107 oven drying. Media, water, and other heat stable liquids were sterilized in the autoclaved at
108 121°C for 15minutes, while glassware were sterilized at 160°C for at least 2 hours. Also, the
109 working environment was always kept neat, tidy and sterile by the process of disinfection.

111 2.4 Enumeration of microbial load in water sample

113 2.4.1 Preparation of the samples

114 Using aseptic method, six (6) different beakers were labelled according to the 6 different
115 brands of waters. Five sachets were mixed from each brand to obtain 100ml homogenous
116 sample in the beaker.

118 2.4.2 Pour plating method

119 One millilitre of appropriate dilution was aseptically pipette into sterile labelled petri dish and
120 this was done in duplicates. Appropriate medium (Nutrient agar, Eosin Methylene Blue,
121 MacConkey agar, Salmonella-Shigella Agar) at 45°C were poured aseptically into the
122 inoculated petri dishes and swirled gently to mix. They were inversely incubated at 37°C for
123 24-48hours. At the end of the incubation period, colonies were counted and the count for
124 each plate expressed as colony forming units per gram (cfu/mL) of the sample suspended.
125 Nutrient agar (NA) to determine the total viable bacterial Count, Eosin Methylene Blue agar
126 (EMB) to enumerate *Escherichia coli*, MacConkey agar (MAC) for coliform count and
127 Salmonella-Shigella agar (SSA) for the determination of *Salmonella* and *Shigella* counts.
128 Culture media were prepared according to the respective Manufacturers specification and
129 sterilized in an autoclave at 121°C at 15 psi for 15 minutes.

130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181

2.4.3 Purification of colonies

Using a fresh nutrient agar medium, 24 hours colonies were picked using a sterile wire loop from the plate and streaked on its surface and incubated for 24 hours at 37°C to obtain pure culture. After incubation, discrete growths were observed on the lines of streak. Distinct colony was picked aseptically and cultured on a fresh nutrient agar slant and incubated for 24 hours at 37°C and stored in a refrigerator at 4°C. The routine laboratory method of Cruickshank *et al.* [12] was used to characterize different isolates. The isolates were identified using their macroscopic, cultural, physiological and biochemical characteristics.

2.5 Morphological characterization (Gram's reaction)

Gram staining was carried out as described by Olutiola *et al.* [13]. Gram stain is one of the differential stains used to characterize bacteria into two main groups: Gram positive and Gram negative. Gram positive stains blue to purple while Gram negative stains pink to red. The colony of the pure cultures of each bacterial isolates was observed for morphological features using Bergey's Manual of Determinative Bacteriology as a standard for comparison. Cell shape was determined under X100 objective of the light microscope after Gram staining procedure. Bacterial smear (not too thick not too thin) was prepared on the slide using an inoculation loop. This was done by introducing a drop of distilled water on grease-free labelled slide followed by the sample and then smeared, air dried and heat fixed. The slide was flooded with crystal violet staining reagent for about 60 seconds, then washed using a gentle indirect stream of tap water for about 2 seconds. The slide was flooded with a mordant (Lugol's iodine) for 15-30 seconds. The slide was decolorized using 70% ethanol for 10 seconds and washed off. Lastly, the slide was flooded with 0.5% counter stain (safranin) for 30 seconds, and then washed using indirect stream of tap water and air dried. A drop of immersion oil was dropped on the stained sample and observed under the microscope.

2.6 Biochemical Characterization and Identification of Isolates

Pure cultures of bacterial isolates were subjected to various biochemical tests according to standard techniques described by Olutiola *et al.* [13] Biochemical tests carried out include; Catalase test, Coagulase test, Indole test, Oxidase test, Citrate test, Fermentation of glucose, lactose, sucrose, maltose and mannitol [14]. Bacterial isolates were identified using Bergey's Manual of Determinative Bacteriology [15].

2.7 Antimicrobial Sensitivity Testing

Commercially available antibiotic impregnated 8mm sensitivity discs (Abtek Biological Ltd, UK) was used to determine the drug sensitivity profile of the isolates. Seventeen different antibiotic discs comprising of Tariviva (OFX), Nalidixic acid (NA), Peflacin (PEF), Gentamycin (CN), Augmentin (AU), Ciproflox (CPX), Septrin (SXT), Ceporex (CEP), Streptomycin (S), Ampicillin (PN) for Gram negative and Levofloxin (Lev), Amoxicillin (Amx), Norfloxacin (NB), Chloramphenicol (CH), Erythromycin (E), Ampiclox (APX), Rifampin (RD), Streptomycin (S), Ciproflox (CPX), Gentamycin (CN) for Gram positive organisms. The antimicrobial sensitivity test of each isolate was carried out as described by the Kirby – Bauer disc diffusion method as recommended by the National Committee for Clinical Laboratory Standards [16]

Procedures: The turbidity of the bacterial suspensions was compared with 0.5 Macfarland's standard by inoculating the organism into 10ml peptone water and incubate. The standardized bacterial suspension was then inoculated on to Muller Hinton Agar and left to dry for 10 minutes, before placing the antimicrobial sensitivity discs. After incubation, the diameter of the zone of inhibition were measured and compared with zone diameter of interpretative chart [17,18] to determine the sensitivity of the isolates to antibiotics.

182 **3. RESULTS**

183

184 All the water samples were National Agency for Food and Drug Administration and Control
185 (NAFDAC) approved and had factory addresses on them (Table 1). They were all odourless,
186 colourless and clear in appearance and had no batch number, also non had production and
187 expiration dates meaning that the duration between production and consumption cannot be
188 determined. Only FD contained little particles in it. Lastly, all were the same net volume of 50
189 cl.

190

191 Table 2 shows the Total viable count (TVC) after 48 hours of water samples on different
192 media. All the water samples were contaminated with bacteria. A higher value of TVC on
193 Nutrient agar (NA) was 1.34×10^2 cfu/ml from sample FD, Eosin Methylene Blue agar (EMB)
194 plate was 3.1×10^1 cfu/ml from sample ML, MacConkey agar (MAC) plate was 25cfu/ml from
195 sample ML and on Salmonella Shigella agar (SSA) plate it was 5.0 cfu/ml from sample FD.
196 The highest number of organisms (on all the media) was 1.77×10^2 cfu/ml in FD sachet water
197 and the lowest was 7.6×10^1 cfu/ml in CV sachet water.

198

199 Out of 29 bacteria isolate, 7 distinct isolates were obtained while others were replicates of
200 the seven. *Klebsiella* sp. had the highest frequency showing 7 out of 29 representing
201 24.14%, followed by both *Staphylococcus* sp. and *Pseudomonas* sp. with the frequency of 5
202 out of 29 isolates representing 17.24%. Other bacteria isolated included; *Escherichia* sp.
203 with the frequency of 4 out of 29 representing 13.79%, *Salmonella* sp. and *Citobacter* sp.
204 with frequency of 3 out of 29 representing 10.34% and *Bacillus* sp. with the least frequency
205 2 out 29 representing 6.90% as shown in Figure 1.

206

207 Six brands of sachet water were analyzed and a total of seven bacterial isolates were
208 identified from the sachet water samples. The isolates were initially differentiated on the
209 basis of the cultural and morphological studies after which they were subjected to various
210 biochemical characterization tests. These tests revealed their probable identity as *Klebsiella*
211 sp., *Escherichia* sp., *Staphylococcus* sp., *Salmonella* sp., *Pseudomonas* sp., *Citrobacter* sp.,
212 *Bacillus* sp.

212

213 *Klebsiella* sp. was most resistant (70%), followed by *Escherichia* sp and *Salmonella* sp. *E*
214 *Escherichia* sp was resistant to 6 (NA, CN, AU, SXT, S, PN and CEP) out of the 10
215 antibiotics tested against it. Same number of antibiotic resistance was recorded for
216 *Salmonella* sp. (NA, CN, AU, S, PN and CEP). The least resistant gram negative isolate was
217 *Citrobacter* sp. (NA, CPX, S, and PN) and *Pseudomonas* sp. All the Gram's negative
218 isolates were resistant to PN and NA. The Gram's positive organisms were less resistant to
219 all the antibiotics they were exposed to. *Bacillus* sp. was resistant to on ciproflox while
219 *Staphylococcus* sp. was resistant to amoxicillin and Gentamycin (Table 3)

220

221

222

223

224

225

226

227

228

229

230

Table 1: Physical examination of the sampled Sachet water brands sold in Uyo metropolis for compliance. Table pattern according to Dada, 2009.

SAMPLE CODE	NAFDAC	PRODUCTION./ BEST FORE DATE	PRODUCERS' NAME & ADDRESS	COLOUR	APPEAR-ANCE	ODOUR	FLOATING PARTICLES	BATCH NO:	NET VOLUME
BC	+	-	+	-	-	-	None	-	50CL
FD	+	-	+	-	-	-	Few	-	50CL
RS	+	-	+	-	-	-	None	-	50CL
CV	+	-	+	-	-	-	None	-	50CL
ML	+	-	+	-	-	-	None	-	50CL
GO	+	-	+	-	-	-	None	-	50CL

231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249

KEY: +: displayed on sample sachet; -: not displayed on sample sachet

250
251
252

Table 2: Total viable count (TVC) after 48hours of culturing sachet water samples on different media


Sample/ Media	NA	EMB	MAC	SSA	Total no. of organism (cfu/ml)
BC	110	30	15	0	1.55×10^2
FD	134	29	9	5	1.77×10^2
RS	70	8	14	0	9.20×10^1
CV	25	31	20	0	7.60×10^1
ML	20	45	25	2	9.20×10^1
GO	118	18	12	1	1.48×10^2

253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286

KEYS: NA: Nutrient Agar; EMB: Eosin Methylene blue agar; MAC: MacConkey agar; SSA: Salmonella Shigella Agar

UNDER PEER REVIEW

287
288



289
290
291
292

FIG. 1: Percentage frequency of bacteria isolates obtained from sachet water sold in Uyo metropolis

293
294
295
296
297
298

TABLE 3. Antibiotics susceptibility pattern of bacterial isolate from sachet water sold in Uyo metropolis.

S/N	Isolate	Gram Positive Isolates										Gram Negative Isolates							% RESISTANCE			
		AMX	S	NB	CPX	CH	E	LEV	CN	APX	RD	OFX	NA	PEF	CN	AU	CPX	SXT		S	PN	CEP
1	<i>E. coli</i>											S	R	S	S	R	S	R	R	R	R	60
2	<i>K. pneumonia</i>											S	R	S	R	R	R	S	R	R	R	70
3	<i>Bacillus sp.</i>	S	R	S	S	S	S	S	S	S	S											10
4	<i>S. aureus</i>	R	S	S	S	S	S	S	R	S	S											20
5	<i>Pseudomonas sp.</i>											S	R	S	S	R	S	S	S	R	R	40
6	<i>Citrobacter sp.</i>											S	R	S	S	S	S	R	R	R	S	40
7	<i>Salmonella sp.</i>											S	R	S	R	R	S	S	R	R	R	60

299
300
301
302
303
304

KEY: Tariviam (OFX), Nalidixic acid (NA), Peflacin (PEF), Gentamycin (CN), Augumentin (AU), Ciproflox (CPX), Septrin (SXT), Ceporek (CEP), Streptomycin(S), Ampicillin(PN) for Gram negative and Levoxin (Lev), Amoxicillin (Amx), Norfloxacin (NB), Chloramphenicol (CH), Erythromycine (E), Ampiclox (APX), Rifampin (RD), Streptomycin (S), Ciproflox (CPX), Gentamycin (CN).

305

306 4. DISCUSSION

307 This experiment was carried out to determine the microbial quality and the antibiotics
308 resistance pattern among the bacterial isolates from sachet water sold in Uyo with the view
309 of creating public health awareness concerning drinking such water sample. In Nigeria,
310 sachet water is largely taken and they are obtained either from surface or underground
311 sources, and are subjected to various treatment to make it fit for individual consumption, but
312 unfortunately, most of them still fall below the WHO standard from the physical and
313 microbiological analysis [19]. From this analysis, one out of six water samples had particles
314 in it. Meanwhile, all were odourless, colourless, and registered with NAFDAC. Bacteria
315 occurrence was recorded in all the sachet-water samples and the TVC for some were higher
316 than what is acceptable for drinking water (1.0×10^1 cfu/ml) [20].

317 The presence of pathogenic bacteria (opportunistic pathogens) was recorded which is above
318 the WHO standard for portable water [4]. High occurrence of *Klebsiella* sp. was recorded,
319 followed by *Staphylococcus* sp. Others included *Pseudomonas* sp., *Escherichia* sp.,
320 *Salmonella* sp., *Citrobacter* sp. and the least frequent was *Bacillus* sp. Total Viable Count on
321 EMB and MAC for coliform bacteria and the various values obtained for each water sample
322 signified possible faecal contamination. This indicates that the sachet-water samples were
323 contaminated especially with faecal materials, thereby not safe for drinking. Presence of
324 coliforms (*Escherichia* sp. and *Klebsiella* sp. and *Citrobacter* sp.) maybe that some of the
325 water were prepared from shallow and contaminated boreholes. Most of these bacteria are
326 indigenous to aquatic environments [20]. *Salmonella* is also as a result of contaminated
327 water and improper treatment, *Pseudomonas* sp. were also found in water and are
328 considered opportunistic pathogens and *Staphylococcus* sp. isolated from the water samples
329 may have entered the water during packaging or handling since the organism is a normal
330 flora of the human skin [21]. The ingestion of these bacteria with contaminated water
331 constitute public health risks to the immunocompromised members of the population,
332 especially newborn babies, elderly and sick people [22]. The presence of relative heavy load
333 of bacteria in water packaged for drinking purposes has been previously documented in
334 literature [23, 24, 25, 26]. The result of the antibiotics susceptibility testing showed various
335 percentages of antibiotic resistance among the bacterial isolates from packaged water
336 samples. *Escherichia* sp. was highly resistant to six antibiotics and sensitive to only four
337 antibiotics which were; Tarivia (OFX), Gentamycin (CN), Peflacin (PEF) and Ciproflox
338 (CPX). *Klebsiella* sp. was resistant to seven antibiotics and sensitive to Tarivia (OFX),
339 Peflacin (PEF) and Septrin (SXT). *Bacillus* sp. was sensitive to all antibiotics and resistant
340 to only Streptomycin (S). *Staphylococcus* sp. was also highly sensitive to all the antibiotics
341 except Amoxicillin (AMX) and Gentamycin (CN). *Pseudomonas* sp. was also sensitive to
342 most antibiotic except Nalidixic acid (NA), Augmentin (AU), Ampicillin (PN) and Ceporex
343 (CEP). *Citrobacter* sp. was more sensitive to the antibiotics and resistant to only four
344 antibiotics; Nalidixic acid (NA), Septrin (SXT), Streptomycin (S), Ampicillin (PN). *Salmonella*
345 sp. was highly resistant to all the antibiotics except four; Tarivia (OFX), Peflacin (PEF),
346 Ciproflox (CPX) and Septrin (SXT). Generally most of the isolates were resistant to Amoxil,
347 Ceporex, Augmentin, Ampicillin, Nalidixic acid and Stretomycin. The resistance exhibited by
348 *Pseudomonas aeruginosa* and *E. coli* to some of the antibiotics corroborates earlier report
349 from South Eastern Nigeria [27]. The presence of the same type of faecal bacteria in almost
350 all brands shows common source of contamination. It is documented that bacteria harbour
351 series of antibiotic resistant genes which can be transferred to others horizontally [28].

352 Therefore from observation, a lot of sachet water producers and sellers have emerged
353 making it their major source of income. With this, appropriate health authorities should
354 ensure that producers comply with the government regulations since some of these
355 packaged water may have been produced under unhygienic conditions. Water can be seen
356 as one of the most important, as well as one of the most abundant of those compounds and
357 it is particularly, vital to living organisms [29]. Also, water is like the life wire of the body and

358 as the basis of life; it is a critical part of human diet. Water constitutes about 90% by weight
359 of the human body [30]. So, water should be treated and the necessary biochemical and
360 microbiological test should be carried out to protect the general public from water-borne
361 disease outbreak.

362

363 5. CONCLUSION

364

365 The study revealed that bacteriological quality of the sachet water brands sold failed to meet
366 the standard for drinking water, even though the bacterial load did not exceed the maximum
367 range of microbial load. However, the bulk of sachet water brands were contaminated by
368 coliform bacteria. It is therefore necessary for sachet water brands to be properly treated and
369 handled to meet the WHO standard for drinking water. To minimise the problem of poor
370 quality of sachet water, government agencies like the NAFDAC and the Environmental
371 Protection Agency should ensure that packaged water manufacturers comply with good
372 manufacturing practices. It is a serious threat to the people of the area if proper
373 measurements are not taken by the concerned authorities. The water sources were
374 contaminated with *Klebsiella* sp., *Escherichia* sp., *Staphylococcus* sp., *Salmonella* sp.,
375 *Pseudomonas* sp., *Citrobacter* sp., and *Bacillus* sp. thus posing a very serious threats to the
376 society. Antibiotic resistance is considered a major problem because many disease causing
377 bacteria are becoming more resistant to the commonly used antibiotics. The bacteria
378 *Klebsiella* sp., *Escherichia* sp., *Citrobacter* sp., showed greater antibiotic resistances. The
379 overuse and misuse of antibiotics can create the conditions for the development of antibiotic
380 resistant bacteria.

381

382 5.1 Recommendation

383 There is need for NAFDAC to intensify efforts in the routine monitoring of activities in the
384 packaged drinking water industry ensuring the safety of sachet drinking water through
385 comprehensive regulatory programs at both the federal and state levels. Also, sample
386 collection and testing of market samples will be a good way of detecting if the water is truly
387 pure as claimed by these producing companies. High emphasis should also be placed on
388 enforcing compliance with Good Manufacturing Practice (GMP) with emphasis on
389 management of raw water source to the consumer product point. Hence, routine monitoring
390 of producers of sachet water should be enforced to ensure adherence to drinking water
391 standard.

392

393

394 COMPETING INTERESTS

395

396 Authors have declared that no competing interest exist.

397

398

399

400 REFERENCES

- 401 1. Spellman FR, Drinan J. The Drinking Water Handbook. Lancaster, Pennsylvania,
402 USA: Technomic Publishing Company Incorporated. 2000. Pp. 260.
- 403 2. World Health Organization. Drinking Water Quality Guideline 4th Edition. World
404 Health Organization (WHO), Geneva, Switzerland. 2011;1-28.
- 405 3. Linda OA, Uchenna CO, Moses NI, Chinelo KU, Charles, OE. Microbial Evaluation
406 and Antibiotic Susceptibility Profile of Isolates of Popular Sachet Water Brands Sold
407 in Anambra State. *British Microbiology Research Journal*, 2016;12(4), 1-9.
- 408 4. Bitton G. *Wastewater microbiology*. 3rd Edition. Wiley series in ecological and
409 Applied Microbiology. 2005

- 410 5. Barrell R, Hunter PG. Microbiological Standards for Water and their Relationship to
411 Health Risk. *Communicable Diseases and Public Health*, 2000;3(1): 8-13.
- 412 6. Finch RG, Greenwood D, Norrby SR, Whitley RJ. Antibiotic and chemotherapy: Anti-
413 infective agents and their use in therapy. 8th ed. Edinburgh: *Churchill Livingstone*.
414 2003. 964.
- 415 7. Nichol K, Zhanel GG, Hoban DJ. Molecular epidemiology of penicillin resistant and
416 ciprofloxacin resistant *Streptococcus pneumoniae* in Canada. *Antimicrobial Agents*
417 *and Chemotherapy*. 2003;47:804-808.
- 418 8. Kummerer K. Resistance in the environment. *Journal of Antimicrobial Chemotherapy*
419 2004;54:311–320.
- 420 9. Sharma BC, Rai B. Incidence of multi-drug resistance in *Escherichia coli* Strains
421 isolated from three lakes of tourist attraction (Mirik Lake, Jorepokhari Lake and
422 Nakhapani Lake) of Darjeeling Hills, India. *Indian Journal of Fundamental and*
423 *Applied Life Sciences*. 2012; 2(2):108-114.
- 424 10. Swartz MN. Use of antimicrobial agents and drug resistance. *The New England*
425 *Journal*. 1997; 45-68.
- 426 11. Scott ME, Melton-Celsa AR, O'Brien AD. Mutations in hns reduce the adherence of
427 Shiga toxin-producing *E. coli* 091:H21 strain B2F1 to human colonic epithelial cells
428 and increase the production of hemolysin. *Microbial Pathogenesis*, 2003;34:155–
429 159.
- 430 12. Cruickshank R, Duguid JP, Marmion BP, Swain, RHA. Medical Microbiology,
431 Volume II, 12th edition. Churchill Livingstone, Edinburgh, London and New York.
432 1975.
- 433 13. Olutiola PO, Famurewa O, Sonntag HG. *An Introduction to general microbiology: A*
434 *practical approach*: HeidelbergerVerlagsansalt und DruckereiGmbH. Heidelberg.
435 1991; P. 267
- 436 14. Harrigan WF, McCance EM. *Laboratory Methods in Food and Dairy Microbiology*.
437 Academic Press, London, New York, San Francisco, 1976; 452 pp.
- 438 15. Buchanan RE, Gibbon NE. *Bergey's Manual of Determinative Bacteriology*. 9th Edition,
439 Williams and Wilkins Co., Baltimore. 1974
- 440 16. National Committee for Clinical Laboratory Standards. Performance standards for
441 antimicrobial disk susceptibility tests. Approved standard M2-A4. Wayne, Pa:
442 National Committee for Clinical Laboratory Standards; 1990.
- 443 17. CLSI 2009. Performance standards for antimicrobial susceptibility testing; 19th
444 informational supplement M100–S19. Clinical and Laboratory Standards Institute,
445 Wayne, PANational Committee for Clinical Laboratory Standards. Performance
446 standards for antimicrobial disk susceptibility tests. Approved standard.
- 447 18. NCCLS document M2-A5. Wayne, Pa: National Committee for Clinical Laboratory
448 Standards; 1993
- 449 19. WHO 2003
- 450 20. Berger, P.S., Oshiro, R.K. Source water protection: Microbiology of source water.,
451 *In: Encyclopedia of Environmental Microbiology*. G.Bitton, Editor-in-chief, Wiley
452 InterScience, New York. 2002; Pp2967-2978
- 453 21. Ollos PJ, Huck PM, Slawson, RM. Factors Affecting Biofilm Accumulation in Model
454 Distribution Systems. *Journal of American Water Works Association*, 2003; 95: 87–
455 97.
- 456 22. LeChevallier MW, Seidler RJ, Evans TM. Enumeration and characterization
457 of standard plate count bacteria in chlorinated and raw water supplies. *Applied*
458 *Environmental Microbiology* 1980; 40:922–930.
- 459 23. Onifade AK, Ilori RM. Microbiological Analysis of Sachet Water Vended in Ondo
460 State, Nigeria. *Environmental Research Journal*, 2008; 2: 107-110.
- 461 24. Oladipo IC, Onyenika IC, Adebisi AO. Microbial analysis of some vended sachet
462 water in Ogbomoso, Nigeria. *African Journal of Food Science*, 2009; 3(12): 406-412.

- 463 25. Oyedeji O, Olutiola PO, Moninuola MA. Microbiological quality of packaged drinking
464 water brands marketed in Ibadan metropolis and Ile-Ife city in South Western
465 Nigeria. *African Journal of Microbiology*. 2010; 4: 96-102.
- 466 26. Onilude AA, Adesina FC, Oluboyede OA and Adeyemi BI. Microbiological
467 quality of sachet packaged water vended in three local governments of Oyo State,
468 Nigeria. *African Journal of Environmental Microbiology*. 2013; 4(9): 195-200.
- 469 27. Nwachukwu E, Emeruem CM. Presence of antibiotic resistant bacteria in sachet
470 water produced and sold in the eastern Nigeria. *Research Journal of Microbiology*,
471 2007; 2(10):782-786
- 472 28. Piddock, L. J. Clinically relevant chromosomally encoded multidrug resistance
473 efflux pumps in bacteria. *Clinical Microbiology Review*, 2006; **19**, 382–402.
- 474 29. Tortora JG, Funke RB, Case LC. Microbiology An introduction. *Media update of 7*
475 *Edition including bibliography and index publisher*. Daryl Fox. 2002; 258-260.
- 476 30. William C, Sonzogoni P, Standridge J, Bussen M. Madison Preservation and
477 survival of *Escherichia coli* in well water sample. *Wisconsin State Laboratory of*
478 *Hygiene, University of Wisconsin*. 2002; 4-10.
- 479
480