

IN-VITRO ANTIOXIDANT AND ANTIBACTERIAL POTENTIAL OF MANNOSE/GLUCOSE-BINDING *PTEROCARPUS OSUN* CRAIB. SEEDS LECTIN**ABSTRACT**

Objective: This study was carried out to purify and characterize a carbohydrate-binding and cell-agglutinating protein, lectin, from *Pterocarpus osun* seeds and also to evaluate its antioxidant and antibacterial potential.

Methods: Isolation and purification of the lectin were done by ammonium sulphate precipitation and size exclusion chromatography on Sephadex G-100. Physicochemical properties of the lectin were determined and antioxidant activity was evaluated by DPPH radical scavenging, lipid peroxidation inhibition and ferric reducing antioxidant potential assay. Disc diffusion method was used for antibacterial effect.

Results: Lectin was detected in the seeds and was able to agglutinate native and enzyme-treated rabbit erythrocytes but only enzyme-treated erythrocytes of human blood were agglutinated. Mannose, Maltose and α -methylmannoside inhibited the divalent cation independent hemagglutinating activity, which was stable up to 60°C and at pH range of 3-13. It showed antioxidant activity with IC₅₀ of 1.17 ± 0.08, 0.58 ± 0.03, and 2.51 ± 0.03 mg/ml for those methods respectively. No antibacterial activity was observed.

Conclusion: *Pterocarpus osun* seeds lectin possess properties similar to other lectins from *Dalbergieae* tribe and its ability to scavenge free radical and inhibit lipid peroxidation show the presence of a valuable health promoting agent in the seeds.

Keywords: Lectin, Hemagglutinating activity, *Pterocarpus osun*, antioxidant, agglutination.

1. INTRODUCTION

Lectins are sugar-binding proteins or glycoproteins which agglutinate erythrocytes and are widely distributed in nature. Lectins have been isolated from various biological sources such as plants, animals and micro-organisms [1,2]. Lectins have been a subject of intense study for more than a decade because they possess various biological activities such as mitogenic and antiproliferative, antiinflammatory, antitumor, antifungal, antibacterial, vasorelaxant, antioxidant

Comment [C1]: Corrected

27 and antihemolytic among others [2-8]. Lectins have the ability to recognize carbohydrate or glycoconjugate and reversibly
28 bind to it through its carbohydrate-binding sites. The binding is with high affinity and specificity and without any chemical
29 modification because lectin has no enzyme-catalytic activity. Lectin can also agglutinate other cells apart from red blood
30 cells. These distinguishes lectin from other carbohydrate-binding proteins and make them valuable tools in
31 biotechnological, pharmacological and therapeutic applications [5,9].

32 The richest sources of lectins in plants are mature seeds, especially those of the legumes, where lectins may
33 constitute one tenth of the seed total protein. Legume lectins are model system of choice to study the molecular basis of
34 protein-carbohydrate interactions because they are not only easy to purify in large quantities, but also exhibit a wide
35 variety of carbohydrate specificities despite strong sequence conservation [10]. The large majority of the leguminous
36 lectins that have been isolated and characterized are from plants belonging to the tribe of *Phaseoleae* and *Dalbergieae* of
37 the *Papilionoideae* subfamily of leguminosae [11]. Worthy of mention is *Pterocarpus angolensis* seeds lectin which has
38 been purified and physicochemically, biochemically and structurally characterized [12,13]. Other seed lectins that have
39 been purified and biochemically characterized from this tribe include galactose-binding lectins from *Lonchocarpus*
40 *capassa* [14], *Vatairea macrocarpa* [11] and *Vatairea guianensis* [15] and mannose-binding lectins from *Platymiscium*
41 *floribundum* [16] and *Centrolobium tomentosum* [17]. Amino acid sequence of *P. floribundum* [16], *C. tomentosum* [17]
42 and *Centrolobium microchate* [18] lectins has been determined partially. Generally, legume lectins are structurally
43 homologous and at time have similar physicochemical properties but display biological activity that are distinctly differ.
44 Consequently, each of the lectins has the potential for different application and deserves to be independently studied.

45 *Pterocarpus osun* Craib belongs to the *Dalbergieae* tribe of Papilionaceae subfamily. *P. osun* is endemic to
46 Southern Nigeria, Equatorial Guinea, Gabon, Cameroun and Zaire [19]. It exists as a tree of about 30 meters height and
47 2.5 meters girth with a spreading crown and the wood marketed as African Padauk. The leaves of *P. osun* are used in the
48 treatment of skin disease such as eczema, candidiasis and acnes [20]. The crude extract of *P. osun* has also been found
49 useful in the treatment of chicken pox in children in the eastern part of Nigeria [21]. The antioxidant potential and the
50 attenuation of acetaminophen-induced redox imbalance by *P. osun* were reported recently [22]. Adewuyi *et al.* [23]
51 showed that the acetonides prepared from the seed oil of *P. osun* has no antibacterial activities but the leaves ethanolic
52 extract of the plant does.

53 In our preliminary study [24], the presence of hemagglutinin in the crude protein extract of *P. osun* seeds was
54 established but the lectin was not purified and characterized. The present study, therefore, focused on purification and

55 physicochemical characterization of lectin from *P. osun* seeds and also, we investigated its *in-vitro* antioxidant and
56 antibacterial activities.

57 2. MATERIALS AND METHODS

58 2.1 Preparation of crude extract

59 The dried mature seeds of *Pterocarpus osun* were removed from the pods and ground into powder using seed
60 blender, after which 50 g of the powder was defatted using petroleum ether and later was extracted in ten volumes of 25
61 mM phosphate buffer saline (PBS, pH 7.2) containing 10mM sodium chloride. After stirring on magnetic stirrer for about
62 10 hrs, the mixture was centrifuged at 10,000 rpm for 20 min and the supernatant was collected into a sample bottle and
63 stored at – 20 °C until used.

64 2.2 Erythrocytes glutaraldehyde fixation and trypsinization

65 Glutaraldehyde was used to fixed the erythrocytes of human and animal bloods following the methods described by Kuku
66 and Eretan, [25]. Heparinized bottles were used to collect the blood samples, which was centrifuged at 3,000 rpm for 15
67 min. The erythrocytes were thoroughly washed with PBS, pH 7.2. 2% of the erythrocyte was prepared with chilled 1%
68 glutaraldehyde-PBS (v/v) solution. The suspension was incubated at 4 °C for 1 hr with occasional mixing. This was
69 followed by centrifugation at 3,000 rpm for 5 min and several washing of the fixed blood cells with PBS to remove
70 glutaraldehyde. The fixed cells were suspended in PBS to a final concentration of 2%. Trypsinization of the erythrocytes
71 was carried out as described by Occena *et al.* [26]. 2% erythrocytes suspension in PBS was obtained by thoroughly
72 washing the whole blood samples of blood groups A, B, O and animals with PBS. Equal volume of 2% erythrocytes
73 suspension and 1% trypsin solution was mixed and incubated for 1 hr at 37 °C. The trypsinized cells were washed five
74 times with PBS and finally diluted to obtained 2% (v/v) trypsinized cells in PBS. This was stored until further use.

75 2.3 Hemagglutination assay and blood group specificity

76 A two-fold serial dilution of *P. osun* seeds lectin solution (100 µl) was performed in U-shaped microtitre plates. This was
77 mixed with 50 µl of a 2% suspension of human as well as animal (rabbit and rat) or 2% trypsinized erythrocytes in
78 phosphate buffered saline, pH 7.2 at room temperature. The erythrocytes have been previously fixed with 1%
79 glutaraldehyde. The plate was left undisturbed for 2 hr for agglutination to take place. The hemagglutination titre of the
80 lectin expressed as the reciprocal of the highest dilution of the lectin exhibiting visible agglutination of erythrocytes was

81 equivalent to one hemagglutinating unit. Specific activity was the number of hemagglutination units per mg protein. The
82 blood group specificity of the crude lectin extract was determined using erythrocytes from different blood groups of the
83 ABO system and those of the rabbit and rat.

84 **2.4 Sugar specificity test**

85 The sugar specificity of the lectin was investigated by the ability of sugars to inhibit the agglutination of human
86 erythrocytes [27]. A serial dilution of the crude lectin sample was made until the end-point causing hemagglutination was
87 obtained. 0.2 M of each sugar solution was added to each well at 50 μ l per well and allowed to stand for 1 hr undisturbed
88 on the laboratory bench and then mixed with 50 μ l of 2% rabbit erythrocyte suspension. The hemagglutination titre
89 obtained were compared with a non-sugar containing blank. The sugars tested are: maltose, D (+)-mannose, lactose, L
90 (+)-arabinose, sorbose, D (+)-glucose, sucrose, galactose, mannitol, N-acetyl-D-glucosamine, mannosamine, 2-deoxy-D-
91 glucose, dulcitol, xylose, α -D-methyl glucopyranoside and D (+)-glucosamine HCl, α -D-methyl-mannoside.

92 **2.5 Temperature, pH and EDTA Effect on hemagglutinating activity**

93 Thermal stability of the lectin was tested by incubating the purified lectin at different temperature ranging from 30 $^{\circ}$ C – 100
94 $^{\circ}$ C in a water bath for 1 hr. At 15 min interval, for each temperature, hemagglutinating activity of aliquots taken was
95 determined by hemagglutination assay. Control was the lectin kept at room temperature and represent 100%
96 hemagglutinating activity.

97 Hemagglutinating activity of the lectin at both basic and acidic condition was tested. The purified lectin was incubated with
98 buffers of different pH values ranging from pH 3.0 – 13.0. for 1 hr. hemagglutinating activity of the lectin was determined
99 and compared with the control which was lectin incubated in PBS (pH 7.2). Buffers used were 0.2 M citrate buffer (pH 3.0
100 - 5.0), 0.2 M Tris-HCl buffer (pH 6.0 - 8.0) and 0.2 M glycine-NaOH buffer (pH 9.0 - 13.0).

101 To determine if the lectin require divalent metal ion for its hemagglutinating activity, lectin was dialyzed against 10 mM
102 and 100 mM EDTA for 24 hrs. Hemagglutinating activity of the resulting lectin was determined. This was followed by
103 incubating the treated lectin with 10 mM of each of the following divalent cation salts: CaCl_2 , MgCl_2 , MnCl_2 , BaCl_2 and
104 SnCl_2 for 2 hrs in order to determine if the hemagglutinating activity be restored.

105 **2.6 Purification of *P. osun* lectin**

106 **2.6.1 Ammonium sulphate precipitation**

The crude lectin extract of the *P. osun* seeds was subjected to 70% ammonium sulphate precipitation. The ammonium sulphate equivalent to 70% precipitation was slowly and continuously added to the crude extracts on magnetic stirrer to aid dissolution of the salt. The mixture was centrifuged after 24 hrs at 3500 rpm for 15 min to obtain the precipitate. The precipitate was dialyzed exhaustively against several changes of PBS to remove the salt. The dialysate was centrifuged at low speed to remove undissolved materials.

2.6.2 Gel-filtration on Sephadex G-100

The dialysate of ammonium sulphate precipitate of *P. osun* crude lectin extract was applied on Sephadex G-100 column (2.5 x 40 cm) previously equilibrated with PBS, pH 7.2. The protein was eluted with the same buffer at a flow rate of 15 ml/hr and 5 ml fractions were collected. The fractions were monitored for protein by measuring the absorbance at 280 nm and assayed for hemagglutinating activity.

2.6.3 Determination of protein concentration

Protein concentration of the crude extract, dialysate and other fractions were determined by the method of Lowry *et al.* [28] using Bovine Serum Albumin (BSA) as standard protein. The absorbance at 280 nm was also used to monitor protein elution in the chromatographic fractions.

2.7 Physicochemical characterization of purified lectin

2.7.1 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay

Ability of the lectin to scavenge DPPH radical was evaluated by method described by Brand-Williams *et al.*[28] with slight modification. Equal volume (1 ml each) of 0.3 mM DPPH and varying concentration of lectin or standard (ascorbic acid) were mixed. The mixture was incubated in the dark for 30 min. Negative control was prepared by addition of 1 ml methanol instead of lectin. Absorbance of the test and control was read at 517 nm. The percentage of DPPH radical scavenging activity inhibition was obtained using this equation.

$$\text{DPPH radical scavenging inhibition \%} = \left[1 - \left(\frac{\text{Abs}_{\text{sample}}}{\text{Abs}_{\text{control}}} \right) \right] \times 100$$

Where:

$\text{Abs}_{\text{sample}}$ = Absorbance of the lectins

$\text{Abs}_{\text{control}}$ = Absorbance of the control at 517 nm

Sample concentration providing 50% inhibition (IC_{50}) was calculated from the graph by plotting inhibition percentage against sample concentration.

2.7.2 Lipid peroxidation inhibition assay

135 Lipid peroxidation was carried out according to the methods of Ohkawa *et al.* [30] as described by Hattori *et al.* [31] with
136 slight modification and BHT was used as standard. 10% egg yolk homogenate was prepared in 150 mM Tris-HCl buffer
137 (pH 7.2). Five hundred microlitres (500 μ l) of the egg yolk homogenate was added to 0.1 ml of varying concentration of
138 the purified lectin and standard (BHT) separately. Then, 50 μ l of 1% ascorbic acid was added to the reaction mixture,
139 followed by 50 μ l of 0.07 M FeSO₄ to induce lipid peroxidation. The reaction mixture was vortexed and incubated at 37 °C
140 for 1h. Sequential addition of 0.5 ml of 0.1N HCl and 2 ml of 0.67% (w/v) Thiobarbituric acid prepared in 9.8% SDS
141 followed incubation. The resulting mixtures were heated in water bath at 95 °C for 1 h, cooled and 2.0 ml of butan-2-ol was
142 added and later centrifuged at 3,000 rpm for 10 min. The control was run as above with the lectin/standard replaced with
143 distilled water. The supernatant was collected and measured at 532 nm. Percentage inhibition of lipid peroxidation was
144 calculated as:

$$145 \quad \% \text{ Inhibition} = \frac{\text{Abs}_{\text{control}} - \text{Abs}_{\text{test}}}{\text{Abs}_{\text{control}}} \times 100$$

146 where Abs_{control} = MDA produced by fenton reaction in the absence of extract (control);

147 Abs_{test} = MDA produced by fenton reaction in the presence of extract.

148 **2.7.3 Ferric reducing antioxidant power (FRAP) assay**

149 Ferric reducing antioxidant power assay was carried out spectrophotometrically adopting the method described by Benzie
150 and Strain [32] with minor modification. The FRAP working reagents was prepared by mixing ten parts of 300 mM acetate
151 buffer (pH 3.6), one part of 10 mM TPTZ (2,4,6-tripyridyl-s-triazine) and one part of 20 mM of ferric chloride solution in the
152 dark. Fifty microlitres of varying concentration of the lectin and standard solution of the ascorbic acid was added to 1 ml of
153 FRAP working reagent. The mixture was vortexed before incubating at 37 °C for 30 min in the dark. The absorbance was
154 taken at 593 nm against the reagent blank containing 1 ml of the FRAP working reagent and 50 μ l of methanol. All
155 measurements were taken at room temperature and the reducing power was expressed as equivalent concentration
156 which is defined as the concentration of antioxidant that gave a ferric reducing ability equivalent to that of the ascorbic
157 acid standard (AAE).

158 **2.8 Antibacterial assay**

159 **2.8.1 Antibacterial sensitivity test**

160 The *in vitro* sensitivity tests of the bacteria to the purified lectins were done by disc diffusion method described by
161 Akinpelu *et al.* [33] with little modification. About 1 ml of the standardized 24 hrs old culture of the test organisms in

162 nutrient broth was inoculated into pre-sterilized molten Mueller-Hinton agar medium in MacCartney bottle. The medium
163 was poured into a sterile Petri dish and allowed to set. With the aid of a sterile cork borer, three wells of about 6 mm in
164 diameter were bored on the plates equidistant from the centre of the plates. About 0.1 ml of each of the purified lectins (5
165 mg/ml) was dispensed into the wells in each of the Petri dishes. The same volume of antimicrobial standard drugs-
166 streptomycin (1 mg/ml) was dispensed into the third well in the Petri dishes. The plates were incubated at 37 °C overnight.
167 At the end of the incubation period, zones of inhibition formed on the agar plates were measured. Zones of inhibition
168 indicate susceptibility of the test bacteria to the lectin suspension and were evaluated in mm.

169 **2.8.2 Bacterial agglutination test**

170 Bacteria were tested for agglutination with the purified lectins. Both Gram negative and Gram-positive bacteria
171 were grown in nutrient broth for about 24 hrs. The cells were harvested by centrifugation at 3000 rpm for 2 minutes and
172 washed with PBS three times. The packed cells were suspended in 0.5% formalin solution and shaken at 25 °C for 24 hrs.
173 Formalin-killed cells were collected by centrifugation, washed with PBS and resuspended in PBS to 1.5×10^8 colony
174 forming unit/ml (McFarland 0.5 standard). Agglutination assay with the formalin-killed cells was performed in microtitre
175 plates. An equal volume of each bacterial suspension was mixed with a two-fold serial dilution of the lectin in a microtitre
176 plate and incubated at room temperature for one hour. The bacterial agglutination titre was expressed as the reciprocal of
177 the highest dilution giving a visible agglutination upon illumination of the microtitre plates [34].

179 **3. RESULTS AND DISCUSSION**

180 *Pterocarpus osun* seeds lectin (POSL) was easily purified by combination of salt precipitation using ammonium
181 sulphate and size exclusion chromatography on Sephadex G-100. The soluble crude protein extract obtained by PBS
182 extraction of *P. osun* seeds powder was initially precipitated by addition of ammonium sulphate up to 70% saturation and
183 active dialysate obtained after exhaustive dialysis was layered on Sephadex G-100 gel filtration column. The elution
184 profile (Figure 1) presented three distinct protein peaks (GO1, GO2, GO3), where only the third peak (GO3) displayed
185 hemagglutinating activity against rabbit erythrocyte. Similar purification procedures were employed by Galbraith and
186 Goldstein [35] and eLacerda *et al.* [36]. In both studies, ammonium sulphate precipitation of the protein preceded size
187 exclusion chromatography on Sephadex G-200 and Sephadex G-100, respectively. Three distinct protein peaks were
188 obtained by eLacerda *et al.* [36] who worked on Brazilian lima bean variety and only the first peak exhibited
189 hemagglutinating activity. The specific activity of the purified lectin was 119.1 HU/mg protein leading to protein purification
190 of 46-fold (Table 1).

191 Phosphate buffer saline extraction produced a soluble crude lectin extract that showed measurable
192 hemagglutinating activity against trypsin-treated and native rabbit erythrocyte with higher hemagglutinating titre for
193 enzyme-treated erythrocyte. The crude lectin extract was unable to agglutinate native human erythrocyte but trypsinized-
194 human erythrocyte of all ABO blood group was considerably and non-specifically agglutinated. The results are shown in
195 Table 2. Similar results were reported for lectins from *Platymiscium floribundum* [16], *Centrolobium microchaete* [18] and
196 *Canavalia virosa* [37]. Lis and Sharon [38] revealed that trypsin can be used to modify the erythrocytes surface to
197 enhance its affinity for lectins without affecting the total number of lectin binding sites on the erythrocytes. In supporting
198 this statement Singh and Saxena [39] stated that trypsinization of red blood cells may removes the sialoglycopeptide of
199 the cells; thus, demolishing the negative charge on the surface of the cells, which may lead to decrease in repulsive force
200 between the cells and hence increase in agglutination.

201 Inhibition of hemagglutination by several different sugars showed that the lectin activity was strongly inhibited by
202 glucose, its epimer – mannose and their derivatives like, 2-deoxy-D-glucose, N-acetyl-D-glucosamine, α -D-methyl-
203 mannoside, α -D-methyl-gluocopyranoside and a disaccharide (maltose). Complete inhibition of the hemagglutinating
204 activity was noticed with mannose, α -D-methyl-mannoside and maltose. Maltose exhibited most potent inhibitory effect
205 with minimum inhibitory concentration of 260 μ M followed by α -D-methyl-mannoside and mannose (Table 3). These
206 results indicate that presence of another glucose unit at the carbon-1 of the first glucose in the disaccharide increases the
207 interaction with the hydrophobic regions of the carbohydrate-binding site, thereby increased the affinity of the POSL when
208 compared with glucose. Availability of methyl group on α -methyl-mannoside may also causes the same interaction that
209 resulted in higher affinity of the POSL for α -methyl-mannoside than mannose. POSL belongs to the mannose/glucose
210 specificity group of lectins from Dalbergieae tribe, which have specificity for different sugars. Among well studied member
211 of the tribe that belong to mannose/glucose specificity group are lectins from *Pterocarpus angolensis* [3], *Platymiscium*
212 *floribundum* [16], *Centrolobium microchaete* [18], and *Platypodium elegans* [40]. Though, other members of the tribe that
213 have specificity for other carbohydrates especially galactose have also been reported (*Vatairea marcocarpa* [11];
214 *Lonchocarpus capassa* [14]; *Vatairea guianensis* [15]). The biological importance of mannose-binding lectin also has been
215 stretched [41].

216 POSL was thermostable, demonstrating full activity up to 70 °C during 15 min of heating. Fifty percent of the full
217 activity was lost when heated for 60 min at 70 °C while retaining 100% activity at 60 °C for 60 min and no hemagglutinating
218 activity was detected when the lectin was heated at 80 °C for 15 minutes (Figure 2 A and B). This implies that the lectins
219 undergo conformational changes under extreme temperatures resulting in the loss of activity. The loss of activity of the
220 lectins with increased temperature is due to destabilization of sporadic weak interactions of tertiary structure responsible

221 for native conformation of lectin [39]. These results are comparable to the reported results of lectins purified from *Vatairea*
222 *marcocarpa* [11], *Vatairea guianensis* [15], *Platymiscium floribundum* [16], and *Canavalia oxyphylla* [42]. In contrast,
223 extremely thermostable lectins have been reported from *Bauhinia forficata* [43] and *Apuleia leiocarpa* [44]. These lectins
224 retain maximum hemagglutinating activity when heated at 100 °C. High thermostability possessed by these lectins may be
225 advantageous, as stable bioactive substance is more efficient during all phases of their processing and on the other hand
226 this is considered as antinutritional factors that cause many adverse phenomena in animals if ingested [36].

227 Lectin, generally, are found stable in harsh conditions such as extreme pH. POSL was subjected to
228 hemagglutination assay at different pH and the lectin retained maximum activity within a broad pH range (pH 3-13) (Figure
229 3). The result suggest that the lectin was insensitive to acidic and basic pH. *Phaseolus lunatus* seeds lectin exhibited
230 hemagglutinating activity within a broad range, remaining stable between pH 2 and 11 [36]. Other lectins with similar pH
231 optimum have been reported [45,46]. Some lectins have shown that extreme pH is less favorable conditions for their
232 hemagglutinating activity. They are found to display maximum activity at around neutral pH. Lectins from *Vatairea*
233 *guianensis* [15], *Platymiscium floribundum* [16] and *Centrolobium microchaete* [18] retained full hemagglutinating activity
234 at pH 6-9. They all belong to the same Dalbergieae tribe with the *P. osun*.

235 The hemagglutinating activity of POSL remain unchanged after dialysis against or incubation with high
236 concentration of EDTA and addition of divalent cations to the EDTA-treated lectin also did not alter the activity. These
237 probably suggest that POSL does not need divalent cations for it to be active or the metal ions are tightly bound to the
238 lectin. The hemagglutinating activity of lectin from *Vatairea guianensis* [15] and *Platymiscium floribundum* [16] among
239 others showed similar results when incubated with EDTA. Although, this is in contrast to the *P. angolensis* [13] and
240 *Centrolobium microchaete* [18] lectins that completely lost their activity after treatment with EDTA and only addition of
241 metal ions restored their full lectin activity.

242 Apart from peptides, obtainable by either enzymatic hydrolysis, chemical hydrolysis or bacterial fermentation, that
243 have been established to possess strong antioxidant ability, numerous evidences exist that proteins possess antioxidant
244 activity and that these antioxidant proteins have been closely linked to the control of some neurodegenerative and
245 cardiovascular diseases because of its ability to ameliorate the harmful effect of free radicals and reactive oxygen species
246 produced during oxidative stress. DPPH radical scavenging, lipid peroxidation inhibition and ferric reducing antioxidant
247 power assays were used to assess the antioxidant potential of POSL. The results revealed that POSL possess significant
248 antioxidant activity, which were concentration dependent (Figure 4A and B). The lectin showed an IC₅₀ of 1.17 ± 0.08, 0.58
249 ± 0.03, and 2.51 ± 0.03 mg/ml for those methods respectively. These results give support to reported studies that

250 detected lectins with antioxidant potential in some leguminous seeds [35, 46-47] and also to other reported antioxidant
251 proteins from other plant family [48-50]. Though, antioxidant activity in protein possibly will not be ascribed to a single
252 mechanism. Elias *et al.* [51] stated some plant proteins can inhibit lipid oxidation via numerous pathways and inactivate
253 reactive oxygen species and other free radicals, chelate transition metals and reduce hydroperoxides. Presence of some
254 amino acids in the primary structure of this lectin may have contributed to the observed antioxidant ability. Therefore,
255 hydrolysis of POSL may ascribe more antioxidant potential to the peptides that will be generated.

256 The antimicrobial roles of lectins as stated by Coelho *et al.* [52] include blockade of invasion and infection, inhibition
257 of growth and germination, regulation of microbial cell adhesion and migration. There is an increasing interest in
258 investigation of the lectin's involvement in the interaction between eukaryotic cells and pathogens in infectious disease
259 development and their antimicrobial potential [53]. Carvalho *et al.* [43] reported that *Apuleia leiocarpa* seed lectin (ApulSL)
260 demonstrated bacteriostatic effects on the Gram-positive bacteria *Staphylococcus aureus*, *Streptococcus pyogenes*,
261 *Enterococcus faecalis*, *Micrococcus luteus*, *Bacillus subtilis* and *Bacillus cereus*, and on the Gram-negative bacteria
262 *Xanthomonas campestris*, *Klebsiella pneumoniae*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Salmonella enteritidis*.
263 ApulSL was also bactericidal against three varieties of *Anthomonas campestris*. Also, in their studies Mishra *et al.* [54]
264 showed that *Bauhinia variegata* lectin (BVL) demonstrated a remarkable antibacterial activity against the pathogenic
265 bacteria *P. aeruginosa*, *S. aureus*, *E. coli*, and *B. subtilis*. BVL also shows a significant antifungal activity
266 against *Aspergillus niger* and *Penicillium crysogenum*. The results of the present antibacterial study are contrary to the
267 majority of the lectins from leguminous family. The present study showed that purified POSL have no antibacterial activity
268 against both gram-positive and gram-negative bacteria strain used and was unable to agglutinate these pathogens. But
269 the crude protein extract demonstrated significant antibacterial effect against gram-positive bacteria (*B. cereus*, *S. aureus*
270 and *B. subtilis*) and gram-negative bacteria (*Pseudomonas fluorescens*, *K. pneumoniae*, *E. coli*, *P. aeruginosa* and
271 *Proteus vulgaris*). But also, could not agglutinate them. I can therefore be concluded that the antibacterial activity
272 exhibited by the crude protein extract is not due to the presence of lectin but possibly to another antibacterial proteins or
273 peptides. Several antibacterial peptides and proteins have been isolated from plants [55,56].

274 4. CONCLUSION

275 *Pterocapus osun* seeds lectin (POSL) was purified and characterized. POSL was inhibited by mannose, maltose
276 and α -methylmannoside, indicating that it belongs to mannose/glucose-binding lectin. POSL agglutinated native rabbit
277 erythrocytes and nonspecifically enzyme-treated human blood group ABO erythrocytes. The lectin was found stable up to

278 70 °C, active in a wide pH range and requiring no divalent cations for its full activity. No antibacterial activity was detected
279 but exhibited significant antioxidant activity.

280 CONFLICT OF INTEREST

281 There are no conflicts of interest among the authors.

282

283

284 REFERENCE

- 285 1. Sharon N, Lis H. History of lectins: From hemagglutinins to biological recognition molecules. *Glycobiology*. 2004;
286 14:53R-62R.
- 287 2. Hamid R, Masood A, Wani IH, Rafiq S. Lectins: Proteins with diverse applications. *J. Appl. Pharma. Sci*. 2013;
288 3(4):S93-S103. DOI: 10.7324/JAPS.2013.34.S18
- 289 3. Lam SK, Ng TB. Lectins: production and practical applications. *Appl. Microbiol. Biotech*. 2011; 89:45–55.
290 DOI:10.1007/s00253-010-2892-9
- 291 4. Kumar KK, Chandra KL, Sumanthi J, Reddy GS, Shekar PC, Reddy B. Biological roles of lectins: A review. *J.*
292 *Orofac. Sci*. 2012; 4(1):20-25. DOI: 10.4103/0975-8844.99883
- 293 5. Santos AFS, da-Silva MDC, Napoleao TH, Paiva PMG, Correia MTS, Coelho LCBB. Lectins: function, structure,
294 biological properties and potential application. In: *Current Topics Pept. Prot. Res*. 2014; 15:41-62.
- 295 6. Dan X, Liu W, Ng TB. Development and applications of lectins as biological tools in biomedical research. *Med. Res.*
296 *Rev*. 2016; 36:221-247. DOI:10.1002/med.21363
- 297 7. Lagarda-Diaz I, Guzman-Partida AM, Vazquez-Moreno L. Legume lectins: Protein with diverse applications. *Int.*
298 *J.Molec. Sci*. 2017; 18(6): E1242-E1256. doi: 10.3390/ijms18061242
- 299 8. Hendrickson OD, Zherdev AV. Analytical application of lectin. *Critical Rev. Analyt. Chem*. 2018; 48(4):279-290.
300 DOI:10.1080/10408347.2017.1422965
- 301 9. Coelho LCBB, Silva PMS, Lima VLM, Pontual EV, Paiva PMG, Napoleao TH, Corriea MTS. Lectins, interconnecting
302 proteins with biotechnological/pharmacological and therapeutic applications. *Evid. Based Complement. Alternat.*
303 *Med*. 2017; 2017: Article ID 1594074. DOI: 10.1155/2017/1594074
- 304 10. Peumans WJ, van Damme EJ. Lectins as plant defense proteins. *Plant Physiol*. 1995; 109(2): 347–352.

- 305 11. Cavada BS, Santos CF, Grangeiro TB, Nunes EP, Sales PV, Ramos RL, et al. Purification and characterization of a
306 lectin from seeds of *Vatairea macrocarpa* Duke. *Phytochem.* 1998; 49: 675 – 680.
- 307 12. Loris R, VanWalle I, De Greve H, Beeckmans S, Deboeck F, Wyns L, Bouckaert J. Structural basis of
308 oligomannose recognition by the *Pterocarpus angolensis* seed lectin. *J. Molec. Biol.* 2004; 335(5):1227-1240.
309 DOI:10.1155/2017/1594074.
- 310 13. Echemendia-Blanco D, Van-Driessche EV, Ncube I, Read JS, Beeckmans S. Stability, subunit interactions and
311 carbohydrate-binding of the seed lectin from *Pterocarpus angolensis*. *Prot. Pept. Lett.* 2009; 16(9): 1120-1134.
- 312 14. Joubert EJ, Sharon N, Merrifield EH. Purification and properties of a lectin from *Lonchocarpus capassa* (apple leaf)
313 seed. *Phytochemistry* 1986; 49:675-680. doi.org/10.1016/S0031-9422(00)85474-6.
- 314 15. Silva HC, Nagano CS, Souza LAG, Nascimento KS, Isidro R, Delatorre P, et al. Purification and primary structure
315 determination of a galactose-specific lectin from *Vatairea guianensis* Aublet seeds that exhibits vasorelaxant effect.
316 *Process Biochem.* 2102; 47:2347-2355. doi.org/10.1016/j.procbio.2012.09.014
- 317 16. Pereira-Junior FN, Silva HC, Freitas BT, Rocha BAM, Nascimento KS, Nagano CS, et al. Purification and
318 characterization of a mannose/N-acetyl-D-glucosamine-specific lectin from the seeds of *Platymiscium floribundum*
319 Vogel. *J. Molec. Recogn.* 2012; 25:443-449. doi.org/10.1002/jmr.2207.
- 320 17. Almeida AC, Silva HC, Pereira-Junior JB, Cajazeiras P, Delatorre CS, Nagano KS, et al. Purification and partial
321 characterization of a new mannose/glucose-specific lectin from *Centrolobium tomentosum* seeds exhibiting low
322 toxicity on *Artemia* sp. *Int. J. Indig. Med. Plants* 2014; 47:1567-1577. DOI: 10.12817/20514263.47.1.27703470
- 323 18. de Vasconcelos MA, Alves AC, Carneiro RF, Dias AHS, Martins FWV, Cajazeiras P, et al. Purification and primary
324 structure of a novel mannose-specific lectin from *Centrolobium microchaete* Mart seeds. *Int. J. Biol.Macromolec.*
325 2015; 81:600-607. doi.org/10.1016/j.ijbiomac.2015.08.059.
- 326 19. Burkill HM. *The useful plants of West Africa*, 2nd edition, Kew, Richmond, UK: Royal Botanic Garden. 1995; 3:431-
327 432
- 328 20. Gill LS. *Ethnomedical uses of plants in Nigeria*. Uniben Press, University of Benin. 1992:200-201
- 329 21. Ezeokonkwo MA, Okoro UC. New dyes for petroleum products. *J. Emerg. Trends Eng. Appl. Sci.* 2012; 3:8-11.
- 330 22. Ajiboye TO, Salau AK, Yakubu MT, Oladiji AT, Akanji MA, Okogun JI. Acetaminophen perturbed redox homeostasis
331 in Wistar rat liver: protective role of aqueous *Pterocarpus osun* leaf extract. *Drug Chem.Toxicol.* 2010; 33: 77-87.
332 DOI: 10.1080/01480540903170746.

- 333 23. Adewuyi A, Fasusi OH, Oderinde RA. Antibacterial activities of acetone extracts prepared from the seed oils of
334 *Calophyllum inophyllum* and *Pterocarpus osun*. J. Acute Med. 2014; 4: 75-80. doi.org/10.1016/j.jacme.2014.02.001
- 335 24. Odekanyin OO, Akeredolu EO. Comparative investigation of the biological activity of the *Pterocarpus osun* and
336 *Bosqueia angolensis* seeds protein extracts. Int. J. Curr. Res. Biosci. Plant Biol. 2016; 3(9):16-26.
337 DOI: 10.20546/ijcrbp.2016.309.003
- 338 25. Kuku A, Eretan OB. Purification and partial characterization of a lectin from the fresh leaves of *Kalanchoe crenata*
339 (Andr.) Haw. BMB Reports 2004; 37(2):229-233. doi.org/10.5483/BMBRep.2004.37.2.229
- 340 26. Occena IV, Mojica EE, Merca FE. Isolation and partial characterization of a lectin from the seeds of *Artocarpus*
341 *camansi* Blanco. Asian J. Plant Sci. 2007; 6(5):757-764.
- 342 27. Eghianruwa Q, Odekanyin O, Kuku A. Physicochemical properties and acute toxicity studies of a lectin from the
343 saline extract of the fruiting bodies of the shiitake mushroom, *Lentinula edodes* (Berk). Int. J. Biochem. Mol. Biol.
344 2011; 2(4):309-317.
- 345 28. Lowry OH, Rosenbrough NJ, Farr AL, Randall R. Protein measurement with the Folin phenol reagent. J. Biol.
346 Chem. 1951; 193:265-275.
- 347 29. Brand-Williams W, Cuvelier ME, Bereset CLWT. Use of a free radical method to evaluate antioxidant activity. LWT-
348 Food Sci. Technol. 1995; 28(1):25-30. doi.org/10.1016/S0023-6438(95)80008-5
- 349 30. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. Anal.
350 Biochem. 1979; 95:351-358. DOI:10.1016/0003-2697(79)90738-3
- 351 31. Hattori M, Yang XW, Miyashiro H, Nabma T. Inhibitory effects of monomeric and dimeric phenylpropanoids from
352 mice on lipid peroxidation *in vivo* and *in vitro*. Phytotherapy Res. 1993; 7:395-401. doi.org/10.1002/ptr.2650070603
- 353 32. Benzie IF, Strain JJ. (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power" the
354 FRAP assay. Anal. Biochem. 1996; 239(1): 70-76. DOI:10.1006/abio.1996.0292
- 355 33. Akinpelu DA, Aiyegoro AO, Okoh AI. Studies on the biocidal and cell membrane disruption potentials of stem bark
356 extracts of *Azela Africana* (Smith). Biol. Res. 2009; 42: 249-349. DOI:/S0716-97602009000300008
- 357 34. Dutta S, Sinha B, Bhattacharya B, Chatterjee B, Mazumder S. Characterization of a galactose binding serum lectin
358 from the Indian catfish, *Clarias batrachus*: Possible involvement of fish lectins in differential recognition of
359 pathogens. Comp. Biochem. Physiol. 2005; 141(1): 76-84. DOI:10.1016/j.cca.2005.05.009
- 360 35. Galbraith W, Goldstein IJ. Phytohemagglutinins: a new class of metalloprotein: Isolation, purification and some
361 properties of the lectin from *Phaseolus lunatus*. FEBS Letter 1970; 9(4):197-201.

- 362 36. e Lacerda RR, do Nascimento ES, deLacerda JTJG, Pinto LS, Rizzi C, Bezerra MM, et al. Lectin from seeds of a
363 Brazilian lima bean variety (*Phaseolus lunatus* L. Var. cascavel) presents antioxidant, antitumor and
364 gastroprotective activities. *Int. J. Biol. Macromolec.* 2017; 95:1072-1081. doi.org/10.1016/j.ijbiomac.2016.10.097
- 365 37. Shanmugavel S, Velayutham V, Kamalanathan T, Periasamy M, Munusamy A, Sundaram J (2016). Isolation and
366 analysis of mannose/trehalose/maltose specific lectin from jack bean with anti-bruchid activity. *Int. J. Biol.*
367 *Macromolec.* 2016; 91:1-14. DOI:10.1016/j.ijbiomac.2016.05.093
- 368 38. Lis H and Sharon N. Lectins as molecules and as tools. *Ann. Rev. Biochem.* 1986; 55: 35-67.
369 DOI:10.1146/annurev.bi.55.070186.000343
- 370 39. Singh AP, Saxena KD. Biological Activity of Purified *Momordica charantia* Lectin. *Chem. Sci. Trans.* 2013; 2:258-
371 262. DOI: 10.7598/cst2013.277
- 372 40. Benevides RG, Ganne G, Simões RC, Schubert V, Niemietz M, Unverzagt C, et al. A lectin from *Platypodium*
373 *elegans* with unusual specificity and affinity for asymmetric complex N-glycans. *J. Biol. Sci.* 2012; 287(31):26352-
374 26363. DOI:10.1074/jbc.M112.375816
- 375 41. Wong JH, Chan HY, Ng TB. A mannose/glucose-specific lectin from Chinese evergreen Chinkapin (*Castanopsis*
376 *chinensis*). *Biochim. Biophys. Acta.* 2008; 1780(9): 1017-1022. DOI:10.1016/j.bbagen.2008.05.007
- 377 42. Santiago MQ, Leitão CC, Pereira-Junior FN, Pinto VR, Osterne JS Lossio CF, et al. Purification, characterization
378 and partial sequence of a pro-inflammatory lectin from seeds of *Canavalia oxyphylla* Standl. & L. O. Williams. *J.*
379 *Molec. Recogn.* 2014; 27(3):117-123. DOI: 10.1002/jmr.2340
- 380 43. Silva MCC, Santana LA, Mentele R, Ferreira RS, de Miranda A, Silva-Lucca RA, et al. Purification, primary structure
381 and potential functions of a novel lectin from *Bauhinia forficata* seeds. *Process Biochem.* 2012; 47(7):1049-1059.
382 doi.org/10.1016/j.procbio.2012.03.008
- 383 44. Carvalho AS, da Silva MV, Gomes FS, Paiva PMG, Malafaia CB da Silva TD, et al. Purification, characterization
384 and antibacterial potential of a lectin isolated from *Apuleia leiocarpa* seeds. *Int. J. Biol. Macromol.* 2015; 75:402-
385 408. doi.org/10.1016/j.ijbiomac.2015.02.001
- 386 45. Sharma A, Ng TB, Wong JH, Lin P. Purification and characterization of a lectin from *Phaseolus vulgaris* cv.
387 (Anasazi Beans). *J. Biomed. Biotech.* 2009; Volume 2009 Article ID 929568. doi.org/10.1155/2009/929568
- 388 46. Pan WL, Ng TB. A dimeric *Phaseolus coccineus* lectin with anti-oxidative, anti-proliferative and cytokine-inducing
389 activities. *Int. J. Biol. Macromol.* 2015; 81:960-966. DOI:10.1016/j.ijbiomac.2015.09.034


- 390 47. Carrasco-Castilla J, Hernandez-Álvarez AJ, Jiménez-Martínez C, Jacinto-Hernández C, Alaiz M *et al.*, Dávila-Ortiz
391 G. Antioxidant and metal chelating activities of *Phaseolus vulgaris* L. var. Jamapa protein isolates, phaseolin and
392 lectin hydrolysates. Food Chem. 2012; 131(4):1157-1164. doi.org/10.1016/j.foodchem.2011.09.084
- 393 48. Boonmee A, Srisomsap C, Karnchanat A, Sangvanich P. An antioxidant protein in *Curcuma comosa* Roxb.
394 rhizomes. Food Chem. 2011; 124: 476-480. doi.org/10.1016/j.foodchem.2010.06.057
- 395 49. Manukumar HM, Madhu CS. Comparative evaluation of fractional efficiency on antioxidant activity of red gram
396 (*Cajanus cajan*) seed coat crude protein extracts. Int. J. Recent Sci. Res. 2003; 4(9), 1395-1399.
- 397 50. Olodude SO, Odekanyin OO, Fatokun AB. Evaluation of antioxidant and carbohydrate hydrolysing enzymes
398 inhibitory activities of *Trilepisium madagascariense* Ficalho seeds hemagglutinin. IOSR J. Pharm. Biol. Sci.2017;
399 12(6):29-41. DOI: 10.9790/3008-1206042941
- 400 51. Elias RJ, Kellerby SS, Decker EA. Antioxidant activity of proteins and peptides. Crit. Rev. Food Sci. Nutr. 2008;
401 48(5): 430-441. DOI:10.1080/10408390701425615
- 402 52. Coelho LCBB, Silva PMS, Oliveira WF, Moura MC, Pontual EV, Gomes FS, et al. Lectin as antimicrobial agents. J.
403 Appl. Microbiol. 2018; 125(5):1238-1252. DOI: 10.1111/jam.14055
- 404 53. Dias RO, Machado LS, Migliolo L, Franco OL. Insights into animal and plant lectins with antimicrobial activities.
405 Molecules 2015; 20(1): 519-541. DOI:10.3390/molecules20010519
- 406 54. Mishra RP, Ganaie AA, Allaie AH. Isolation and purification of a galactose-specific lectin from seeds of *Bauhinia*
407 *variegata* and evaluation of its antimicrobial potential. Int. J. Pharm. Sci. Res. 2016; 7(2): 804-809.
408 DOI: 0.13040/IJPSR.0975-8232.7(2).804-09
- 409 55. Nawrot R, Barylski J, Nowicki G, Broniarczyk J, Buchwald W, Goździcka-Józefiak A. Plant antimicrobial peptides.
410 Folia Microbiology 2014; 59: 181-196. DOI: 10.1007/s12223-013-0280-4
- 411 56. Salas CE, Badillo-Corona JA, Ramirez-Sotelo G, Oliver-Savador C. (2015). Biologically active and antimicrobial
412 peptides from plants. Biomed. Res. Int. 2015; Volume 2015, Article ID 102129. DOI:10.1155/2015/102129
- 413 
- 414
- 415

Table 1: Summary of purification procedure for *Pterocarpus osun* seeds lectin (POSL)

Fractions	Volume (ml)	Total protein (mg)	Total Activity (HU)	Specific Activity (HU/mg)	Fold Purification
Crude protein extract	50	394.0	1024	2.6	1.0
70% Ammonium Sulphate Precipitate Dialysate	18	77.4	1024	13.23	5.1
Gel Filtration Sephadex G-100	6	12.9	2048	158.8	61.1

Table 2: Hemagglutinating activity of PBS extract of *P. osun* against human and animal erythrocytes

Erythrocyte		Non-trypsinized	Trypsinized
Human	A	2 ⁰	2 ⁵
	B	2 ⁰	2 ⁵
	O	2 ⁰	2 ⁴
Rabbit		2 ¹¹	2 ¹⁵
Rat		2 ⁶	ND

ND Not determined.

455
456
457
458

459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485

UNDER PEER REVIEW

Table 3: Inhibition of hemagglutinating activity of *P. osun* seed lectin by different sugars.

Sugars	Hemagglutinating Titre	Minimum Inhibition Concentration (mM)
Arabinose	2 ¹¹	ND
Xylose	2 ⁹	ND
Glucose	2 ³	0.913 ± 0.345
Galactose	2 ⁹	ND
Mannose	2 ⁰	1.824 ± 0.689
Sorbose	2 ¹⁰	ND
Maltose	2 ⁰	0.260 ± 0.065
Sucrose	2 ⁵	ND
Lactose	2 ¹¹	ND
Mannosamine	2 ⁷	ND
Glucosamine HCl	2 ⁵	ND
2-deoxy-D-glucose	2 ²	3.646 ± 1.378
N-acetyl-D-glucosamine	2 ³	ND
α-D-methyl glucopyranoside	2 ²	1.043 ± 0.261
Mannitol	2 ¹¹	ND
Dulcitol	2 ⁹	ND
α-methyl mannoside	2 ⁰	0.456 ± 0.173
Control	2 ¹¹	ND

Minimum inhibition concentration is the minimum concentration of sugar that inhibits 50% of hemagglutinating activity. Data for minimum inhibition concentration are expressed as mean ± SEM of triplicate determination

ND - Not determined




Figure 1: Gel filtration chromatogram of ammonium sulphate dialysate of crude extract of *P. osun* seeds on Sephadex G-100 column.

The column (2.5 x 40 cm) packed with Sephadex G-100 was equilibrated with 25mM phosphate buffered saline (PBS) pH 7.2 containing 10mM sodium chloride (NaCl). 5 ml of ammonium sulphate precipitate dialysate (4.3 mg) was layered on the column and the lectin was eluted with the same buffer at a flow rate of 15 ml/hr and fractions of 5 ml were collected.

Legend: — Pooled fractions; Hemagglutinating activity; ■■■■ OD₂₈₀;
GO - Protein Peaks




Figure 2: (A) Effect of temperature on *P. osunseeds* lectin (B) Thermostability of *P. osunseeds* lectin.

Lectin samples were incubated at different temperatures (30 - 90 $^{\circ}\text{C}$) for 60 min. Aliquots of the lectin was taken at every 15 min interval and then rapidly cooled in ice and assayed for agglutinating activity. The control was agglutinating activity of lectin sample kept at 20 $^{\circ}\text{C}$.





Figure 3: Effect of pH on the hemagglutinating activity of *P. osunseeds* lectin.

Lectin samples were incubated in the following buffers at different pH values; 0.2 M citrate buffer, pH 2.0 – 5.0; 0.2 M Tris-HCl buffer, pH 6.0 – 8.0; and 0.2 M glycine-NaOH buffer, pH 9.0 – 13.0. After 1 hour, the hemagglutination activity of the lectin was determined. The control values were the agglutination titre of the lectin in PBS, pH 7.2.

531
532
533
534
535
536
537
538
539
540

UNDER PEER REVIEW

541 A




542

543

544

B



545

546

547

548

Figure 4: Antioxidant activity of POSL (A) DPPH radical scavenging activity of *P. osunseeds* lectin (B) Lipid peroxidation inhibition. Data are expressed as mean \pm SEM. (n=3)