

ALLOMETRIC MODELS FOR ESTIMATING SITE INDEX OF TEAK (*Tectona grandis* Linn F.) IN KANYA FOREST PLANTATION, KEBBI STATE, NIGERIA

ABSTRACT

This study aims to develop site index for Teak (*Tectona grandis*) in Kanya Forest Plantation, Nigeria. Site index is defined as the total height of the dominant or co-dominant trees at an arbitrary index age, it is a method used for quantifying site quality for pure even-aged stands which is essential in growth and yield modelling. The data used in this study were obtained from six different age classes. Five sample plots each were selected across all age classes in which a total of 712 trees were measured, variables measured include total height, diameter at the base, middle, top, and diameter at the breast height were taken from 30 temporary sampled plots of 25x25m approximately from the centre, 180 dominant trees were selected from 712 trees. Basal area and volume of sampled trees were computed. Yield values obtained from the dominant trees are (B=249.312m³/ha, D=196.128m³/ha, F=134.976m³/ha, C=119.328m³/ha, E=100.320m³/ ha and A=86.976m³/ha). The results showed that B was the best and A was the poorest. Seventeen models were generated and paired sampled t-test was used for model validation, comparing the actual and predicted height. Two out of 17 were rejected (significant P<0.05). The first model $Hd=12075.346-354.809(Age)+3.448(Age)^2-135193.126(1/Age)$ is the recommended height estimation of Teak in Kanya Forest plantation for its best performance.

Keywords: Site; Site Index; Site Quality; Dominant Trees; Teak.

1. INTRODUCTION

Site index is defined as the average total height of dominant and co-dominant trees (site trees) at a specified reference or base age, which is commonly selected to lie close to the rotation age [1]. The top height is the arithmetic mean height of the 100 trees ha⁻¹ with the greatest diameters [2]. However, the most common objective of site index is to determine the height development pattern that a stand is expected to follow throughout rotation [1]. It is little affected by varying densities and species composition, relatively stable under varying thinning intensities and is strongly correlated with volume.

Site index is a quantitative measure of site quality, and it is generally a reflection of the potential timber productivity of a stand of trees [3]. Dominant and co-dominant trees are used to describe

34 site index because they are assumed to have grown freely throughout their life; thus, the growth
35 of these trees is somewhat independent of other vegetation. A base age is usually used as a
36 reference so that stands of different site quality can be compared. These characteristics of
37 plantation forest - uniformity of crop, the intensity of production, high density, fast growth rate
38 and high productivity have raised concerns that many of the sites on which the plantations are
39 established may be incapable of sustaining their productivity [4]. Site quality assessment is the
40 evaluation of the innate productive capacity of an area of forest land for one or more tree species.
41 Site quality assessment is very important in forest management because a site could support one
42 species excellently while supporting another species poorly.

43 According to 2000-2005 Global Forest Resources Assessment of the Food and Agricultural
44 Organization of the United Nation (FAO), Nigeria has the world's highest annual deforestation
45 rate of primary forest at 55.7%. The country is one of the largest losers of annual natural forest in
46 Africa at 11.1%. Nigeria's annual deforestation rate of natural forest is the highest in the world
47 and put it in the pace to lose virtually all its primary forest within a few years. There is a growing
48 concern about the uncontrolled exploitation of Nigeria's forest resources in accordance to the
49 recent observations that deforestation poses a great risk to sustainable land use and the wellbeing
50 of the people [5].

51 For more than 100 years, site index has remained the world's most widely used measure of site
52 productivity, many decisions in forestry rely on estimates of the land's inherent ability to grow
53 trees and yield timber. These site productivity estimates serve as a baseline for land-use
54 decisions, land appraisals, silvicultural investment analyses, and growth and yield predictions.
55 The importance of site quality assessment remains imperative in quantitative forestry, simply for
56 its potential and possibility of determining the productive capacity of the plantations area for

57 sustainable management [6]. The main objective of this research is to develop site index for Teak
58 (*Tectona grandis* Linn F.) in Kanya Forest Plantation

59 **2. MATERIALS AND METHODS**

60 **2.1 The Study Area**

61 The study was conducted in Kanya Forest Plantation in Danko Wasagu Local Government,
62 Kebbi State, **Nigeria**. It is located on Latitude 11.339⁰N to 11.348⁰ and Longitude 5.606⁰E to
63 5.641⁰E, occupying about 4,208km². It is bordered in the South by Sakaba Local Government, in
64 the West by Zuru Local Government both in Kebbi State and in the North by Bukkuyum Local
65 Government Area of Zamfara State. Danko Wasagu has an estimated population of about
66 265,271 people [7]. The vegetation falls under Northern Guinea Savannah. The topography is
67 said to be flat or low land with fertile soil covered by sandy soils, sometimes coarse in texture
68 with Madama and alluvial plain suitable for agricultural activities. The weather is marked by a
69 single rainy season and long dry season; the average rainfall is 720mm, the rainy season is about
70 four to five months, the mean temperature ranges from 31⁰C and 38⁰C. From November to
71 February cold weather is usually experienced due to the dry harmattan wind and from March to
72 May, the weather is generally hot and wet as in the tropics [8]

73 **2.2 Sampling Procedure**

74 The area was stratified into different age classes based on the years of establishment (1979,
75 1980, 1981, 1982, 1983, and 1989) on which five temporary sample plots of 25 x 25m
76 (0.0625ha) were marked at random from each age block close to the centre. Measurements were
77 taken on all trees within the selected plots. Stand age was obtained from plantation records.

78 **2.3 Data Collection**

79 The data obtained include: Counting and recording of individual trees per plot, Measuring the
80 total height of six dominant trees in all selected plots using Haga Altimeter (this represented the
81 100 largest trees per ha), Diameter at breast height (DBH) of all individual trees was measured at
82 1.3m above ground level. The flexible measuring tape was used to determine the circumference
83 of the boles, Diameters at three different points (Base, middle, Top) were determined with the
84 aid of Spiegel Relascope.

85 **2.4 Development of Site Index Equations**

86 After due consideration of the rotation age and the age of culmination of mean annual increment
87 as recommended by various researchers [9-11] 30 years was adopted as the appropriate base age
88 for the determination of site index of Teak plantations in the study area. Among the various
89 techniques for developing site index equations, the proportional or guide curve method was
90 adopted for this study because the data were obtained from temporary sample plots, thus
91 permitting only the use of this method [1,12]. Various linear and non-linear equations commonly
92 used for site index studies in forestry were selected from forestry literature
93 [13,14,15,16,11,17,18,19]. These equations or models followed the order as follows.

94
$$Hd = a_0 + a_1 (\text{Age}) + a_2 (\text{age})^2 + e_i \quad (4)$$

95
$$Hd = a_3 + a_4 (\text{Age}) + a_5 (1/\text{Age}) + e_i \quad (5)$$

96
$$Hd = a_6 + a_7 \ln (\text{Age}) + e_i \quad (6)$$

97 **2.5 Model Selection and Validation**

98 Different criteria for choosing the best model are available; the highest coefficient of
99 determination (R^2) and the lowest root mean square error (RMSE) were considered appropriate
100 criteria in selecting the best model. Model validation was achieved by dividing the data into two
101 sets; (75%) of the data to calibrate the models and the other set (25%) to validate the models,

102 testing for the significant differences in mean predicted and observed values of the dependent
 103 variables in all cases was achieved using paired sample T-test.

104 3. RESULTS

105 The summaries of growth and yield characteristics of 180 sampled dominant trees are presented
 106 in Tables 1 and 2. Mean, minimum and maximum values of Dbh, height, BA and volume are
 107 recorded for all the age series. The standard error of the mean was also attached to all the mean
 108 values to see the variability distribution of the sampled data from the population.

109 **Table 1: Summary Statistics of Dominant Trees (Sampled Trees)**

| Age (years) | Plots | Trees | Dbh (cm) | | | Height(m) | | |
|----------------|-------|-------|----------|-------|------------|-----------|-------|------------|
| | | | Min | Max | Mean* | Min | Max | Mean* |
| 38 | 5 | 6 | 12.51 | 36.98 | 23.77±0.29 | 9.85 | 15.25 | 15.61±0.44 |
| 37 | 5 | 6 | 20.53 | 27.05 | 25.10±0.75 | 11.30 | 19.60 | 15.19±0.51 |
| 36 | 5 | 6 | 19.26 | 37.91 | 26.62±0.93 | 10.70 | 20.00 | 15.58±0.42 |
| 35 | 5 | 6 | 16.23 | 37.91 | 30.07±1.39 | 11.55 | 19.60 | 22.61±0.46 |
| 34 | 5 | 6 | 19.89 | 48.09 | 24.91±0.89 | 18.80 | 28.25 | 15.07±0.39 |
| 28 | 5 | 6 | 16.87 | 39.15 | 25.59±0.41 | 12.90 | 19.80 | 16.06±0.29 |

110 *Mean± standard error

111 **Table 2: Summary of yield characteristics of Dominant Trees (Sites Trees) *Tectona grandis*.**

| AC | P | Trees | Basal Area (m ²) | | | Volume (m ³) | | | | |
|----|---|-------|------------------------------|------|-----------|--------------------------|-------|-------|-------------------|---------|
| | | | Min | Max | Mean | Min | Max | Mean | Mean volume/ha | |
| A | 5 | 6 | 0.01 | 0.11 | 0.04±0.01 | 4.29 | 0.240 | 0.980 | 0.906±0.04 | 86.976 |
| B | 5 | 6 | 0.03 | 0.06 | 0.50±0.03 | 8.03 | 0.410 | 1.310 | 2.597±0.20 | 249.312 |
| C | 5 | 6 | 0.29 | 1.11 | 0.11±0.04 | 10.34 | 1.260 | 5.630 | 1.243±0.18 | 119.328 |
| D | 5 | 6 | 0.02 | 1.11 | 0.08±0.01 | 7.20 | 0.480 | 5.470 | 2.043±0.18 | 196.128 |
| E | 5 | 6 | 0.03 | 0.18 | 0.05±0.01 | 4.70 | 1.150 | 5.300 | 1.045±0.08 | 100.320 |
| F | 5 | 6 | 0.02 | 0.12 | 0.14±0.02 | 13.14 | 0.580 | 2.290 | 1.406±0.08 | 134.976 |

112 *Mean± standard error

113
 114
 115
 116
 117

118 **3.1 Site Index Parameter Models.**

119 Different model forms/structures were considered in developing site index equations and
 120 estimated regression parameters are presented in Table 3.

Table 3: General models

| | Model Expression | Estimated parameters | | | |
|----|--|----------------------|------------|----------|----------|
| | | a | b | c | d |
| 1 | $Hd = a + b (\text{Age})$ | 23.84 | -0.225 | - | - |
| 2 | $Hd = a + b (\text{Age})^2$ | 20.821 | -0.004 | - | - |
| 3 | $Hd = a + b \ln (\text{Age})$ | 38.441 | -6.311 | - | - |
| 4 | $Hd = a + b \ln (\text{Age})^2$ | 27.990 | -0.948 | - | - |
| 5 | $Hd = a + b (\text{Age}) + c (\text{Age})^2$ | -238.092 | 15.996 | 0.248 | - |
| 6 | $Hd = a + b (\text{Age}) + c (1/\text{Age})$ | 597.471 | -9.017 | 9225.302 | - |
| 7 | $Hd = a + b (1/\text{Age}) + c (\text{Age})^2$ | 296.410 | -5903.004 | -0.089 | - |
| 8 | $Hd = a + b (1/\text{Age}) + c \ln (\text{Age})$ | 2716.098 | -19260.413 | -604.20 | - |
| 9 | $Hd = a + b (\text{Age})^2 + c \ln (\text{Age})$ | -772.306 | -0.129 | 266.731 | - |
| 10 | $Hd = a + b \ln (\text{Age}) + c \ln (\text{Age})^2$ | -3452.704 | 2004.657 | -289.249 | - |
| 11 | $Hd = a + b (\text{Age}) + c (1/\text{Age}) + d (\text{Age})^2$ | -238'183 | 15.999 | 10.00 | -0.248 |
| 12 | $Hd = a + b (\text{Age}) + c(1/\text{Age}) + d \ln (\text{Age})$ | -1349.746 | -17.299 | 15.000 | 555.078 |
| 13 | $Hd = a + b (\text{Age}) + c \ln (\text{Age}) + (\text{Age})^2$ | -545.676 | -23.907 | 15.000 | 110.529 |
| 14 | $Hd = a + b (\text{Age}) + c \ln (\text{Age})^2$ | -544.228 | -23.919 | 110.729 | - |
| 15 | $Hd = a + b (1/\text{Age}) + c \ln (\text{Age})^2$ | 1287.809 | -14801.435 | -66.976 | - |
| 16 | $Hd = a + b (\text{Age}) + c \ln (\text{Age})^2$ | -3427.703 | 1990.238 | -287.173 | - |
| 17 | $Hd = a + b (\text{Age}) + c (\text{Age})^2 + d (1/\text{Age})$ | 12075.346 | -354.809 | 3.448 | 1351.126 |

121 Hd = dominant height, a, b, c, d = Regression coefficients, Age = Age of the plantation

122 **3.2 Model Evaluation and Validation**

123 Fifteen equations were ranked based on some statistics generated in the course of modelling.

124 Models with the highest coefficient of determination (R^2) and Smaller Root mean square error

125 (RMSE) were considered as the better fit models and were subsequently ranked higher than those

126 with lower R^2 and higher RMSE values (Table 4). The selected candidate models were equations

127 1,2, and 3 based on ranking with equation 1 having the highest R^2 and lowest RMSE of 0.760

128 and 1.384. Furthermore, plots of residual values of dominant height for the selected models are

129 shown in Figures 1, 2, and 3. The selected models have their scatter plot normally distributed


130 having perfect prediction of the estimated dominant height.

131


Table 4: General Equations Developed for Site Index Estimation

| Developed Equations | R ² | RMSE | Ranking |
|--|----------------|-------|---------|
| Hd = 12075.346 - 354.809 (Age) + 3.448 (Age) ² - 135193.126 (1/Age) | 0.760 | 2.384 | 1 |
| Hd= 2716.098 - 19260.413 (1/Age) - 604.020ln (Age) | 0.634 | 3.592 | 2 |
| Hd = 1287.809 - 1401.435 (1/Age) - 66.976ln (Age) ² | 0.632 | 3.607 | 3 |
| Hd= - 3452.704 + 2004.657ln (Age) - 289.249ln (Age) ² | 0.627 | 3.657 | 4 |
| Hd = - 597.471 - 9.017 (Age) - 9225.302 (1/Age) | 0.627 | 3.660 | 5 |
| Hd = - 3427.703 + 1990.238ln (Age) - 289.173ln (Age) ² | 0.626 | 3.695 | 6 |
| Hd = 296.410 - 5903.004 (1/Age) - 0.089 (Age) ² | 0.620 | 3.730 | 7 |
| Hd = - 1349.746 - 17.299 (Age) + 15.00 (1/Age) + 555.078ln (Age) ² | 0.620 | 3.745 | 8 |
| Hd = - 772.306 - 0.129 (Age) ² + 266.731ln (Age) | 0.613 | 3.794 | 9 |
| Hd = - 238.092 + 15.996 (Age) - 0.248 (Age) ² | 0.607 | 3.860 | 10 |
| Hd = - 238.183 + 15.999 (Age) + 10.00 (1/Age) - 0.248 (Age) ² | 0.607 | 3.882 | 11 |
| Hd = 20.821 - 0.004 (Age) ² | 0.071 | 9.055 | 12 |
| Hd = 23.884 - 0.225 (Age) | 0.055 | 9.216 | 13 |
| Hd= 27.990 - 0.498ln (Age) ² | 0.045 | 9.315 | 14 |
| Hd = 38.441 - 6.311ln (Age) | 0.041 | 9.352 | 15 |


132 Hd = dominant height (m) Age = Age of the plantation, R² = coefficient of determination, RMSE = Root mean square error.



133 **Figure 1: Residual plots for the dominant height of *Tectona grandis* using first equation**



135 **Figure 2: Residual plot for the dominant height of *Tectona grandis* using second equation**



137
138 **Figure 3: Residual Plot for Dominant Height of *Tectona grandis* using third Equation**

139 To validate the selected models, paired-sample T-test was carried out comparing heights of
140 dominant trees measured from the field and the estimated values from the equations generated
141 (Table 5). Pairs that showed non-significant difference were considered as valid models for
142 application, while pairs that yielded significant differences were rejected ($p < 0.05$).

Table 5: Results of Model Validation for Site Index Estimation

| Paired Samples | Mean difference | T-Value | P-Value | Decision |
|----------------|-----------------|----------|----------|----------|
| Hd(m) v Eqn 1 | 0.00789 | 0.028 | 0.978 ns | Accepted |
| Hd(m) v Eqn 2 | 0.12022 | 0.425 | 0.671 ns | Accepted |
| Hd(m) v Eqn 3 | -0.00193 | -0.007 | 0.995 ns | Accepted |
| Hd(m) v Eqn 4 | -0.00203 | -0.007 | 0.994 ns | Accepted |
| Hd(m) v Eqn 5 | 0.31456 | 1.391 | 0.166 ns | Accepted |
| Hd(m) v Eqn 6 | 0.00181 | 0.008 | 0.994 ns | Accepted |
| Hd(m) v Eqn 7 | 0.00033 | 0.001 | 0.999 ns | Accepted |
| Hd(m) v Eqn 8 | 0.29726 | 1.319 | 0.189 ns | Accepted |
| Hd(m) v Eqn 9 | 0.00039 | 0.002 | 0.999 ns | Accepted |
| Hd(m) v Eqn 10 | -0.42848 | -1.908 | 0.058 ns | Accepted |
| Hd(m) v Eqn 11 | 0.01019 | 0.045 | 0.96 ns | Accepted |
| Hd(m) v Eqn 12 | -3.59844 | -16.004 | 0.00 sig | Rejected |
| Hd(m) v Eqn 13 | -49.59120 | -199.945 | 0.00 sig | Rejected |
| Hd(m) v Eqn 14 | 0.01658 | 0.074 | 0.941 ns | Accepted |
| Hd(m) v Eqn 15 | -0.00152 | -0.007 | 0.995 ns | Accepted |
| Hd(m) v Eqn 16 | 0.00317 | 0.140 | 0.989 ns | Accepted |
| Hd(m) v Eqn 17 | -0.30048 | -1.454 | 0.145 ns | Accepted |

143 Hd = Dominant height (m), Eqn = Equations, Ns = No significant different, Sig.= Significant difference.

144 4. DISCUSSION

145 4.1 Site Index Models

146 The parameter and statistics of regression models 1-17 generated are being presented. Parameters
147 (a, b, c, d,) varied with the model. The selected height-age models are presented according to
148 criteria for selecting the most suitable prediction model, model 17 (Ranked 1st) performed better
149 than all for height-age prediction in Kanya Forest Reserve having highest R^2 and the lowest
150 RMSE of 0.760 and 2.384. Although, the R^2 value was higher than that reported by [20] in
151 height-diameter models and slightly lower R^2 in volume-Dbh models $R^2 = 0.51$ and $R^2 = 0.85$
152 respectively. The result implies greater accuracy for dominant height prediction due to higher
153 modelling efficiency. [18] carried out a similar study on Site Index Equation for *Pinus caribaea*
154 Plantation in Erosion Prone Enugu Ngwo, Nigeria having R^2_{adj} 0.911 and RMSE 0.023
155 respectively. [21] Site Quality Assessment for *Tectona grandis* Linn.F Plantations in Gambari
156 Forest Reserve, Nigeria reported R^2 value of 0.95 and SEE value of 0.0835 this shows higher
157 accuracy than the best model in this research. A study by [22] on Site Quality Assessment of
158 Degraded *Quercus fraianetto* stand in central Greece obtained slightly higher R^2 values (0.799,
159 0.788, and 0.700) for the first two equations and the third equation reporting lower R^2 value, two
160 of the equations have higher accuracy than the best equation in this study and the third equation
161 having low accuracy in predicting dominant height than what was presented in this study. Paired
162 sampled t-test was used to validate the models by comparing the height estimated by the
163 equations and actual height measured. The validation showed mean differences, T-value, P-value
164 and also decisions on the accepted and rejected models. All equations with higher P-values
165 ($P > 0.05$) showed no significant difference between the actual measured and estimated dominant
166 height and are accepted, while those with P-value ($P < 0.05$) showing significant differences were

167 automatically rejected. [20] also used paired sampled t-test to validate the models comparing
168 actually measured and estimated volume of the stand recording three equations that were
169 significantly different ($p < 0.05$).

170 **5. Conclusion and Recommendation**

171 From the results of the study, it can be concluded that the indexed age of 30 was used to
172 determine the site index of *Tectona grandis* in Kanya Forest Plantation, Nigeria. Similarly, the
173 site index equation developed was appropriate for the determination of the site quality of the
174 *Tectona grandis* plantation and thus can be relevant in assessing the productivity of the Teak
175 stand. This will enhance the proper management of the stand for sustainable timber yield
176 production.

177 It is therefore recommended that Equation 1. $Hd = 12075.346 - 354.809 (Age) + 3.448 (Age)^2 -$
178 $135193.126 (1/Age)$, should be used for site index estimation of teak plantation in Kanya Forest
179 Plantation for its best performance and simple structure

180 **REFERENCES**

- 181 1. Clutter JL, Fortson JC, Pienaar LV, Brister GH, Bailey RL. *Timber Management: A*
182 *Quantitative Approach*; John Wiley & Sons.: New York, NY, USA. 1983;30–62.
- 183 2. Hagglund B. Evaluation of forest site productivity. *Forest Abstracts*. 1981; 42(11): 515–
184 527.
- 185 3. Hanson EJ, Azuma DI, Hiserote BA. Site Index equations and mean annual increment
186 equations for Pacific Northwest Research Station Forest Inventory and Analysis
187 Inventories, 1985 2001. USDA Forest Service Research Note PNW-RN-533.2002
- 188 4. Onyekwelu JC, Mosandl R. and Stimm B. Productivity, site evaluation and state of
189 nutrition of *Gmelina arborea* plantations in Oluwa and Omo forest reserves, Nigeria.
190 *Forest Ecology and Management*, 2006; 229: 214-227.
- 191 5. Schulte-Bisping H, Bredemeier M, Beese F. Global availability of wood and energy
192 supply from fuel wood and charcoal. *Ambio*, 1999; 28(7): 592–594.
- 193 6. Skovsgaard JP, Vanclay JK. Forest site productivity: a review of the evolution of
194 dendrometric concepts for even-aged stands. *Forestry Advance Access* published
195 November 22, 2007.
- 196 7. **National Population Commission. Provisional Census Figure. Abuja Nigeria. 2006; 1-3.**

- 197 8. Girma SA. Agro-climatology of Millet Production in Desert Fringe Zone of Nigeria, A
198 Case Study of Kebbi State. M.Sc. dissertation.; Federal University of Technology Minna,
199 Niger state.2008; 1-97.
- 200 9. Curtis RO, Demars DJ, Herman FR. Which dependent variable in site index-height-age
201 regression? Forest Science,1974; 20: 74–87.
- 202 10. Trousdell KB, Beck DE, Llyod FT. Site index for loblolly pine in the Atlantic Central
203 Plain of the Carolina and Virginia. USDA Forest Service Research. Paper SE-1974; 115.
- 204 11. Teshome T, Petty JA. Site index equation for Cupressus lusitanica stands in Munessa,
205 Ethiopia. Forest Ecology and Management. 2000; 126: 339–347.
- 206 12. Nanang DM, Nunifum TK. Selecting a functional form for anamorphic site index curve
207 estimation. Forest Ecology and Management. 1999; 118: 211–221.
- 208 13. Carmean WH. Site index curves for upland oaks in the Central States. Forest Science,
209 1972; 18: 109–120.
- 210 14. Malende YH, Temu AB. Site index curves and volume growth of teak (*Tectona grandis*)
211 at Mtibwa, Tanzania. Forest Ecology and Management. 1990; 31: 91–99
- 212 15. Akindele SO. Development of a site index equation for teak plantations in south-western
213 Nigeria. Journal of Tropical Forest Science.1991; 4(2): 162–169.
- 214 16. Onyekwelu JC, Fuwape JA. Site index equation for *Gmelina arborea* pulpwood
215 plantations in Oluwa Forest Reserve, Nigeria. Journal of Tropical Forest Science. 1998;
216 10(3): 337–345.
- 217 17. Onyekwelu JC. Choosing appropriate index age for estimating site index of *Gmelina*
218 *arborea* timber plantations in Oluwa forest reserve, Nigeria. Journal of Food, Agriculture
219 and Environment. 2003; (3&4): 286–290.
- 220 18. Oyebade BA, Popo-ola FS, Aguma S. Site Index Equation for *Pinus Caribaea* Plantation
221 in Erosion Prone Enugu Ngwo, Nigeria. International journal of science and nature. 2012;
222 3(3): 586-591
- 223 19. Kitikidou K, Milios E, Tsirekis E, Pipinis E, Stampoulidis A. Site quality assessment of
224 degraded *Quercus frainetto* stands in central Greece. iForest – Biogeosciences and
225 Forestry: 2014; e1-e6.
- 226 20. Adeyemi AA. Site Quality Assessment and Allometric Models for Tree Species in Orban
227 Forest, Nigeria. Journal of Sustainable forestry. 2016; 36(4): 280-289.
- 228 21. Ige PO, Akinyemi GO. Site Quality Assessment for *Tectona grandis* Linn.f Plantations
229 in Gambari Forest Reserve, Nigeria. Journal of Forestry Research and Management.
230 2015; 12: 58-67.

231

232