

1
2
3
4
5
6
7
8
9
10
11

The A Comparative Study of the Daily Activity Patterns of Dog Faced Baboon (*Papio anubis*) in Captivity at: A Case Study of the Kano University Zoo and Kano Zoological Garden

ABSTRACT

Aim: This study seeks to observed the daily activity patterns of dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Wudil Zoo and Kano Zoological Garden.

Materials and methods: This study of the activities of dog faced baboon (*Papio anubis*) in Kano University of Science and Technology Wudil Zoo and Kano Zoological Garden was carried out daily from between 6:00am to 6:00pm between from December 2016 to January 2017. Digital camera was also attached to cages at the two sites. The observation in the activity patterns of dog face baboon (*Papio anubis*) were recorded in on the recording sheet, observation is done three times a week at 20 minutes intervals.

Results: The findings on activity pattern of dog faced baboon (*Papio anubis*) in captivity shows that the day time activities decrease from morning to evening. Resting activities was 47.5%, movement and feeding were carried out in the morning, followed by afternoon and evening with 33.3% and 19.1% activities respectively. The results from of the activities of dog faced baboon in Kano Zoological Garden, indicated that 42.7% of the activities perform by dog faced baboon in captivity are resting, this is followed by movement which accounted for 34.9% of the activities, while feeding activities account for the least with 22.4%. Similarly, it shows that about 43.2% of the recorded activities carried out by dog faced baboon in Kano University of Science and Technology Wudil, Zoo was Resting, followed by the Movement which constituted with 34.8% of the activities and the feeding activity which accounted for 22%.

Conclusion: Due to the fact that majority of the baboons activities usually take place between ~~m~~Morning and afternoon, it is recommended that visitors [interested in baboons](#) should ~~plan~~pay their visitation to the Zoo pen during that time. [It is also recommended that](#) ~~f~~Feeding and ~~chasing-harassing~~ of ~~a~~Animals by the visitors should be discouraged [in order to ensure consistency in their behaviour.](#)

12

13 | *Keywords: Papio ~~a~~Anubis, feeding, movement, resting, Kano University of Science and*
14 | *Technology Wudil and Kano Zoological Garden*

15

16

17

18

19 | **1. INTRODUCTION**

20

21 | Activity patterns have been studied in several primate taxa including hominoids [1,2]
22 | cercopithecines [3, 4, 5] [and](#) colobines, [6]. Time is limited for most animals [7, 8]. Thus,
23 | animals are faced with the challenge of allocating the limited time to different activities.
24 | According to the optimality theory, “the amount of time that an organism spends engaged in
25 | various activities depends on the cost of the activity relative to the derived benefits in that
26 | organism’s habitat” [9].

27 | The amount of time spent on foraging activities therefore relates to the energy content of the
28 | food relative to the costs of obtaining the food plus the cost of all other activities (resting,
29 | moving or socialising). Thus, specifically, food availability and energy content are critical
30 | determinants of an animals’ daily activity pattern. Therefore, factors that influence the
31 | availability of food have a strong bearing on time allocation profiles in baboons.

32 | Due to the different costs and benefits of specific activities animals have varying time
33 | allocation profiles based on age and sex for certain activities [10]. Furthermore, since these
34 | activities cannot be performed simultaneously some individuals may allocate time between
35 | various behaviours better than others [7, 8]. The costs and benefits of these activities
36 | change with changes in the ecological and social state of the environment as well as the
37 | physiological state of the animal. This gives rise to temporal and spatial variation in
38 | individual activity budgets of the animal. Baboons allocate the greater proportion of their time
39 | to foraging activities [11, 12, 13, 14, 15]. De Hoop and Mkuzi baboon troops spent 69.8 %
40 | and 66.5 % [respectively](#) of their time foraging [respectively](#) [14]. **In a study of [12] that they**

41 spend 69.8 %, 75.2 % and 43 % of their time foraging, respectively. The Lodge troop spent
42 relatively less time foraging than Alto and Hook groups.

43 Weather patterns have both direct and indirect influences on the activity pattern of primates.
44 Rainfall and temperature have pervasive effects on animals [16] and so influence time
45 allocation patterns both temporally and spatially.

46 This study seeks to observed the daily activity patterns of dog faced baboon (*Papio anubis*)
47 in Kano University of Science and Technology Wudil Zoo and Kano Zoological Garden.

48 2. MATERIAL AND METHODS

49

50 2.1 Study area

51 The study was carried out in Kano University of Science and Technology Zoo and Kano
52 Zoological Garden. Wudil is located between the latitude $11^{\circ} 37'N$ and longitude $8^{\circ} 58'E$ has
53 a total area of $362km^2$ and is located within Sudan savannah region of Nigeria. The annual
54 maximum rainfall is between 850mm-870mm with a minimum and maximum temperature of
55 $26^{\circ}c$ - $30^{\circ}c$. The relative humidity of the region is always low and ranges between 40% -
56 51%.

57 2.2 MATERIALS

58 Standardized data collection sheet, stop clock, Recording sheet and Digital camera

59 2.3 DATA COLLECTION

60 Sampling method was used to study the activities of dog faced baboon (*Papio anubis*) in
61 Kano University of Science and Technology Wudil Zoo and Kano Zoological Garden from
62 6:00am to 6:00pm between December to January 2016. The observations in the activity
63 patterns of dog face baboon (*Papio anubis*) are were recorded in standard data the
64 recording sheets, observation is done three times a week at 20 minutes intervals in each of
65 the cages under study. Note: this research is limited to period when the temperature is
66 extremely low (Hammattan period). The activity parameters recorded include: Feeding,
67 Moving, and Resting and are described as follows:

68 **Feeding:** the feeding began when the animal first made contact with any part of food or
69 other food substances, feeding bout terminated when the either moved more than one full
70 stride, even if it was carrying some food material on its hand and mouth or stopped looking
71 at the food material, by this definition, a switch to a new food type in the absence of either of
72 these condition was not for bout to be consider terminated, thus a single feed bout could
73 include more than one food type [17, 9].

74 **Resting:** this includes behavior during which an animal was neither feeding, moving or
75 engaged in other social behavior that include sleeping auto-grooming, looking around etc [9,
76 10].

77 **Moving:** this includes all locomotion activities like walking, running, climbing, jumping and
78 leaping but excluding short movements during feeding and locomotion during social behavior
79 e.g when primates chased one another [9, 10].

80 **Other activities:** other social behavior including all other activities which an animal's
81 attention and behavior where clearly directed toward another individual. These include allo-
82 grooming, mounting, mating, chasing, playing, aggressive or agnostic behaviours [9, 10].

83 All the activities such as resting, movement and feeding are carried out in the morning,
84 afternoon and evening.

85 2.4 DATA ANALYSIS

86 The data collected are was subjected to descriptive statistics which includes frequency
87 distribution and percentage.

88

89 3. RESULTS AND DISCUSSIONS

90 The result of the day time activities of dog faced baboon (*Papio anubis*) in captivity is
91 presented in Table 1. The result shows that the day time activities decrease from morning to
92 evening. Resting was 47.5% of the activities, movement and feeding were carried out in the
93 morning, followed by afternoon and evening with 33.3% and 19.1% activities respectively.

94 **Table 1 Variation in day time activities of dog faced baboon (*Papio anubis*) in**
95 **captivity.**

96

97 DAY TIME	TOTAL NUMBERS OF ACTIVITIES	PERCENTAGE
98 Morning	67	47.52
99 Afternoon	47	33.33
100 Evening	27	19.15
101 Total	141	100.00

102

103 The result of the activities of dog faced baboon in Kano Zoological Garden shown in
104 Tables 2 and 3, indicated that 42.7% of the activities perform by dog faced baboon in
105 captivity are resting. This was followed by movement which accounted for 34.9% of the
106 activities, while feeding activities account for the least with 22.4%. The result of this study is
107 in variance with the finding of [9] who reported 50.00% for feeding and 8.50% for resting for
108 the Kkwano Forest baboons. In his study, Kkwano Forest baboons spent relatively higher
109 proportion of time feeding and lesser proportion of time resting and movement, this is
110 probably due to the level of availability and distribution of food resources at the site compare
111 to captive environment.

112

113

114

115

116 **Table 2: Variation in the activities of dog faced baboon (*Papio anubis*) in Kano**

117 **Zoological garden/day.**

118

119 **NUMBER OF DAYS FREQUENCY (ACTIVITIES) PERCENTAGE (%)**

120

121 **DAY 1**

122 Feeding 32 22.7

123 Moving 50 35.5

124 Resting 59 41.8

125 **DAY 2**

126 Feeding 32 23.0

127 Moving 49 35.2

128 Resting 58 41.7

129 **DAY 3**

130 Feeding 30 21.4

131 Moving 52 37.1

132 Resting 58 41.4

133 **DAY 4**

134 Feeding 31 22.5

135 Moving 46 33.3

136 Resting 61 44.2

137 **DAY 5**

138 Feeding 30 21.3

139 Moving 53 37.6

140 Resting 58 41.1

141 **DAY 6**

142 Feeding 31 22.5

143 Moving 47 34.0

144 Resting 60 43.5

145 **DAY 7**

146 Feeding 32 23.3

147 Moving 43 31.4

148 Resting 62 45.2

149 **Total 974 100**

150

151

152 **Table 3: variation in the activities of dog faced baboon (*Papio anubis*) in Kano**

153 **Zoological garden/week**

154 **ACTIVITIES TOTAL NUMBERS OF ACTIVITIES PERCENTAGE**

155

156 Feeding 218 22.4

157 Movement 340 34.9

158 Resting 416 42.7

160
 161 The result of the dog faced baboon activities is indicated in Table 4 and 5. It shows that
 162 43.2% of the activities carried out by dog faced baboon in Kano University of Science and
 163 Technology Wudil, Zoo is Resting, followed by the Movement with 34.8% of the activities
 164 and the feeding activity account for 22%. This result of the dog faced baboon activities in
 165 captivity indicated in the table above is however, agrees with the finding of [10] who reported
 166 highest resting period than feeding and movement period

167 The Dog faced Baboon activities are higher in day time period. The baboon was observed to
 168 be very active in the morning followed by afternoon and evening. These activities which is
 169 made up of mostly movement and feeding may be due to the presence of visitors in the
 170 morning and afternoon his activity pattern morning, afternoon and evening have been
 171 commonly reported among arboreal species [17]. The daily activities of dog faced baboon
 172 (*Papio anubis*) in Kano University of Science and Technology Zoo and Kano Zoological
 173 Gerden ranged between 6:00am in the morning to 6:00pm in the evening in which most of
 174 the visitors usually pay their visit. However, [17] was of the opinion that adaptive
 175 significances of diurnal variability in primate's activities budget are poorly understood. With
 176 regards to individual activities, resting which include sleeping, looking about etc. was the
 177 most frequent activity carried out by the dog faced baboon in captivity. This may be due to
 178 the confinement in which the baboons were kept. Most of the baboon's time was spent in
 179 sitting postures, standing or playing posture. In this position, the hind limb may be placed in
 180 variety of positions and the fore limbs of the baboon often at rest on the knees or between
 181 hind limb. However, despite the confinement, movement also constitute the substantial
 182 percentage of the dog faced baboon activities in Kano University of Science and Technology
 183 and Kano Zoological Garden. The movement which accounted for 34.9% in Kano Zoological
 184 Garden and 34.8 in the University Zoo include walking, running, climbing, leaping and riding.
 185 This significance percentage might be due to the fact that dog faced baboon are usually
 186 regarded as one of the most entertaining animal within the Zoo.

187 **Meanwhile, of both the animals studied, the Baboon in the Kano Zoological Garden has the**
 188 **slightly higher frequency of activities. This might be also due to more number of visitor.**

189
 190 **Table 4: Variation in the activities of dog faced baboon in Kano University of Science**
 191 **and Technology Wudil, Zoo/day.**

192	NUMBER OF DAYS	FREQUENCY (ACTIVITIES)	PERCENTAGE
193			
194	DAY 1		

195	Feeding	29	21.0
196	Moving	37	26.8
197	Resting	72	52.1
198	DAY 2		
199	Feeding	27	19.8
200	Moving	35	25.7
201	Resting	74	54.4
202	DAY 3		
203	Feeding	30	21.9
204	Moving	42	30.6
205	Resting	65	47.4
206	DAY 4		
207	Feeding	30	22.2
208	Moving	45	33.3
209	Resting	60	44.4
210	DAY 5		
211	Feeding	31	22.3
212	Moving	56	40.3
213	Resting	52	37.4
214	DAY 6		
215	Feeding	32	23.3
216	Moving	60	43.8
217	Resting	45	32.8
218	DAY 7		
219	Feeding	32	23.7
220	Moving	58	43.0
221	Resting	45	33.3
222	Total	957	100

223

224

225 **Table 5: Variation in the activities of dog faced baboon (Papio anubis) in Kano**

226 **University of Science and Technology Zoo/week**

227	ACTIVITIES	FREQUENCY (ACTIVITIES)	PERCENTAGE
228	Feeding	211	22.0
229	Movement	333	34.8
230	Resting	413	43.2
231	Total	957	100

232

233 **CONCLUSION**

234 This study was designed to gather information on the daily activity pattern of Dog faced

235 Baboon in Kano University of Science and Technology and Kano Zoological Garden. From

236 the study, the following conclusions can be made. The dog faced baboons are most active in

237 the morning. Also Resting constitute the most frequent activity of dog faced baboon in

238 captivity. Most of the baboon activities have short duration.

239

240

241 **COMPETING INTERESTS**

242 Authors have declared that no competing interests exist.

243

244

245 **REFERENCES**

246 1. Goodall, J. The chimpanzees of Gombe: patterns of behaviour, Belknap Press,
247 Cambridge, 1986.

248 2. Clutton-Brock, T.H.C. and Harvey P. Species differences in feeding and ranging
249 behavior in primates. In: primate Ecology (Ed. T.H.C. Clutton-Brock), New York:
250 Academic press, 1977. pp. 557-584

251 3. Post, D.G. Activity Patterns of Yellow Baboons (*Papio cynocephalus*) In the
252 Amboseli National Park, Kenya. *Animal Behaviour* 1981, 29: 357-374.

253 4. Van Schaik C.P. Why are diurnal primates living in groups? *Behaviour* 1983,
254 87:120-144.

255 5. Isbell, L.A., Preutz J.D. and Young T.P. Movements of vervet (*Cercopithecus*
256 *aethiops*) and patas monkeys (*Erythrocebus patas*) as estimators of food resource
257 size, density and distribution. *Behaviour Ecology Sociobiology* 1993, 42:123-133.

258 6. Strushaker, T.T. *The red colobus monkey*, University of Chicago Press, Chicago,
259 1975.

260 7. Dunbar, R.I.M. *Primate Social Systems*, Cornell University Press, Ithaca, 1988.

261 8. Dunbar, R. I. M. Time: A Hidden Constraint on the Behavioural Ecology of
262 Baboons, *Behaviour Ecology Sociobiology*, 1992;31: 35-49.

263 9. Gessaman, J.A. Ecological energetics of homeotherms: a view compatible with
264 ecological modelling, Utah University Press *Monograph series*, 1973 20: 1-155.

265 10. Melle, E.M. socio-ecology of Olive baboons thesis Department of wildlife and
266 fisheries management, university of Ibadan, 2004.

267 11. Altmann, J and Altmann, S *Baboon Ecology: African Field Research*. Karger, Basel,
268 1970.

269 12. Amboseli, Kenya, Bronikowski and Altmann . Guarding Constrains Foraging Activity
270 of Male Baboons, *Animal Behaviour*, 1996; 51: 1269–1277.

271 13. Barton, R.A, Whitten A, Strum S.C, Byrne, R. W and Simpson, A.J Habitat and
272 Resource Availability in Baboons, *Animal Behaviour*, 1992; 43: 831-844.

- 273 14. Gaynor, D Foraging and feeding behaviour of chacma baboons in a woodland
274 habitat, PhD thesis, University of Natal, 1994.
- 275 15. Johnson, S.E. & Bock, J. (tradeoffs in skill acquisition and time allocation among
276 juvenile chacma baboons, *Human Nature* Volume, 2004; 15 (1): 45-62.
- 277 16. Bronikowski, A.M and Altmann, J Foraging in variable environment: weather
278 patterns and the behavioural ecology of baboons, *Behaviour Ecology Sociobiology*,
279 1996; 39: 11-25
- 280 17. Post, D.G Feeding Behaviour of Yellow Baboons (*Papio Cynocephalus*) In
281 Amboseli National Park, Kenya, *International Journal of Primatology*, 1982; 3(4): 403-
282 430.
- 283 18. Clutton-Brock, T.H.C. Some aspect of intra-specific variation in feeding and ranging
284 behavior in primates. In: primate Ecology (Ed. By T.H.C. Clutton-Brock), pp. 529-
285 536. New York: Academic press, 1977.
- 286

UNDER PEER REVIEW