

3
4 **Hydric-stress tolerance in cocona (*Solanum***
5 ***sessiliflorum* Dunal)**
6
7
8
9

10
11
12 **ABSTRACT**
13

Aims: The present work aims to assess hydric stress tolerance in cocona (*Solanum sessiliflorum*).

Study design: Four cocona genotypes were planted in completely randomized blocks design with three replicates. Each replicate was irrigated with different water volumes, tantamount to 50, 100 and 150% of evapotranspiration (ET) respectively.

Place and Duration of Study: The present study was developed in National Institute of Amazonian Research at the agricultural experimental station, which is located on Km 14 AM-10 roadway, from September 2013 to April 2014.

Methodology: The fruits were harvested each 15 days by three months. The assessed characters were plant stand, stem diameter, plant height, fruit yield, number of fruits per plant; fruit mass, length, diameter and length/diameter ratio.

Results: Irrigation treatments, both 50 and 150% ET, reduced height plant, fruit mass and length. Other characters were not affected by the hydric stress.

Conclusion: Cocona is tolerant to both hydric stress, being the major hydric stress effect fruit size and mass decreasing. Other studies must be performed to determinate the hydric stress threshold which lead to decrease fruit yield and dead plant.

14
15 *Keywords: abiotic stress; Amazon crop; cubiu; drip irrigation; water management*
16

17 **1. INTRODUCTION**
18

19 Cocona (*Solanum sessiliflorum* Dunal) belongs to Solanaceae family and Lasiocarpa section. This
20 section holds 13 cultivate species distributed from northern Andes region to Amazon. Cocona is
21 distributed in the Amazon region, which includes Peru, Ecuador, Brazil, Colombia and Venezuela [1, 2].
22 This region presents heavy high rainfall (>2500mm). But recently an interest has recently been
23 demonstrated on having this species grown under subtropical conditions with lighter rainfall. In the future,
24 perhaps it will be adapted in a greenhouse system. Thus, hydric tolerance studies in cocona are need to
25 face future cocona cultivation challenges.

26 Cocona is also well adapted to an acid, low nutrient soil and high temperatures [3]. Its fruits look resemble
27 tomato and its plant architecture is like that a large-leafed eggplant. Its fruit tastes like a citric fruit
28 combination. It is used for making ice cream, juice [2], meat dishes, sauce, jelly and desserts [3].

29 Cocona researches have focused on assessing genotypes [3, 4], outcrossing rate studies [3, 5, 6],
30 chemical characterization for food processing industry [7, 8]. However, there is paucity of regarding its
31 physiology; especially on what concern hydric tolerance. In spite of this fact, there are some studies on
32 eggplant hydric tolerance. These studies can be help to understand hydric tolerance in cocona, on
33 account of, both species being phylogenetically related [9].

34 In eggplant, irrigation with 85% of evapotranspiration (ET) had no effect on fruit yield, but 65% and 40%
35 one reduced it by 35 and 46% respectively [10]. Water management can be raise fruit yield and quality of
36 several species [11-14]. Therefore, cocona hydric stress studies can help to manage irrigation of this
37 species, specially, during dry season in the Amazon (June to October).

38 The present paper aim to assess the hydric stress effect on fruit yield by over and under irrigating cocona,
39 150 and 50% ET respectively.

40 41 **2. MATERIAL AND METHODS**

42
43 The experiment was conducted in greenhouse at INPA experimental field "Dr. Alejo von der Pahlen" (02°
44 59'48.2" S and 60° 01' 22.4" W), during September 2013 to April 2014. The mean annual rainfall was
45 2450 mm (mainly from November to June) [15] and mean temperature 27°C. The soil was non-flooded
46 land, red-yellow argisil, sandy texture and pH=6.0. This is a typical Amazon soil, which is poor in organic
47 material; therefore, it was fertilized using 2kg of compost per plant.

48 Cocona genotypes were CUB-10, CUB-11, CUB-12 and CUB-13. These genotypes were originating from
49 Santa Isabel do Rio Negro municipality, Amazonas State (00° 24'50" S; 65° 01' 08" W), from Acariquara,
50 Abianai, Matozinho and Naraná do Enxixi cities, respectively.

51 The genotypes were planted in a completely randomized block design with three replications. Each
52 replication was irrigated with one type of irrigation regime, which were 50, 100 (control) and 150% of
53 evapotranspiration (ET). The fruits were harvested during three months and assessed characters were
54 stand of plant, stem diameter (cm), plant height (cm), fruit yield (t.ha⁻¹), number of fruits per plant, fruit
55 mass (g), length (cm), diameter (cm) and length/diameter ratio.

56 The drip irrigation system combining with evaporation data were used to adjust daily water quantity in
57 each block. We used three type of drip irrigation lines, which had emitters spaced in 10, 20 and 40 cm.
58 The climatic data and evapotranspiration are presented in Table 1. The climatic data were obtained using
59 digital thermohygrometer Incoterm®. The evapotranspiration (ET) was estimated via Ivanov equation:

$$60 \quad ET = 0.006 \times (25 + T)^2 \times (1 - RH/100) \times Kc$$

61 ET= Evapotranspiration (mm.day⁻¹), T=Mean temperature (°C), RH=Relative humid, Kc= Crop coefficient,
62 which has four values, depending of growing stage. This coefficient (Kc) was adapted from eggplant [16].

63

64 **Table 1. Temperature, relative humid, and evapotranspiration per month. Manaus 2013-2014**

Month	Temperature (°C)	Relative humid (%)	Evapotranspiration (mm/day)	Rainfall (mm/day)
2013				
Sep	35.8	77	4.04	0.89
Oct	32.8	52	7.66	4.51
Nov	31.3	51	5.75	12.00
Dec	31.9	46	6.48	4.22
2014				
Jan	31.4	46	5.99	7.60
Feb*	30.2	51	4.86	9.10
Mar*	29.9	57	4.59	13.95
Apr*	31.4	51	5.60	10.86

65 *The fruit harvests were performed during these months.


66
 67 Data were submitted to analysis of variance, and Duncan test ($P < 0.05$) using SAS Software, and
 68 procedure PROC GLM. In addition, it was made quadratic equations to predict characters behavior. The
 69 equation vertex was estimated by $-b/(2a)$, which indicates the equivalent irrigation that maximize fruit
 70 mass, number per plant and yield.

71 To show the relationship among characters and irrigation treatments was make a biplot graphic using
 72 GGEBiplotGUI package in R software (R Core Team). For this purpose, the data were scaled by standard
 73 deviation of each character.

74 75 **3. RESULTS AND DISCUSSION**

76
 77 We had found no previous studies on cocona hydric stress, but there are in other Solanaceae such as
 78 tomato [17], hot pepper [18] and eggplant [19]. In them, the evapotranspiration (ET) method seems
 79 appropriate to measure the effect of hydric stress.

80 Equivalent irrigation to 100% ET indicates that irrigation restores evaporated water. Thus, water quantity
 81 above or below 100% ET would be lead to hydric stress. The irrigation accuracy based on ET was
 82 observed through Biplot analysis ["which won where what" method] (Fig. 1), where 50, 100 and 150% ET
 83 were far apart from each other with all high-valued characters associated with 100% ET. In other words, it
 84 would indicate 100% ET to be optimal to maximize every character expression. Far apart points indicated
 85 the contrasting effect of irrigation treatments on characters. Therefore, this irrigation management
 86 showed to be optimal to assess hydric stress. In addition, these results suggest that eggplant crop
 87 coefficient may be used in Ivanov equation. Probably eggplant and cocona have similar physiology, on
 88 account of both are similar phylogenetically [9].



89

90 **Fig. 1. Biplot graphic shows a relationship among irrigation volume and morphological characters**
 91 **by “which won where what” method.**

92

93 Eleven $t \cdot ha^{-1}$ was the maximum fruit yield, which is very low comparing with other studies. Silva Filho and
 94 Yuyama [4] reported fruit yield from 40 to 100 $t \cdot ha^{-1}$ in Manaus. Low fruit yield may be accounted for by
 95 the fruits having to be harvested for three months, due *Sclerotium rolfsii* infestation. Normally, the
 96 harvesting is performed following four to five months. Nevertheless, the results were sufficient to show the
 97 effect of hydric stress on early yield.

98 The findings showed significant effect of both hydric stresses (50% and 150% ET) mainly on plant height,
 99 fruit mass and length (Table 2). However, it was found no significance difference to stand of plants, stem
 100 diameter, fruit yield, fruit number per plant, fruit diameter and length/diameter ratio (Table 2). Therefore
 101 irrigation of 50% and 150% ET would be utilized without decrease the potential fruit yield.

102

103 **Table 2. Duncan test for irrigation regime of 50, 100 and 150% of evapotranspiration (ET)**
 104 **considering various fruit characters. Manaus 2013-2014.**

Irrigation based on ET (%)	Plant stand	Stem diameter (cm)	Plant height (cm)	Fruit yield ($t \cdot ha^{-1}$)	Fruits Number per plant	Fruit mass (g)	Fruit length [L] (mm)	Fruit diameter [D] (mm)	L/D ratio
50	4.5 a ¹	3.9 a	128.5 b	7.9 a	7.1 a	132.4 ab	62.5 b	59.0 a	1.04 a
100	4.8 a	4.2 a	140.3 a	10.6 a	8.6 a	153.4 a	68.6 a	61.7 a	1.19 a
150	4.0 a	3.9 a	132.8 ab	7.3 a	8.6 a	107.4 b	64.6 ab	60.0 a	1.07 a

105

¹Means with the same letter in the column are not significantly different ($P > 0.05$)

106 Generally, hydric stress led to the decrease of cocona growing and developing. Over and under irrigation,
 107 150 and 50% ET, decreased fruit mass by 30 and 13% respectively. Plant height decreased by 5 and 8%
 108 respectively. Fruit length decreased 8 and 6% respectively. Despite irrigation treatments having
 109 presented no significant differences on the fruit yield, they presented a tendency to lower it by 31 and
 110 25% respectively. These facts would support the former observation of Silva Filho [20], which over
 111 irrigation would decrease fruit yield. Comparatively, these yield decreases are minor than in eggplant [10],
 112 which were 35% for 60% ET. It suggests cocona has more tolerance to hydric stress than it.


113 On the other hand, cocona genotypes showed difference in fruit mass, length and L/D ratio (Table 3).
 114 Indicating there to be genotypic diversity. Therefore, these results concerning hydric stress may be valid
 115 to cocona species.

116 **Table 3. Duncan test for cocona genotypes considering various fruit characters. Manaus 2013-**
 117 **2014**

Genotype	Plant number per plot	Stem diameter (cm)	Plant height (cm)	Fruit yield (t.ha ⁻¹)	Fruits number per plant	Fruit mass (g)	Fruit length [L] (mm)	Fruit diameter [D] (mm)	L/D ratio
CUB-10	4.3 a	4.2 a	130.3 a	8.1 a	7.7 a	157.7 ab	71.3 ab	67.9 a	1.04 ab
CUB-11	4.3 a	3.9 a	137.6 a	10.4 a	10.5 a	164.8 ab	79.5 a	64.3 a	1.23 a
CUB-12	5.0 a	3.8 a	134.4 a	11.0 a	9.2 a	182.4 a	83.1 a	67.3 a	1.23 a
CUB-13	4.0 a	4.0 a	133.1 a	5.4 a	5.1 a	140.6 b	62.0 b	69.2 a	0.90 b

119 [†]Means with the same letter in the column are not significantly different (P> 0.05)

120 Usually, quadratic equations are used to find maximum yield points [21]. Fruit mass and number of fruits
 121 per plant had quadratic behavior (Figure 2).



122


123 **Fig. 2. Quadratic behavior of fruit mass and fruit number per plant for different irrigation volumes**
 124 **in cocona.**

125

126 In the same figure is showed fruit mass is more sensible to hydric stress while fruit numbers per plant is
 127 more stable. The vertex equation shows the high fruit mass and fruit number per plant would be found at

128 91 and 125% ET respectively. In other words, irrigations from 50 up to 91% ET tend to increase both fruit
 129 mass and fruit number. Irrigations from 91 up to 125% ET tend to reduce the fruit mass, but to increase
 130 fruit number per plant. Irrigations from 125 up to 150% ET decrease both characters. This quadratic
 131 behavior was observed in “gigante cocona” [22].

132 On the other hand, when fruit mass was compared with fruit yield (Figure 3) was observed fruit mass is
 133 more sensitive to the hydric stress than fruit yield. Vertex equation showed that around 98% ET led to
 134 high fruit yield.



135


136 **Fig. 3. Quadratic behavior of fruit mass and yield for different irrigation volume in cocona.**

137

138 Since genotypic point view there was genetic variability for fruit mass (140.6 – 182.4 g), fruit length (62.0
 139 – 83.1 mm) and length diameter (L/D) ratio (0.9 – 1.2). CUB-12 showed the highest fruit mass (182.4 g)
 140 with elongated fruits and yield of 11 t ha⁻¹ (Table 3). In contrast, CUB-13 showed the lowest values to fruit
 141 mass (140.6 g) with flat-round fruits and yield of 5.4 t ha⁻¹.

142 Biplot analysis (Figure 4) accounted for the 99% of variation. It indicates the interpretations are highly
 143 reliable. The vectors represent each character and their direction the behavior. Thus, the biplot analysis is
 144 doing by vectorial comparisons. All vector directions were predominantly towards 100% ET. It shows a
 145 positive association between 100% ET with all high values of characters, in other words, this irrigation
 146 level increased all character values. In contrast, 50 and 150% ET negatively affected the character
 147 expression.

148



149

150 **Fig 4. Biplot graphic shows a relationship among irrigation volume and morphological characters**
 151 **(50, 100 and 150% ET represent water volumes used in cocona irrigation based on**
 152 **evapotranspiration)**

153 However, 50% ET would be slightly associated with high plant stand and fruit mass. At the same time,
 154 150% ET would be associated with high fruit number per plant, which is agreeing with maximum
 155 quadratic curve point estimated by vertex formula (125% ET).

156 **4. CONCLUSION**

157

158 Cocona is tolerant to hydric stress both excessive irrigation and its shortage. Its characters more sensible
 159 to this stress were plant height, fruit mass and length.

160

161 Further studies will have to be conducted in order to test more extreme hydric stress, such as 25 and
 162 175% of evapotranspiration, considering different phases of growing: seedling, vegetative and
 163 reproductive phase.

164

165

166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217

COMPETING INTERESTS

Authors have declared there not to be any competing interests.

REFERENCES

1. Schuelter AR, Grunvald AK, Amaral Junior AT, da Luz CL, Luz CL, Gonçalves LM, et al. *In vitro* regeneration of cocona (*Solanum sessiliflorum*, Solanaceae) cultivars for commercial production. *Genetics and Molecular Research*. 2009; 8(3):963-75.
2. Melgarejo TA, Fribourg CE, Russo M. Properties of a tobusvirus that infects cocona (*Solanum sessiliflorum*) in the Peruvian jungle. *Journal of Plant Pathology*. 2003; 85(2):105-10.
3. Salick J. Crop Domestication and the Evolutionary Ecology of Cocona (*Solanum sessiliflorum* Dunal). In: Hecht M, Wallace B, Macintyre R, editors. *Evolutionary Biology*, vol 26. New York: Plenum Press; 1992.
4. Silva Filho DF, Yuyama LKO, Aguiar JPL, Oliveira MC, Martins LHP. Caracterização e avaliação do potencial agrônomo e nutricional de etnovarietades de cubiu (*Solanum sessiliflorum* Dunal) da Amazônia. *Acta Amazônica*, 2005; 35(4): 399-406. Portuguese
5. Paiva WO. Taxa de polinização cruzada em cubiu. *Pesquisa Agropecuária Brasileira*. 1999; 34(1):145-49. Portuguese
DOI: <http://dx.doi.org/10.1590/S0100-204X1999000100020>.
6. Pizzinato JR, Shuelter AR, Junior ATA, Rocha ACS, Silva JM, Volkweis CR, et al. Crossing and diagnostic methods of cubiu hybrid plants based on genetic markers. *Crop Breeding and Applied Biotechnology*. 2008; 8(4): 283-90.
DOI:10.12702/1984-7033.v08n04a05
7. Pires AMB, Silva PS, Nardelli PM, Gomes JC, Ramos AM. Caracterização e processamento de cubiu (*Solanum sessiliflorum*). *Revista Ceres*. 2006; 53(307):309-16. Portuguese
8. Yuyama LKO, Macedo SHM, Aguiar JPL, Filho DS, Yuyama K, Fávaro DIT, et al. Quantificação de macro e micro nutrientes em algumas etnovarietades de cubiu (*Solanum sessiliflorum* Dunal). *Acta Amazônica*. 2007; 37(3): 425-430.
DOI <http://dx.doi.org/10.1590/S0044-59672007000300014>
9. Bohs L. A chloroplast DNA phylogeny of *Solanum* section *Lasiocarpa*. *Systematic Botany*. 2004;29(1):177-187.
DOI: <https://doi.org/10.1600/036364404772974>
10. Chartzoulakis K, Drosos N. Water use and yield of greenhouse grown eggplant under drip irrigation. *Agricultural Water Management*. 1995;28(2):113-120.
DOI: [https://doi.org/10.1016/0378-3774\(95\)01173-G](https://doi.org/10.1016/0378-3774(95)01173-G)
11. Gao QH, Yu JG, Wu CS, Wang ZS, Wang YK, Zhu DL, et al. Comparison of drip, pipe and surge spring root irrigation for Jujube (*Ziziphus jujuba* Mill.) fruit quality in the Loess Plateau of China. *Plos One*. 2014;9(2):e88912.
DOI: <https://doi.org/10.1371/journal.pone.0088912>
12. He H, Ma F, Yang R, Chen L, Jia B, Cui J, et al. Rice performance and water use efficiency under plastic mulching with drip irrigation. *Plos One*. 2013;8(12):e83103.
DOI: <https://doi.org/10.1371/journal.pone.0083103>
13. Aujla MS, Thind HS, Buttar GS. Fruit yield and water use efficiency of eggplant (*Solanum melongena* L.) as influenced by different quantities of nitrogen and water applied through drip and furrow irrigation. *Scientia Horticulturae*. 2007; 112(2):142-48.
DOI: 10.1016/j.scienta.2006.12.020
14. Leogrande R, Lopodota O, Vitti C, Ventrella D, Montemurro F. Effects of irrigation volumes and organic fertilizers on eggplant grown in Mediterranean environment. *Acta Agriculturae Scandinavica, Section B – Soil and Plant Science*. 2014; 64(6):518-28.
DOI: 10.1080/09064710.2014.927526

- 218 15. Alvares CA, Stape JL, Sentelhas PC, de Moraes JLG, Sparovek G. Koppen's climate classification
219 map for Brazil. *Meteorologische Zeitschrift*. 2013; 22(6): 711-28.
220 DOI: 10.11.27/0941-2948/2013/0507
- 221 16. Marouelli WA, Silva WLDC, Silva HRD. Irrigação por aspersão em hortaliças. Brasília DF: Embrapa
222 Informação Tecnológica; 2008. Portuguese
- 223 17. Argerich CA, Poggi LM, Lipinski VM. The influence of poultry manure and irrigation during fruit-setting
224 critical period on processing tomatoes. *Acta Horticulturae*. 1999; 487(1):557-62.
225 DOI: 10.17660/ActaHortic.1999.487.91
- 226 18. Guang-Cheng S, Na L, Zhan-Yu Z, Shuang-en Y, Chang-ren C. Growth, yield and water use efficiency
227 response of greenhouse-grown hot pepper under Time-Space deficit irrigation. *Scientia Horticulturae*.
228 2010; 126 (2):172-79.
229 DOI:10.1016/j.scienta.2010.07.003
- 230 19. Bletsos FA, Thanassouloupoulos CC, Roupakias DG. Water stress and verticillium wilt severity on
231 eggplant (*Solanum melongena* L.). *Journal of Phytopathology*. 1999; 147(4): 243-48.
232 DOI: <https://doi.org/10.1046/j.1439.1999.147004243.x>
- 233 20. Silva Filho DF. Cocona (*Solanum sessiliflorum* Dunal): Cultivo y utilización. Caracas: Tratado de
234 Cooperación Amazonica; 1998. Spanish
- 235 21. Marouelli WA, Silva WLC. Irrigação por gotejamento do tomateiro industrial durante o estágio de
236 frutificação, na região de Cerrado. *Horticultura Brasileira*. 2006; 24(3): 342-46. Portuguese
237 DOI: <http://dx.doi.org/10.1590/S0102-05362006000300014>
- 238 22. García A, Barreira J, Vargas G, Melgarejo LM, Hernandez MS, Quintero L, et al. Ecofisiología y
239 respuestas al ambiente de producción de cocona (*S. sessiliflorum*) en la Amazonia norte, in: García
240 JAB, Gómez MSH, Melgarejo LM, (Eds.), *Estudios Ecofisiológicos en la Amazonia Colombiana*. 2.
241 Cocona. Instituto Amazônico de Investigaciones Científicas-Sinchi, 2001. 77-92.
242