

1                   **DISTRIBUTION OF ECOSYSTEM HEALTH INDICATORS FOR**  
2                   **BIOMONITORING OF OIL POLLUTION IN THE WESTERN NIGER DELTA,**  
3                   **NIGERIA**

4  
5    **ABSTRACT**

6    As a result of the limitations of physical and chemical methods for monitoring pollution,  
7    interest on the more reliable biological monitoring intensified over the past four decades.  
8    Soil microarthropods, specifically the free-living mites (Cryptostigmata, Mesostigmata,  
9    Prostigmata) and Collembolans were used as monitor (ability to withstand pollutants) and  
10   indicator (sensitive to pollutants) species in the Eastern Niger Delta. Study was undertaken  
11   in the Western Niger Delta (Delta State) across three eco-vegetational zones (freshwater  
12   swamp forest, Mangrove swamp forest, Lowland rainforest) in the area to determine if these  
13   ecosystem health indicators were widely distributed in these zones. Collections were made  
14   during the rainy season over a 4-month period. A modified Berlese-Tullgren funnel was  
15   used for extraction of microarthropods. Free-living mites: Cryptostigmata (Oribatida) –  
16   *Archogozettes magnus*, *Opiida sp.*, *Annecticarus sp.*, *Bicyrthermania negeriana*, *Cephalida*  
17   *sp.*, *Schelorbates sp.*, *Galumnida sp.*, Mesostigmata (Gamasida) - *Asca sp.*, *Trichuropodida*  
18   and Collembolan – *Paronella sp.*, were widely distributed across the eco-vegetational  
19   zones. **Oribatids were most abundant across eco-vegetational zones.** These mesofauna  
20   contained the full complement of monitor and indicator species. It is therefore possible to  
21   use these mesofauna for biomonitoring of oil pollution across the Niger Delta (eastern and  
22   western sectors), Nigeria.

23   Key Words: Pollutants, Ecosystem Health Indicators, Mites, Collembolans, Biomonitoring,  
24   Niger Delta, Nigeria.

25   **INTRODUCTION**

26   The effective dose of a pollutant in an individual, determined by physical or chemical  
27   methods may be much lower than the result obtained. The total concentration may be raised  
28   by high levels of surface contamination or the binding of pollutant to inert sites. Thus the  
29   biological significance of the concentration in the individual may be overestimated. In  
30   contrast, biological monitoring assesses the significance of a pollutant for an organism in its  
31   habitat and other individuals in the community. Monitor and indicator species are used to

32 measure pollutant impact (Martin and Coughtrey, 1982). Monitor species have the ability to  
33 withstand pollutants and they are used to assess the scale and distribution of the pollutant.  
34 In contrast, indicator species are sensitive to the pollutant and their presence or absence  
35 indicates a significant level of contamination.

36 Mites of the suborder Oribatida, also called “beetle” or “moss” mites are the world’s most  
37 numerous arthropods living in the soil (Seastedt, 1984). There are several thousand species,  
38 yet the fauna of much of the tropics is still unknown. They are the most abundant group of  
39 microarthropods in most forested, grassland and desert ecosystem (Seastedt, 1984). Mites  
40 and other microarthropods (Including Collembolans), part of the mesofauna play a crucial  
41 role in the context of soil biodiversity, decomposition and mineralization processes  
42 (Seastedt, 1984; Tiann *et al.*, 1998). Among the microarthropods, the feeders (mycophages)  
43 are dominant Collembolans free-living astigmatid mites and most oribatids  
44 (Cryptostigmmites) have well-developed mouthparts, capable of fragmenting organic matter,  
45 while feeding on the microflora adhering to detritus. Fragmentation and communitation are  
46 important to the decomposition and mineralization processes by creating new surface area  
47 for microbial colonization (Fountain and Hopkin, 2005). The decomposer community  
48 received greater interest within soil ecology in the past six decades (Bardgett, 2002). The  
49 free-living mites (Cryptostigmata, Mesostigmata, Prostigmata) and Collembolans have been  
50 used as monitor and indicator species to determine ecosystem health in Eastern Niger-Delta:  
51 Rivers (Gbarakoro *et al.*, 2010; Okiwelu *et al.*, 2011a & b; Gbarakoro *et al.*, 2011;  
52 Gbarakoro and Anianu, 2016; Gbarakoro and Umoren, 2016) and Bayelsa (Gbarakoro *et al.*,  
53 2017) States. They have also been used for biomonitoring in spent automobile lubricant  
54 and heavy metal-polluted sites (Gbarakoro *et al.*, 2015; Gbarakoro and Chukumali, 2016).

55 This study was undertaken to produce baseline data on the species composition of mites and  
56 Collembolans in relatively undisturbed habitats in mangrove swamp forest, freshwater  
57 swamp forest, lowland rainforest in Delta State, Western Niger Delta, Nigeria. If they are  
58 distributed across these zones, they can be used for biomonitoring of oil pollution across the  
59 Niger Delta.

## 60 MATERIALS AND METHODS

61 The mangrove swamp forest is located on the bank of the Isaba River, Warri North Local  
62 Government Area (LGA) and the freshwater swamp forest, at Merogun, Warri South LGA,  
63 Delta State. Collections were made from two sites in each of in each of the LGAs. At Isaba

64 in Warri North LGA, collections were made from the mangrove forest and in lowland  
65 rainforest approximately 200m away. In Warri South LGA, at Merogun, collections were  
66 from the freshwater swamp forest and approximately 200km away, in farm bush located in  
67 lowland rainforest.

68 Studies were conducted over a 4-month period, June-September, during the rainy season. At  
69 each location, an area 30.00cmx30.00cm was delineated. Each delineated area was divided  
70 into 4 sub-plots and collections made monthly in rotation from each sub-plot. Collections  
71 were made at 08.00-09.00hrs from litter and depths of 0.50cm, 5.0-10.0cm, 10.0-15.0cm;  
72 15.0-20.0 cm, 20-25cm and 25-30cm. Samples were placed in labeled transparent bags. A  
73 modified Berlese-Tullgren funnel was used to extract the microarthropod species were  
74 identified to family levels by keys and illustrations provided by Badejo (1994) and type  
75 specimens in the Entomology and Pest Management Laboratory, University of Port  
76 Harcourt.

## 77 RESULTS

78 In the mangrove swamp forest, six species of Cryptostigmata (Oribatida) were collected:  
79 *Archogozettes magnus*, *Annecticarus sp.*, *Bicyrthermania negeriana*, *Cephalida sp.*,  
80 *Scheloribates sp.*, *Galumnida sp.*; a species of Mesostigmata (*Gamasida*) *Asca sp.* and a  
81 species of Collembola, *Paronella sp.* (Table 1). In the freshwater swamp forest, 4  
82 Cryptostigmata (*Oribatida*) species - *Opiida sp.*, *Galumnida sp.*, *Cephalid sp.*,  
83 *Scheloribates*, 1 Mesostigmata (*Gamasida sp.*) were collected (Table 2).

84 In lowland rain forest - Warri North LGA (1), 4 species of Cryptostigmata (*Oribatida*) -  
85 *Archeozettes magnus*, *Galumnida sp.*, *Cephalid sp.*, *Scheloribates sp.*; 2 species of  
86 Mesostigmata (*Gamasida*), *Asca sp.* and *Uropodida sp.* and 1 species of Collembola -  
87 *Paronella sp.* were collected (Table 3). In lowland rainforest of Warri South LGA, 4  
88 Cryptostigmata (*Oribatida*) - 4 species *Scheloribates sp.*, *Galumnida sp.*, *Cephalida sp.*,  
89 *Opiida sp.*; 2 Mesostigmata (*Gamasida*) - *Asca sp.*, *Uropodida sp.*, and 1 Collembola *sp.*-  
90 *Paronella sp.* were collected (Table 3). Oribatids were most abundant across eco-  
91 vegetational zones. In correlation and paired t-test statistics, among Oribatids/Gamasids,  
92 Oribatids/Collembola and Gamasids/Collembola, correlations were 0.970, 0.962 and 1.000  
93 respectively, but only significant between Gamasids/Collembola (0.19; t=4.914; p<0.01)  
94 Lowland rainforest yielded the highest number of soil microarthropods. In correlation and  
95 paired t-test statistics, Mangrove Forest/Freshwater Swamp forest, Mangrove

96 forest/Lowland Rainforest and Freshwater Swamp forest/Lowland Rainforest, correlations  
97 were 0.985, 1.000 and 0.983 respectively, but only significant between Mangrove  
98 forest/Lowland Rainforest (0.007; t-test=-985; p<0.01) (Table 4).

## 99 DISCUSSION

100 In the lowland rainforest, the species composition of mites and Collembolans was more  
101 limited than that of Okiwelu *et al.* (2011) from undisturbed habitat in lowland rainforest of  
102 eastern Niger Delta. This was probably due to the significantly reduced period for  
103 collection. The absence of any Prostigmata sp. was also probably due to the limited  
104 collection period. However, both monitor species of Cryptostigmata (*Galumnida sp.*;  
105 *Scheloribates sp.*) and indicator species of Cryptostigmata (*Cephalides sp.*, *Archeogozettes*  
106 *magnus*, *Oppia sp.*) and Mesostigmata (*Asca sp.*, *Trachyllropodida sp.*, *Uropodida sp.*)  
107 were encountered. The Collembolan monitor sp. (*Paronella sp.*) was encountered.

108 In the mangrove swamp forest, there were six spp. that consisted of both monitor spp.  
109 (*Bichrthermannia nigeriana*, *Scheloribates sp.*, *Galumnida sp.*) and indicator spp.  
110 Cryptostigmata - (*Archeogozettes magnus.*, *Cephalida sp.*) and Mesostigmata - (*Asca sp.*). The  
111 Collembolan monitor species - *Paronella sp.* was also identified. In the freshwater swamp  
112 forest, there were 2 monitor spp. - Mesostigmata (*Galumnida sp.*, *Scheloribates sp.*) and 3  
113 indicator species – (*Oppia sp.*, *Cephalida sp.*) and Mesostigmata (*Asca sp.*). In lowland  
114 rainforest, 2 monitor species- Mesostigmata (*Galumnida sp.*, *Scheloribates sp.*) and 1  
115 indicator species - Mesostigmata (*Asca sp.*) were identified. The Collembolan monitor  
116 species - *Paronella* was also identified.

117 In a series of studies in lowland rainforest, Rivers State, eastern Niger Delta (Gbarakoro *et*  
118 *al.*, 2010, 2011; Okiwelu *et al.*, 2011 a & b), it was established that a full complement of  
119 soil microarthropods of monitor and indicator species was adequate for bio-monitoring to  
120 assess ecosystem health. Monitor species are used to assess the scale and distribution of the  
121 pollutants while indicator species are sensitive to the pollutant and their presence or absence  
122 indicates a significant level of contamination. The mangrove, freshwater swamp forests and  
123 lowland rainforests of the western Niger Delta also have these complements of soil  
124 microarthropods. Oil pollution biomonitoring with soil microarthropods is therefore feasible  
125 across all eco-vegetational zones in the Niger Delta and it is thus recommended.

## 126 CONCLUSION

127 The full complement of soil micro-arthropod ecosystem health indicators (monitor and  
 128 indicator species) were widely distributed across the major eco-vegetational zones  
 129 (Lowland rainforest, Mangrove swamp forest, Freshwater Swamp forest) of the western  
 130 Niger Delta. They had been found extensively distributed in the eastern Niger Delta.  
 131 Consequently, their use for biomonitoring of oil pollution in the Niger Delta is feasible and  
 132 advisable.

133  
 134  
 135  
 136  
 137

Table 1. Species of microarthropods from Mangrove Swamp Forest

Vegetation type	Mites		Collembola
	Oribatida	Gamasida	
Mangrove Swamp Forest	<i>Archezogettes magnus</i> <i>Annecticarus sp.</i> <i>Bicyrthermannia nigeriana</i> <i>Cephalida sp.</i> <i>Galumnida sp.</i> <i>Scheloribates sp.</i>	<i>Asca sp.</i>	<i>Paronella sp.</i>

138  
 139  
 140  
 141

Table 2. Species of microarthropods from Freshwater Swamp Forest

Habitat type	Mites		Collembola
	Oribatida	Gamasida	
Freshwater Swamp Forest	<i>Cephalida sp.</i> <i>Galumnida sp.</i> <i>Oppia sp.</i> <i>Scheloribates sp.</i>	<i>Ascidae sp.</i>	

142  
 143  
 144  
 145

Table 3. Species of microarthropods from Lowland Rainforest

Vegetation type	Mites		Collembola
	Oribatida	Gamasida	
Lowland Rainforest - Warri North LGA	<i>Archezogettes magnus</i> <i>Cephalida sp.</i> <i>Galumnida sp.</i> <i>Scheloribates sp.</i>	<i>Asca sp.</i>	<i>Paronella sp.</i>
Lowland Rainforest (Farm bush) - Warri South LGA	<i>Cephalida sp.</i> <i>Galumnida sp.</i> <i>Oppia sp.</i> <i>Scheloribates sp.</i>	<i>Asca sp.</i> <i>Uropodidae sp.</i>	<i>Paronella sp.</i>

146

147  
148  
149

Table 4. Mite and Collembolan abundance across eco-vegetational zones

Order	Sub-Order	Mangrove forest	Freshwater swamp forest	Lowland rainforest
Acarina	Oribatida	42	11	77
	Gamasida	6	3	15
Collembolan		2	0	9

150 REFERENCES

- 151 Badejo M.A. 1996. Measuring the diversity of soil microarthropods and microfauna in an  
152 area of conservation of biodiversity. *In* Biosphere resources for diversity conservation and  
153 sustainable development in Anglophone Africa (BRAAF), Assessment and monitoring  
154 techniques in Nigeria, Abeokuta, Nigeria, 120pp.
- 155 Bardgett R.D. 2002. Causes and consequences of biological diversity in soil. *Zoology* 105:  
156 367-374.
- 157 Fountain M.T., Hopkins S.P. 2005. *Folsornia candida* (Collembolan): a standard soil  
158 arthropod. *Annual Review of Entomology* 50:201-222.
- 159 Gbarakoro T.N., Anianu G.O. 2016. Contributions of Soil Microarthropods to Ecosystem  
160 Recovery at Tai communities, Rivers State, Nigeria. *Resources and Environment* 6: 136-  
161 142.
- 162 Gbarakoro T.N., Chukumati J. 2016. Impact of spent Mushroom substrate on soil  
163 microarthropods in spent automobile lubricant habitat types at University farm, University  
164 of Port Harcourt, Rivers State, Nigeria. *International Journal of Environment and Pollution*  
165 *Research* 4: 13-23.
- 166 Gbarakoro T.N., Umoren E.I. 2016. Soil microarthropod- Indicators of effects of living  
167 Mulches in Sole and Mixed Cropping Systems at University Park Farm, Port Harcourt,  
168 Nigeria. *European Journal of Agriculture and Forestry Research* 4:13-21.



- 169 Gbarakoro T.N., Ogele E.J., Umeozor O.C. 2017. Impact of tillage remediation on soil  
170 microarthropods and nutrients in hydrocarbon contaminated soil ecosystem at Ikarama  
171 community, Bayelsa State, Nigeria. *Nigerian Journal of Entomology* 33:111-120.
- 172 Gbarakoro T.N., Okiwelu S.N., Badejo M.A., Umeozor O.C. 2010. Soil microarthropods in  
173 a secondary rainforest in Rivers State, Nigeria -I- Seasonal variations in species richness,  
174 vertical distribution and density in an undisturbed habitat. *Scientia Africana* 9:48-54.
- 175 Gbarakoro T.N., Okiwelu S.N., Umeozor O.C., Badejo M.A. 2011. Soil microarthropods in  
176 a secondary rainforest -III- partial recovery after an oil spill. *International Journal of*  
177 *Ecosystem* 1: 1-4.
- 178 Gbarakoro T.N., Oneche E., Umeozor C.O. 2015. Impact of Heavy metals on species  
179 richness, abundance and vertical distribution of soil microarthropods in waste dump sites at  
180 Delta Steel Company, Warri, Nigeria. *International Journal of Environment and Resource* 4:  
181 9-14.
- 182 Martin M.H., Coughtrey P.J. 1982. Biological monitoring of heavy metal pollution.  
183 Pollution monitoring series Ad. (Mellanby K., ed.). Applied Sciences Publishers Ltd.  
184 London 475pp.
- 185 Okiwelu S.N., Gbarakoro T.N., Badejo M.A. 2011a. Soil microarthropods in a secondary  
186 rainforest, Rivers State: Ecosystem health indicators of oil pollution. *Journal of Ecology*  
187 *and Natural Environment* 3:29-32.
- 188 Okiwelu S.N., Gbarakoro T.N., Umeozor O.C., Badejo M.A. 2011b. Soil microarthropods  
189 in a secondary rainforest, Rivers State, Nigeria -IV- the impact of oil pollution on their  
190 distribution. *Resources and Environment* 1:1-4
- 191 Seastedt T.R. 1984. The role of microarthropods in decomposition and mineralization  
192 processes. *Annual Review of Entomology* 29:25-46.
- 193 Tian G., Adejuyigbe C.O., Adeoye G.O., Kang B.T. 1998. Role of soil microarthropods in  
194 leaf decomposition and N release under various land-use practices in the humid tropics.  
195 *Pedobiologia Jena* 42:33-42.